



Effects of Substitution of Fish Meal (FM) and Macroalgae (MA) with Soybean Meal and Rice Bran in a Commercial Juvenile Abalone (*Haliotis duscus hannai*) Diet on Growth Performance

Ki Wook Lee¹, Hee Sung Kim¹, Dong Gyu Choi¹, Bok Il Jang¹, Hyeon Jong Kim¹, Ahyeong Yun¹, Sung Hwoan Cho^{1*}, Byeong-Hee Min², Kyoung-Duk Kim³, Hyon-Sob Han³

¹Korea Maritime and Ocean University, Department of Convergence Study on the Ocean Science and Technology, Busan 49112, Korea.

²Busan Marine Fisheries Resources Research Institute, Busan 46763, Korea.

³Aquafeed Research Center, National Fisheries Research and Development Institute, Gyeongsangbuk-do 37517, Korea.

* Corresponding Author: Tel.: +82.51 4104755; Fax: +82.51 4044750;
E-mail: chosunh@kmou.ac.kr

Received 28 June 2016
Accepted 22 November 2016

Abstract

The effects of fish meal (FM) and macroalgae (MA) substitution with fermented soybean meal and rice bran were tested on a commercial diet in juvenile abalone. Abalone (21600) individuals were distributed equally to four diet treatments. Four experimental diets were prepared: a Std diet, FM50, FM50+MA50 and FM50+MA100. The standard diet (Std) consists of FM (140 g/kg), fermented soybean meal (250 g/kg), corn gluten meal (34 g/kg), shrimp meal (30 g/kg), wheat flour (200 g/kg), dextrin (50 g/kg) and MA (250 g/kg). *Undaria* and *Laminaria* were also prepared. The feeding trial lasted for 16 weeks. Survival of abalone fed the formulated diets was higher than that of abalone fed the *Undaria* and *Laminaria*. Weight gain and specific growth rate of abalone fed the Std diet were higher than all other diets. Substitutability of the fermented soybean meal for FM seemed to be rather limited in the commercial diet for abalone farms. However, another 500 g/kg MA in the commercial diet could be substituted with rice bran as long as 500 g/kg MA were substituted with rice bran. The FM50+MA50 diet is the most recommendable for abalone farm in considering feed cost.

Keywords: Abalone (*Haliotis discus hannai*), fish meal (FM), macroalgae (MA), fermented soybean meal, rice bran.

Introduction

Annual abalone aquaculture production in Korea was about 20 metric tons in 2000 and reached 8,982 metric tons in 2014 (FishStatJ, 2015). This trend is likely to continue due to its continuing high consumer demand and continued farm expansions to meet the demand.

Fish meal (FM), which has been commonly used as a primary protein source in aquafeeds, prices have increased 285% since 2001 (FAO, 2014) and are projected to continue increasing due to a shortage of catchable FM sources in wild and continuing expansion of global fish meal fed aquaculture. Macroalgae (MA), which has been used as a replacement or feed additive, prices are also becoming more costly due to expansion of the biofuel industry to develop MA for ethanol extraction.

In Korea, abalone farmers are likely to continue to feed abalone on MA such as *Undaria pinnatifida* Harvey or *Laminaria japonica* Areschoug during the winter season on their farms as these MA are grown naturally and harvested in the wild. Although its nutrient content, such as protein (amino acid) and lipid (fatty acid) do not satisfy the complete dietary

requirement for abalone, *Haliotis discus hannai* Ino (Uki, Kemuyama, & Watanabe, 1986a; Mai, Mercer, & Donlon, 1995a, b), farmers prefer feeding abalone on MA for the convenience of farm management. This may result in poorer growth rates and increase the production cost of abalone over commercial feeds. Generally during summer, the dry or salted MA is commonly fed to abalone. However, the nutrients in the dry or salted MA are easily destroyed in the drying or salting process and this can also result in poorer growth rates.

One of the largest production costs in aquaculture operations is feed and a compromise needs to be made between a nutritionally balanced and cost effective feed (Neori & Nobre, 2012). As FM and MA are the largest components of commercial abalone feeds and prices continue to increase, alternatively sources need to be identified and their performances evaluated against existing commercial feeds. Globally research has shown that a well formulated feed will produce better growth rates than a single algal species diet (Lee, 1998; Cho, Park, Kim, Yoo, & Lee, 2006; Kim, Lee, Han, Kim, & Park, 1998; Cho, Park, Kim, & Yoo, 2008; Naidoo, Maneveldt, Ruck, & Bolton, 2006; Gracia-Esquivel &

Felbeck, 2009; Dang, Li, Speck, & Benkendorff, 2011; Kim *et al.*, 2015; Myung *et al.*, 2016).

Studies by Uki, Kemuyama, & Watanabe (1985a, b, 1986a) have shown that plant protein (soybean and cottonseed meal) are good alternative sources for casein, and can produce the best weight gain of abalone, *H. discus hannai*, compared to other plant proteins tested (Lee, Yun, & Hur, 1998). Cho *et al.* (2008) also reported that growth of abalone fed a combined fish and soybean meal diet or a combined fish, soybean and crustacean meal diet was comparable to that of abalone fed a casein-basal diet. In addition, FM at 350 g/kg could be completely replaced with soybean meal at 580 g/kg with an additional 5 g/kg methionine supplementation, as it is lacking in the plant protein source (Cho, 2010). Therefore, soybean meal seems to be a promising alternative protein source for FM in abalone feed.

Fermentation of soybean meal increases protein content, eliminates trypsin inhibitors, and reduces peptide size (Hong, Lee, & Kim, 2004; Kook, Cho, Hong, & Park, 2014). Dietary inclusion of fermented plant protein sources, such as soybean and cottonseed meals also produced a promising effect on performance of Nile tilapia, *Oreochromis niloticus* Linne and black sea bream, *Acanthopagrus schlegelii* Bleeker (Lim & Lee, 2011; Zhou *et al.*, 2011). A study by Kim *et al.* (2015) on abalone, *H. discus* Reeve revealed that 1000 g/kg substitution of *L. japonica* with rice bran, [an agriculture byproduct and rich in the nutrients such as crude protein and vitamins (Gao, Kaneko, Hirata, Toorisaka, & Hano, 2008)], at 200 g/kg in the diet was successful in terms of abalone weight gain. However, the work needed to be trialed under commercial farm conditions.

The aim of this paper is to investigate the substitution effect of FM and MA with fermented soybean meal and rice bran in commercial abalone diet on growth performance of juvenile abalone at commercial farm.

Materials and Methods

Preparation of Abalone and Rearing Conditions

Juvenile abalone, *H. discus hannai*, were purchased from a private hatchery and transferred to Joeun abalone farm (Wando, Jeollanamdo, Korea). Abalone were acclimated for two weeks and fed with the dry *L. japonica* once a day at the ratio of 2-3% of total biomass. A 6 ton concrete flow-through raceway tank (120 cm×690 cm×50 cm; water volume: 4.14 ton) at a flow rate of 42.3 L/min with sand filtered seawater (temperature range: 17.9 to 22.8°C (mean±SD: 20.2±0.01°C) was divided into 3 sections (120 cm×230 cm×50 cm; water volume: 1.38 ton) by a plastic-framed net (mesh size of 5 mm), and 4 concrete tanks were used for this experiment. Juvenile abalone (n = 21600) individual averaging 3.6 g were randomly distributed into each section (n=1800 per

section). Aeration was supplied into each raceway and the photoperiod followed natural conditions. The experimental diets were fed to abalone once a day (17:00) at a satiation level with a little leftover (about 2-3% of biomass). Dead abalone were removed daily and the tanks were cleaned twice a week. The feeding trial lasted for 16 weeks. At the end of the feeding trial, 200 abalone were randomly sampled from each treatment and collectively weighed.

Preparation of the Experimental Diets

Four experimental diets and the dry *Undaria* and *L. japonica* were prepared (Table 1). Four diets are referred to as follows: a Std, FM50, FM50+MA50 and FM50+MA100 diets. The standard diet (Std) is a commercial juvenile abalone diet and consists of FM and shrimp meal, fermented soybean, what flour and MA (250 g/kg, a mixture of *Undaria* and *Hizikia fusiforme* Harvey at a ratio of 1:1) as the protein and carbohydrate sources and was formulated to satisfy dietary nutrient requirements for (Uki *et al.*, 1986a; Mai *et al.*, 1995a, b). The 500 g/kg FM (70 g/kg of diet), and combined 500 g/kg FM and 500 g/kg MA (125 g/kg of diet), and combined 500 g/kg FM and 1000 g/kg MA (250 g/kg of diet) were substituted with the same amount of fermented soybean meal, and combined fermented soybean meal and rice bran, respectively, in the Std diet, referred to as FM50, FM50+MA50 and FM50+MA100 diets. Four diets were pelletized by an extruded pelletter (Jyoda, Japan) in Ewha Oil and Fat Industry Co. Ltd. (Busan, Korea). Pellets were round-shape, and their sizes were 5 mm in diameter (1.5 mm in thickness). *Undaria* and *L. japonica* were prepared to compare effect of the experimental diets with that of MA on the growth performance of abalone.

Analytical Procedures of the Diets and Carcass

Abalone from each treatment were sampled and frozen for chemical analysis (100 at the beginning and 50 at the trial termination). Prior to examination, all samples were slightly thawed, followed by separation of the shell and soft-body tissue. Shell length and width were measured to 1.0 mm using a digital caliper (Mitutoyo Corporation, Kawasaki, Japan), and the ratio of soft body weight to body weight (the soft body weight+the excised shell's weight) was calculated to determine a condition index for abalone. Specific growth rate (SGR, %/body weight gain day) was calculated using the formula:

$$SGR = [(\ln(W_f) - \ln(W_i))/\text{days of feeding}] \times 100,$$

where $\ln(W_f)$ = natural log of the final mean weight of abalone and $\ln(W_i)$ = natural log of the initial mean weight of abalone Britz (1996).

The pooled separated soft body tissue from the sampled abalone from each diet was then

Table 1. Feed formulation of the experimental diets (g/kg, DM basis)

	Experimental diets					
	Std	FM50	FM50+MA50	FM50+MA100	<i>U. pinnatifida</i>	<i>L. japonica</i>
Fish meal (FM) ¹	140	70	70	70		
Fermented soybean meal ²	250	320	320	320		
Corn gluten meal	34	34	34	34		
Shrimp meal	30	30	30	30		
Wheat flour	200	200	200	200		
Dextrin	50	50	50	50		
Macroalgae (MA) ³	250	250	125			
Rice bran			125	250		
<i>Spirulina</i>	5	5	5	5		
Yeast	10	10	10	10		
Sea aroma	1	1	1	1		
Choline chloride(500g/kg)	5	5	5	5		
Vitamin premix ⁴	5	5	5	5		
Mineral premix ⁵	20	20	20	20		
<i>Nutrients (g/kg)</i>						
Dry matter	987	988	978	981	860	887
Crude protein	363	361	380	383	207	90
Crude lipid	14	14	37	68	8	4
Carbohydrate ⁶	424	435	464	467	424	581
Ash	199	190	119	82	361	325

¹Fish meal (FM): Alaska pollack meal.

²Fermented soybean meal was purchased from CJ CheilJedang Corp. (Seoul, Korea).

³Macroalgae (MA) were the mixture of *Undaria* and *Hizikia fusiforme* at a ratio of 1:1.

⁴Vitamin premix contained the following amount which were diluted in cellulose (g/kg mix): excipient, 317; riboflavin, 23.8; pyridoxine, 4.7; niacin, 95.2; Ca-pantothenate, 33.3; inositol, 476.9; folic acid, 1.5; p-amino benzoic acid, 47.6.

⁵Mineral premix contained the following ingredients (g/kg mix): Excipient, 45.5; MgSO₄, 140.8; NaH₂PO₄, 92.4; KH₂PO₄, 246; Ca(H₂PO₄)₂, 139.5; ZnSO₄, 22.5; Ca-lactate, 310; AlCl₃, 0.15; KI, 0.15; MnSO₄, 2; CoCl₂, 1.

⁶Carbohydrate was calculated as the difference between 1000 and the sum of crude protein, crude lipid and ash contents.

homogenized and used for proximate analysis. Crude protein content was determined by the Kjeldahl method (Auto Kjeldahl System, Buchi B-324/435/412, Switzerland), crude lipid was determined using an ether-extraction method, moisture was determined by oven drying at 105°C for 24 h and ash was determined using a muffle furnace at 550°C for 4 h. All methods were according to AOAC (1990) practices. Amino acid composition of the experimental diets were determined by using a high speed amino acid analyzer (Hitachi L-8800, Tokyo, Japan) after which the samples were hydrolyzed in 6 N HCl for 24 h at 110°C.

Statistical Analysis

One-way ANOVA and Duncan's multiple range test (Duncan, 1955) were used to determine the significance of the differences among the means of treatments by using SAS version 9.3 program (SAS Institute, Cary, NC, USA). Percentage data was arcsine-transformed prior to statistical analysis.

Results

Amino acid profiles of the experimental diets are given in Table 2. The essential amino acids, such as arginine, isoleucine, lysine, methionine, threonine and valine were relatively low in the FM50 diet, but

arginine, histidine, isoleucine, leucine, phenylalanine and valine were relatively high in the FM50+MA50 and FM50+MA 100 diets compared to the Std diet.

Fatty acid profiles of the experimental diets are presented in Table 3. A sum of n-3 highly unsaturated fatty acid (HUFA) content were relatively high in the Std diet compared to that of the FM50, FM50+MA50 and FM50+MA100 diets, especially linolenic acid (18:3n-3), eicosapentaenoic acid (EPA, 20:5n-3) and docosahexanoic acid (DHA, 22:6n-3) decreased in proportion to an amount of substitution of FM and MA with the fermented soybean meal and rice bran, respectively, but linoleic acid (18:2n-6) content increased. No EPA and DHA was also found in MA (*Undaria* and *L. japonica*) diets.

Survival of abalone fed the all formulated diets was significantly higher (P<0.0001) than that of abalone fed the *Undaria* and *L. japonica* (Table 4). Survival of abalone fed the *Undaria* was significantly higher (P<0.05) than that of abalone fed the *L. japonica*. Weight gain (g/abalone) and SGR (%/day) of abalone fed the Std diet were significantly higher (P<0.0001) than those of abalone fed the all other diets, and followed by the FM50, FM50+MA50 and FM50+MA100 diets, *Undaria* and *L. japonica* in that in order.

The longest shell length was obtained in abalone fed the Std diet, followed by the FM50, FM50+MA50 and FM50+MA100 diets, *L. japonica*, and *Undaria* (Table 5). The widest shell width was observed in

Table 2. Amino acid profiles of the experimental diets (g/kg, DM basis)

	Experimental diets					
	Std	FM50	FM50+MA50	FM50+MA100	<i>U. pinnatifida</i>	<i>L. japonica</i>
Alanine	19.0	18.1	19.0	18.7	19.3	4.7
Arginine	21.3	20.5	22.2	22.9	7.3	2.6
Aspartic acid	34.0	34.4	36.4	35.8	14.0	15.6
Cystine	4.5	4.6	5.0	5.1	1.4	1.0
Glutamic acid	59.5	60.2	64.3	64.1	19.8	25.6
Glycine	19.5	17.3	18.3	18.0	8.2	3.4
Histidine	7.4	7.5	8.3	8.5	2.2	1.0
Isoleucine	14.4	14.2	15.7	15.5	6.4	2.4
Leucine	26.9	26.9	28.9	28.5	11.1	4.1
Lysine	18.1	16.9	18.2	18.0	7.2	2.8
Methionine	5.7	5.3	5.6	6.1	2.8	1.1
Phenylalanine	16.1	16.3	17.5	17.2	3.7	2.9
Proline	18.4	18.1	19.2	19.6	6.5	3.1
Serine	16.8	16.9	17.0	16.8	6.0	2.8
Threonine	14.2	14.0	14.3	14.0	6.4	3.0
Tyrosine	10.6	10.5	11.3	11.7	4.2	1.8
Valine	16.2	15.9	17.6	17.5	7.8	3.3

Table 3. Fatty acid compositions of the experimental diets (g/kg of total fatty acid)

	Experimental diets					
	Std	FM50	FM50+MA50	FM50+MA100	<i>U. pinnatifida</i>	<i>L. japonica</i>
14:0	36.1	32.1	16.2	10.0	55.9	88.4
15:0	n.d.	n.d.	n.d.	n.d.	9.6	n.d.
16:0	192.0	194.8	174.2	172.9	273.6	155.0
18:0	25.1	25.4	19.8	18.6	27.0	13.2
20:0	n.d.	n.d.	4.1	4.6	6.3	n.d.
21:0	n.d.	n.d.	n.d.	n.d.	5.8	n.d.
22:0	n.d.	n.d.	2.8	2.9	n.d.	n.d.
24:0	n.d.	n.d.	4.5	4.9	n.d.	n.d.
∑Saturates	253.2	252.3	221.6	213.9	378.2	256.6
14:1n-9	2.9	3.2	n.d.	n.d.	7.1	5.7
16:1n-9	35.6	27.4	13.3	n.d.	8.1	27.5
16:2n-6	n.d.	n.d.	n.d.	8.2	n.d.	n.d.
17:1n-9	n.d.	n.d.	n.d.	n.d.	5.7	6.0
18:1n-9	176.3	170.1	293.7	335.9	115.1	210.6
18:1n-11	42.6	38.8	20.0	14.8	n.d.	n.d.
20:1n-9	29.8	19.0	12.1	9.5	n.d.	n.d.
22:1n-9	18.7	10.2	4.8	2.8	n.d.	6.2
24:1n-9	2.9	n.d.	n.d.	n.d.	n.d.	n.d.
∑Monoenes	308.8	268.7	343.9	371.2	136.0	256.0
C18:2n-6	260.2	312.4	344.0	363.1	61.1	85.1
C22:2n-6	n.d.	n.d.	n.d.	n.d.	5.7	n.d.
C18:3n-3	30.7	39.4	26.5	21.8	61.9	64.4
C18:3n-6	n.d.	n.d.	n.d.	n.d.	8.9	31.4
C18:4n-3	13.9	15.5	9.4	3.6	102.7	70.3
C20:3n-3	5.3	5.1	4.2	n.d.	106.8	145.8
C20:3n-6	n.d.	n.d.	n.d.	n.d.	5.2	6.0
C20:4n-3	2.9	n.d.	n.d.	n.d.	4.9	n.d.
C20:5n-3	60.9	52.4	26.8	14.5	n.d.	n.d.
C22:5n-3	7.3	5.0	n.d.	n.d.	57.6	62.8
C22:6n-3	50.0	43.4	20.8	11.9	n.d.	n.d.
∑n-3 HUFA	126.4	105.9	51.8	26.4	174.5	214.6
Unknown	6.9	5.6	2.7	n.d.	70.9	21.5

n.d. indicates not detected.

abalone fed the Std diet, and followed by the FM50, FM50+MA50 and FM50+MA100 diets, *Undaria*, and *L. japonica*. Shell height of abalone fed the formulated (Std, FM50, FM50+MA50 and

FM50+MA100) diets were not significantly different among treatments, but significantly taller ($P < 0.0009$) than that of abalone fed the *Undaria* and *L. japonica*. The soft body weight of abalone fed the Std diet was

Table 4. Survival (%), weight gain (g/abalone) and specific growth rate (SGR) of juvenile abalone fed experimental diets for 16 week

Experimental diets	Initial weight (g/abalone)	Final weight (g/abalone)	Survival (%)	Weight gain (g/abalone)	SGR ¹ (%/day)
Std	3.6±0.00	7.5±0.01 ^a	85.1±0.48 ^a	3.87±0.011 ^a	0.63±0.001 ^a
FM50	3.6±0.00	6.5±0.01 ^b	84.8±0.75 ^a	2.94±0.005 ^b	0.52±0.000 ^b
FM50+MA50	3.6±0.00	6.1±0.01 ^c	84.0±0.48 ^a	2.49±0.014 ^c	0.46±0.002 ^c
FM50+MA100	3.6±0.00	6.0±0.02 ^c	84.9±0.11 ^a	2.40±0.021 ^c	0.44±0.003 ^c
<i>U. pinnatifida</i>	3.6±0.00	5.6±0.04 ^d	80.4±0.18 ^b	2.00±0.038 ^d	0.38±0.006 ^d
<i>L. japonica</i>	3.6±0.00	4.6±0.06 ^e	72.4±0.27 ^c	1.02±0.063 ^e	0.22±0.012 ^e

Values (means of duplicate±SE) in the same column sharing the same superscript letter are not significantly different (P>0.05).

¹Specific growth rate (SGR, %/day) = [(Ln(Wf) - Ln(Wi))/days of feeding]×100, where Ln(Wf) = natural log of the final mean weight of abalone and Ln(Wi) = natural log of the initial mean weight of abalone.

Table 5. Shell length (mm), shell width (mm), shell height (mm), soft body weight (g) and the ratio of soft body weight to total weight of abalone fed the experimental diets for 16 week

Experimental diets	Shell length (mm)	Shell width (mm)	Shell height (mm)	Soft body weight (g)	Soft body weight/total weight
Std	41.7±0.14 ^a	28.5±0.01 ^a	8.6±0.14 ^a	4.3±0.05 ^a	0.61±0.004 ^a
FM50	40.6±0.05 ^b	28.0±0.33 ^{ab}	8.6±0.01 ^a	4.0±0.02 ^{ab}	0.60±0.003 ^a
FM50+MA50	40.0±0.10 ^c	27.2±0.41 ^b	8.6±0.20 ^a	3.8±0.21 ^b	0.60±0.013 ^a
FM50+MA100	39.7±0.00 ^d	27.1±0.09 ^b	8.3±0.12 ^a	3.7±0.04 ^b	0.61±0.004 ^a
<i>U. pinnatifida</i>	38.1±0.08 ^c	25.9±0.02 ^c	7.2±0.05 ^b	3.1±0.08 ^c	0.61±0.002 ^a
<i>L. japonica</i>	38.2±0.00 ^e	25.9±0.05 ^c	7.2±0.13 ^b	3.1±0.05 ^c	0.60±0.006 ^a

Values (means of duplicate±SE) in the same column sharing the same superscript letter are not significantly different (P>0.05).

significantly heavier (P<0.002) than all other diets, except for the FM50 diet. However, the ratio of the soft body weight to total weight of abalone was not significantly (P>0.4) different among the diets.

Moisture and ash content of the soft body of abalone was not significantly (P>0.6) different among the diets (Table 6). However, crude protein content of the soft body of abalone fed the FM50+MA100 diet was significantly higher (P<0.05) than that of abalone fed the all other diets. Crude protein content of the soft body of abalone fed the FM50+MA50 diet was also significantly higher (P<0.0003) than that of abalone fed the *Undaria* and *L. japonica*, but not significantly different from that of abalone fed the Std and FM50 diets. Crude lipid content of the soft body of abalone fed the FM50+MA50 and FM50+MA100 diets was significantly higher (P<0.0002) than that of abalone fed the all other diets. Crude lipid content of the soft body of abalone fed the Std and FM 50 diets was also significantly higher than that of abalone fed the *Undaria* and *L. japonica*. Crude protein and lipid content of the soft body of abalone was directly reflected from dietary protein and lipid contents.

Discussion

The several essential amino acids, such as arginine, isoleucine, lysine, methionine, threonine and valine were low in the FM50 diet, but arginine, histidine, isoleucine, leucine, phenylalanine and valine were high in the FM50+MA50 and FM50+MA 100 diets compared to the Std diet (Table 2). All essential amino acids were found, but relatively low

in the *Undaria* and *L. japonica* in this study, agreeing with Dawczynski, Schubert, & Jahreis (2007)'s study. Mai, Mercer, & Donlon (1994) also reported that the several essential amino acids, such as arginine, methionine, threonine and histidine were the limiting factor in 6 species of MA, *Ulva lactuca* Linne, *Chondrus crispus* Stackh, *Palmaria palmate* Linnaeus, *Alaria esculenta* Linne, *Laminaria digitata* Hudson and *Laminaria saccharina* Linne, for abalones, *H. tuberculata* Linnaeus and *H. discus hannai*. Histidine was also reported to be a limiting factor when fish meal was substituted with various sources of animal and/or plant protein sources (Cho, 2010).

A sum of n-3 HUFA content were high in the Std diet compared to that of the FM50, FM50+MA50 and FM50+MA100 diets, especially linolenic acid, EPA and DHA decreased in proportion to an amount of substitution of FM and MA with the fermented soybean meal and rice bran, respectively, but linoleic acid content increased. Similar EPA and DHA contents in soybean and rice bran oils were reported by Krishna, Hemakumar, & Khaton (2006) and Grisdale-Helland *et al.* (2002). No EPA and DHA was also found in MA, *Undaria* and *L. japonica*, diets in this study, partially agreeing with Dawczynski *et al.* (2007)'s study showing that 132 and 162 g/kg of EPA were detected in *Undaria pinnatifida* and *Laminaria* sp., but no DHA was detected in all seaweeds investigated. The requirement of n-3 HUFA was reported to be 10 g/kg in the diet containing 50 g/kg lipid, being equivalent to 0.5 g/kg in the diet for abalone, *H. discus hannai* (Uki, Kemuyama, &

Table 6. Chemical composition (g/kg) of the soft body of abalone at the end of the 16 week feeding trial

Experimental diets	Moisture	Crude protein	Crude lipid	Ash
Std	773±2.0 ^a	168±1.6 ^{bc}	9±0.4 ^b	28±1.2 ^a
FM50	778±1.2 ^a	167±0.8 ^{bc}	8±0.8 ^b	27±0.4 ^a
FM50+MA50	775±6.9 ^a	172±0.8 ^b	16±0.8 ^a	25±0.0 ^a
FM50+MA100	769±0.8 ^a	184±1.2 ^a	17±0.8 ^a	25±0.4 ^a
<i>U. pinnatifida</i>	770±0.4 ^a	163±1.6 ^c	6±0.4 ^c	29±0.4 ^a
<i>L. japonica</i>	771±0.0 ^a	158±1.2 ^d	6±0.4 ^c	27±0.4 ^a

Values (means of duplicate±SE) in the same column sharing the same superscript letter are not significantly different (P>0.05).

Watanabe, 1986b). However, the requirement of n-3 HUFA was all satisfied in the all diets, and ranged from 0.9 g/kg (*L. japonica*) to 1.9 g/kg (FM50+MA50 diet). Mai, Mercer, & Donlon (1996) reported that EPA played a prominent role for 2 species of abalone, *H. tuberculata* and *H. discus hannai*, and n-3 and n-6 polyunsaturated fatty acid (PUFA) seemed to be essential for abalone, *H. discus hannai*.

Poor performance in survival, weight gain and SGR of abalone fed the *Undaria* and *L. japonica* could be explained by the poor nutritional quantity and quality of essential amino and fatty acids in these MA (Tables 2 and 3). Similarly, MA commonly used as feed in wild did not produce the optimal performance of abalone (Lee, 1998; Cho et al., 2006; Kim et al., 1998; Naidoo et al., 2006; Cho et al., 2008; Gracia-Esquivel & Felbeck, 2009; Dang et al., 2011; Myung et al., 2016).

Poorer weight gain of abalone fed the FM50, FM50+MA50 and FM50+MA100 diets was observed compared to that of abalone fed the Std diet in this study. However, unlike this study, Cho (2010) showed that FM at 350 g/kg could be completely replaced with soybean meal at 580 g/kg with 5 g/kg methionine supplementation in the diet for juvenile abalone. The differences in performance of abalone between this study and Cho (2010)'s study could have resulted from the lack of methionine supplementation in this study. However, Lee et al. (1998) reported that the plant protein sources (soybean meal or cotton seed meal) without supplementation of amino acids and are likely to be deficient in plant protein sources could replace casein or FM in the diet for abalone, *H. discus hannai*. Fish meal could also be successfully replaced with a combined animal and plant protein sources in abalone diets (Guzman & Viana, 1998; Bautista-Teruel, Fermin, & Koshio, 2003; Cho, 2010).

Another reason for the poorer weight gain of abalone fed the FM50, FM50+MA50 and FM50+MA100 diets compared to that of abalone fed the Std diet could be due to the fermentation of soybean meal in this study. Abalone may not utilize the fermented soybean meal well. This is why all 500 g/kg fish meal substituted diets with the fermented soybean meal regardless of substitution of MA with rice bran produced poorer weight gain in this study. Similarly, juvenile abalone (*H. discus*) fed the FM-basal diet with 250 g/kg fermented soybean meal achieved poorer weight gain than that fed the FM-

basal diet with 250 g/kg soybean meal in the 16-week feeding trial (Cho, 2016; unpublished data). Nutritional values of fermented soybean meal varied depending on physical conditions of fermentation (Kook et al., 2014). More studies to elucidate the reason why abalone fed the diet containing fermented soybean meal showed poorer growth performance than that fed diet containing soybean meal is needed. Unlike this study, however, dietary inclusion of the fermented plant protein sources, such as soybean meal and cottonseed meal produced promising effect on performance of fish (Lim & Lee, 2011; Zhou et al., 2011).

Unlike Kim et al. (2015)'s study, 50% and 100% substitution of MA with rice bran in the FM50+MA50 and FM50+MA100 diets, respectively, produced poorer weight gain of abalone compared to the Std and FM50 diets in this study. The differences could have resulted from those in MA used in both studies. MA used in this study are the mixture of *Undaria* and *Hizikia fusiforme* at a ratio of 1:1, but a single *L. japonica* was used in the former. Abalone fed the diets (Std and FM50 diets) containing the mixture of *Undaria* and *Hizikia fusiforme* in this study outgrew over that fed the control diet containing *L. japonica* in Kim et al. (2015)'s study. The mean SGR was 0.58%/day in this study, but 0.46%/day for the latter. Therefore, substitution effect of MA with rice bran in this study was probably masked. The combined MA produced slightly, but not significantly, improved weight gain of abalone, *H. discus hannai* and *H. laevigata* Donovan, over a single MA (Qi et al., 2010; Dang et al., 2011).

No difference in weight gain and SGR of abalone fed the FM50+MA50 and FM50+MA100 diets (Table 4) in this study could indicate that another 50% MA in the commercial diet could be substituted with rice bran as long as 500 g/kg MA were substituted with rice bran. Abalone are known to be herbivorous and feed mostly on MA, which is usually low in protein and lipid, but high in carbohydrate, 400-500 g/kg in the wild (Thongrod, Tamtin, Chairat, & Boonyaratpalin, 2003). Unlike this study, however, Kim et al. (2015) reported that 400 g/kg substitution of *L. japonica* with rice bran at 200 g/kg in the diet achieved the best specific growth rate of abalone, *H. discus* and concluded that 100% substitution of *L. japonica* with rice bran at 200 g/kg in the diet was successfully made without retardation

of weight gain when juvenile abalone were fed on the experimental diets trailed over *L. japonica*. However, a 16 week trail may not be a sufficiently long enough time give that abalone remain on farms for considerably longer periods. Similarly, the leaf meal, *Moringa oliefera* Lamarck, and freshwater aquatic fern, *Azolla pinnata* R. Brown, are promising alternative feed ingredients for *H. asinina* (Linnaeus) culture (Reyes & Fermin, 2003).

Biological criteria of abalone measured in this study (shell length, shell width, shell height and soft body weight), except for the ratio of the soft body weight to total weight seemed to be closely related to growth rate of abalone. Similarly, biological criteria of abalone agreed with growth rate of abalone (Bautista-Teruel et al., 2003; Cho, 2010; Myung et al., 2016).

Crude protein and lipid content of the soft tissue directly related to dietary protein and lipid contents in this study, and agreed with other studies (Mai et al., 1995a, b; Thongrod et al., 2003; Cho et al., 2008; Garcia-Esquivel & Felbeck, 2009; Cho, 2010; Kim et al., 2015; Myung et al., 2016).

In conclusion, the substitutability of fermented soybean meal for fish meal seems to be rather limited in commercial diets for abalone farm. However, another 500 g/kg MA in the commercial diet could be substituted with rice bran as long as 500 g/kg MA were substituted with rice bran. In considering feed cost, the FM50+MA50 diet is the most recommendable for abalone farm.

Acknowledgments

This research was a part of the projected titled "Development of the granule-type artificial diet for larval fish" funded by the Ministry of Oceans and Fisheries, Korea. In addition, this work was supported by a grant from the National Institute of Fisheries Science (R2016016).

References

- AOAC (Association of Official Analytical Chemists), (1990). Official Methods of Analysis, 15th edn. Arlington, VA, USA.
- Bautista-Teruel, M.N., Fermin, A.C., & Koshio, S.S. (2003). Diet development and evaluation for juvenile abalone, *Haliotis asinina*: animal and plant protein sources. *Aquaculture*, 219(1-4), 645-653. [http://dx.doi.org/10.1016/S0044-8486\(02\)00410-6](http://dx.doi.org/10.1016/S0044-8486(02)00410-6)
- Britz, P.J. (1996). Effect of dietary protein level on growth performance of South African abalone, *Haliotis midae*, fed fishmeal-based semi-purified diets. *Aquaculture*, 140(1-2), 55-61. [http://dx.doi.org/10.1016/0044-8486\(95\)01198-6](http://dx.doi.org/10.1016/0044-8486(95)01198-6)
- Cho, S.H., Park, J., Kim, C., Yoo, J., & Lee, S. (2006). Effect of the various sources of dietary additives on growth, body composition and shell color of abalone *Haliotis discus hannai*. *Journal of Aquaculture*, 19(4), 275-280.
- Cho, S.H., Park, J., Kim, C., & Yoo, J. (2008). Effect of casein substitution with fishmeal, soybean meal and crustacean meal in the diet of the abalone *Haliotis discus hannai* Ino. *Aquaculture Nutrition*, 14(1), 61-66. <http://dx.doi.org/10.1111/j.1365-2095.2007.00505.x>
- Cho, S.H. (2010). Effect of fishmeal substitution with various animal and/or plant protein sources in the diet of the abalone *Haliotis discus hannai* Ino. *Aquaculture Research*, 41(10), e587-e593. <http://dx.doi.org/10.1111/j.1365-2109.2010.02561.x>
- Dang, V.T., Li, Y., Speck, P., & Benkendorff, K. (2011). Effects of micro and macroalgal diet supplementations on growth and immunity of hreenlip abalone, *Haliotis laevigata*. *Aquaculture*, 320(1-2), 91-98. <http://dx.doi.org/10.1016/j.aquaculture.2011.08.009>
- Dawezyński, C., Schubert, R., & Jahreis, G. (2007). Amino acids, fatty acids, and dietary fiber in edible seaweed products. *Food Chemistry*, 103(3), 891-899. <http://dx.doi.org/10.1016/j.foodchem.2006.09.041>
- Duncan D.B. (1955). Multiple range and multiple F tests. *Biometrics*, 11, 1-42.
- Food and Agriculture Organization of the United Nations (FAO). (2014). FAO Globefish reports.
- FishStatJ. (2015). Food and Agriculture Organization of the United Nations. Rome.
- Gao, M., Kaneko, M., Hirata, M.H., Toorisaka, E., & Hano, T. (2008). Utilization of rice bran as nutrient sources for fermentative lactic acid production. *Bioresource Technology*, 99(9), 3659-3664. <http://dx.doi.org/10.1016/j.biortech.2007.07.025>
- Gracia-Esquivel, Z., & Felbeck, H. (2009). Comparative performance of juvenile red abalone, *Haliotis rufescens*, reared in laboratory with fresh kelp and balanced diets. *Aquaculture Nutrition*, 15(2), 209-217. <http://dx.doi.org/10.1111/j.1365-2095.2008.00585.x>
- Grisdale-Helland, B., Ruyter, B., Rosenlund, G., Obach, A., Helland, S.J., Sandberg, M.G., Standal, H., & Rosjo, C. (2002). Influence of high contents of dietary soybean oil on growth, feed utilization, tissue fatty acid composition, heart histology and standard oxygen consumption of Atlantic salmon (*Salmo salar*) raised at two temperatures. *Aquaculture*, 207(3-4), 311-329. [http://dx.doi.org/10.1016/S0044-8486\(01\)00743-8](http://dx.doi.org/10.1016/S0044-8486(01)00743-8)
- Guzman, J.M., & Viana, M.T. (1998). Growth of abalone *Haliotis fulgens* fed diets with and without fish meal, compared to a commercial diet. *Aquaculture*, 165(3-4), 321-331. [http://dx.doi.org/10.1016/S0044-8486\(98\)00271-3](http://dx.doi.org/10.1016/S0044-8486(98)00271-3)
- Hong, K.J., Lee, C.H., & Kim, S.W. (2004). *Aspergillus oryzae* GB107 fermentation improves nutritional quality of food soybean and feed soybean meals. *Journal of Medicinal Food*, 7(4), 430-434. <http://dx.doi.org/10.1089/jmf.2004.7.430>
- Kim, J., Lee, S., Han, S., Kim, B., & Park, S. (1998). Effects of experimental diets, commercial diets and algae (*Undaria*) on growth and body composition among juvenile abalones (*Haliotis discus*, *H. sieboldii* and *H. discus hannai*). *Journal of Aquaculture*, 11(4), 505-512.
- Kim, Y.E., Myung, S.H., Kim, H.S., Jung, W., Cho, S.H., Jwa, M.S., Kim, P.Y., Park, M., & Kim, B. (2015). Effect of dietary substitution of sea tangle (ST), *Laminaria japonica* with rice bran (RB) on growth and body composition of juvenile abalone (*Haliotis discus*). *Aquaculture Research*, 47(4), 1202-1208. <http://dx.doi.org/10.1111/are.12577>

- Kook, M., Cho, S., Hong, Y., & Park, H. (2014). *Bacillus subtilis* fermentation for enhancement of feed nutritive value of soybean meal. *Journal of Applied Biological Chemistry*, 57(2), 183-188.
- Krishna, A.G., Hemakumar, K.H., & Khaton, S. (2006). Study on the composition of rice bran oil and its higher free fatty acids value. *Journal of the American Oil Chemists' Society*, 83(2), 117-120. <http://dx.doi.org/10.1007/s11746-006-1183-1>
- Lee, S. (1998). Evaluation of economical feed formulations for abalones (*Haliotis discus hannai*). *Journal of the Aquaculture*, 11(2), 159-166.
- Lee, S., Yun, S.J., & Hur, B. (1998). Evaluation of dietary protein sources for abalone (*Haliotis discus hannai*). *Journal of Aquaculture*, 11(1), 19-29.
- Lim, S., & Lee, K. (2011). A microbial fermentation of soybean and cottonseed meal increases antioxidant activity and gossypol detoxification in diets for Nile tilapia, *Oreochromis niloticus*. *Journal of the World Aquaculture Society*, 42(4), 494-503. <http://dx.doi.org/10.1111/j.1749-7345.2011.00491.x>
- Mai, K., Mercer, J.P., & Donlon, J. (1994). Comparative studies on the nutrition of two species of abalone, *Haliotis tuberculata* L. and *Haliotis discus hannai* Ino. II. Amino acid composition of abalone and six species of macroalgae with an assessment of their nutritional value. *Aquaculture*, 128(1-2), 115-130. [http://dx.doi.org/10.1016/0044-8486\(94\)90107-4](http://dx.doi.org/10.1016/0044-8486(94)90107-4)
- Mai, K., Mercer, J.P., & Donlon, J. (1995a). Comparative studies on the nutrition of species of abalone, *Haliotis tuberculata* L. and *Haliotis discus hannai* Ino. III. Responses of abalone to various levels of dietary lipid. *Aquaculture*, 134(1-2), 65-80. [http://dx.doi.org/10.1016/0044-8486\(95\)00043-2](http://dx.doi.org/10.1016/0044-8486(95)00043-2)
- Mai, K., Mercer, J.P., & Donlon, J. (1995b). Comparative studies on the nutrition of two species of abalone, *Haliotis tuberculata* L. and *Haliotis discus hannai* Ino. IV. Optimum dietary protein level for growth. *Aquaculture*, 136(1-2), 165-180. [http://dx.doi.org/10.1016/0044-8486\(95\)01041-6](http://dx.doi.org/10.1016/0044-8486(95)01041-6)
- Mai, K., Mercer, J.P., & Donlon, J. (1996). Comparative studies on the nutrition of two species of abalone, *Haliotis tuberculata* L. and *Haliotis discus hannai* Ino. V. The role of polyunsaturated fatty acids of macroalgae in abalone nutrition. *Aquaculture*, 139(1-2), 77-89. [http://dx.doi.org/10.1016/0044-8486\(95\)01158-7](http://dx.doi.org/10.1016/0044-8486(95)01158-7)
- Myung, S.H., Jung, W., Kim, H.S., Kim, Y.E., Cho, S.H., Jwa, M.S., Kim, P.Y., Kim, M.K., Park, M., & Kim, B. (2016). Effects of dietary substitution of fishmeal with the combined dry microalgae, *Nannochloropsis oceanica* (NO) biomass residue and casein on growth and body composition of juvenile abalone (*Haliotis discus*). *Aquaculture Research*, 47(1), 341-348. <http://dx.doi.org/10.1111/are.12562>
- Naidoo, K., Maneveldt, G., Ruck, K., & Bolton, J.J. (2006). A comparison of various seaweed-based diets and formulated feed on growth rate of abalone in a land-based aquaculture system. *Journal of Applied Phycology*, 18(3), 437-443. http://dx.doi.org/10.1007/978-1-4020-5670-3_26
- Neori, A., & Nobre, A.M. (2012). Relationship between trophic level and economics in aquaculture. *Aquaculture Economics and Management*, 16(1), 40-67. <http://dx.doi.org/10.1080/13657305.2012.649046>
- Qi, Z., Liu, H., Li, B., Mao, Y., Jiang, Z., Zhang, J., & Fang, J. (2010). Suitability of two seaweeds, *Gracilaria lemaneiformis* and *Sargassum pallidum*, as feed for the abalone *Haliotis discus hannai* Ino. *Aquaculture*, 300(1-4), 189-193. <http://dx.doi.org/10.1016/j.aquaculture.2010.01.019>
- Reyes, O.S., & Fermin, A.C. (2003). Terrestrial leaf meals or freshwater aquatic fern as potential feed ingredients for farmed abalone *Haliotis asinina* (Linnaeus 1758). *Aquaculture Research*, 34(8), 593-599. <http://dx.doi.org/10.1046/j.1365-2109.2003.00846.x>
- Thongrod, S., Tamtin, M., Chairat, C., & Boonyaratpalin, M. (2003). Lipid to carbohydrate ratio in donkey's ear abalone (*Haliotis asinina*, Linne) diets. *Aquaculture*, 225(1-4), 165-174. [http://dx.doi.org/10.1016/S0044-8486\(03\)00287-4](http://dx.doi.org/10.1016/S0044-8486(03)00287-4)
- Uki, N., Kemuyama, A., & Watanabe, T. (1985a). Development of semipurified test diets for abalone. *Bulletin of the Japanese Society of Scientific Fishers*, 51(11), 1825-1833. <http://dx.doi.org/10.2331/suisan.51.1825>
- Uki, N., Kemuyama, A., & Watanabe, T. (1985b). Nutritional evaluation of several sources in diets for abalone *Haliotis discus hannai*. *Bulletin of the Japanese Society of Scientific Fisheries*, 51(11), 1835-1839. <http://dx.doi.org/10.2331/suisan.51.1835>
- Uki, N., Kemuyama, A., & Watanabe, T. (1986a). Optimum protein level in diets for abalone. *Bulletin of the Japanese Society of Scientific Fisheries*, 52(6), 1005-1012. <http://dx.doi.org/10.2331/suisan.52.1005>
- Uki, N., Kemuyama, A., & Watanabe, T. (1986b). Requirement of essential fatty acids in the abalone *Haliotis discus hannai*. *Bulletin of the Japanese Society of Scientific Fishers*, 52(6), 1013-1023. <http://dx.doi.org/10.2331/suisan.52.1013>
- Zhou, F., Song, W., Shao, Q., Peng, X., Xiao, J., Hua, Y., & Owari, B.N. (2011). Partial replacement of fish meal by fermented soybean meal in diets for black sea bream, *Acanthopagrus schlegeli*, juveniles. *Journal of the World Aquaculture Society*, 42(2), 184-197. <http://dx.doi.org/10.1111/j.1749-7345.2011.00455.x>