



Mesh Size Recommendation for Turkey Pike (*Esox lucius* L., 1758) Gillnet Fishery

Mehmet Cilbiz^{1*}, Rahmi Uysal¹, Ahmet Alp², Vedat Yeğen¹, Meral Apaydın Yağcı¹, Abdulkadir Yağcı¹, Ramazan Küçükbara¹

¹Fisheries Research Institute, Eğirdir - Isparta 32500, Turkey.

²Kahramanmaraş Sütçü İmam University, Agricultural Faculty, Fisheries Department, Kahramanmaraş - 46100 Turkey.

* Corresponding Author: Tel.: +90.246 3133460; Fax: +90.246 3133460;
E-mail: mehmetcilbiz@hotmail.com

Received 11 November 2016
Accepted 23 January 2017

Abstract

The pike is one of the most valuable freshwater fish species in Turkey. This study was carried out in order to estimate the selectivity properties of gillnet used in pike (*Esox lucius* L., 1758) fishing in Çivril Lake, Turkey. The study was performed between January 2005 and December 2005 and selectivity estimates were based on six different monofilament gillnet mesh sizes (i.e. 4, 5, 6, 7, 8 and 9 cm). The SELECT method was used to estimate the selectivity parameters. According to the bi-normal model, optimum model lengths for 4, 5, 6, 7, 8 and 9 cm mesh sizes were determined as 20.99, 26.24, 31.49, 36.73, 41.98 and 47.23 cm respectively. When considering minimum landing size, the findings of the study showed that the use of gillnets of mesh size less than 8 cm should be prohibited for the fishing of *E. lucius* in Çivril Lake.

Keywords: Gillnet selectivity, *Esox lucius*, SELECT method, Çivril Lake, Pike.

Introduction

The Pike (*Esox lucius* Linnaeus, 1758) is widely distributed in Europe, Asia and North America. It is an important species for commercial and recreational fisheries throughout its natural range (Moslemi-Aqdam, Imanpour Namin, Sattari, Abdolmalaki, & Bani, 2014; Paukert, Stancill, DeBates, & Willis, 2003).

E. lucius is also a main target species for inland fisheries in Turkey due to the high commercial value. There has been an increase in fishing pressure on *E. lucius* and a significant reduction in its landings in Turkish water. For instance, landing amount reached 350 t in 1997 but decreased to 203 t in 2015 (Anonymus 2016). Aside from fishing pressure, other reasons for that reduction in landing might be pollution, diseases, habitat degradation and invasive species.

Apaydın Yağcı, Alp, Uysal, Yeğen, and Yağcı (2009) reported the first maturity length as 24.9 cm for female and 22.9 cm for male *E. lucius* caught from the Lake Işıklı. In another study carried out by Balık, Çubuk, Özkök, and Uysal (2004) in the Lake Karamık, the first maturity length was reported as 18 cm (fork length) for female *E. lucius*.

One of the main principles for sustainable fisheries management is to let fish to reproduce at least once in natural habitat. In the other words, catch size should be larger than the length at first maturity

(L_m). Thus, it is necessary to know the selectivity properties of fishing gear used in commercial fisheries. Fisheries management decision makers can make legal regulation for mesh size to catch fish above first breeding by using the size selectivity data. That is to say it is very easy to implement the results of studies on selectivity of gill nets within the scope of minimum mesh size.

Gear selectivity is an important tool for fisheries managers who, by regulating the minimum mesh sizes of a fishing fleet, can more or less determine the minimum sizes of the target species of certain fisheries (Sparre & Venema, 1998). An ideal selection of a fishing gear for optimum fish length should aim to catch larger fish than the first maturity (Yüksel, Gündüz, & Demiroğlu, 2014). Therefore, result of this study are very valuable for improving the management of commercial pike fishery in Çivril Lake and other inland waters of Turkey.

One of the most selective fishing gears is gillnet and it allow fisherman to catch certain sized species at an optimum level (Hamley, 1975; Hoşsucu, 2011; Kiyaga, 2008; Özekinci, 1995). Generally, fishermen use gillnet for fishing *E. lucius* in the Lake Çivril because of easy use, affordability, durability and productivity.

Legislative regulations for both minimum landing size and minimum mesh size are the case in commercial pike fisheries. However, gillnet

selectivity properties for current minimum landing size (40 cm total length) is unclear. This present results would be very valuable for fisheries management authorities who are responsible for the determination of minimum mesh size for commercial pike fisheries.

There are very limited studies selectivity characteristics of gillnets in *E. lucius* fishing. Pierce, Tomcko, and Kolander (1994) compared size selectivity of gill net by indirect and direct methods. Authors reported that indirect estimates show that gill nets were most effective for retaining northern pike when fish length/mesh perimeter ratio was between 3.5 and 3.7 besides increasing selectivity with increasing fish length by the direct estimates. In the another study which was conducted by Balık (2008), investigated selectivity properties of gillnets by SELECT method for 3.6, 4, 4.4, 5 and 6 cm mesh sizes. In this study, we improved former research in respect to used model type and large mesh size range in accordance with commercial fishery. The purpose of this study was to estimate the selectivity properties of gillnets for commercial fishery of *E. lucius* in the Lake Çivril.

Materials and Methods

The Lake Çivril is located within the boundaries of Denizli province (west Anatolia) (Figure 1). Surface of the lake is 64.53 km², maximum depth is 7 m and altitude is 821 m (İlhan & Balık, 2003). Fish samplings were carried out monthly in two different stations with a total of 24 trials between from January and December 2005. Gillnets were made of monofilament material with 4, 5, 6, 7, 8 and 9 cm stretched mesh sizes, 0.20 mm twine thickness and 0.50 of hanging ratio. Depths of all nets were 50 meshes and panel length of them 100 m. All nets were set in the afternoon and retrieved in the next morning. Fish were classified according to mesh sizes of the nets and total fish length was measured using a

measurement board with 1 mm precision.

The SELECT (Share Each Length's class Catch Total) method was used to determine selectivity as an indirect estimation method (Millar, 1992; Millar & Fryer, 1999; Millar & Holst, 1997). Data obtained from field studies were analyzed with RStudio (version 0.99.903) (RStudio, 2016). R codes are developed by Millar (2010, 2011). Length selectivity of each mesh size was described by five different models (normal location, normal scale, gamma, lognormal and bi-normal) of the SELECT method (Millar & Fryer, 1999; Park *et al.*, 2011). The Kolmogorov-Smirnov (*K-S*) test was used to determine differences between pairs of size frequency distributions per mesh size of the net (Karakulak & Erk, 2008; Siegel & Castellan, 1989).

Results

A total of 317 specimens of *E. Lucius* was caught over the study period. Fish lengths ranged between 22.9 and 52.9 cm. Catch composition and length frequency distributions of specimens are shown in Table 1 and Figure 2 for each mesh size of the nets, respectively.

A comparison of the deviances of five models of SELECT method revealed that bi-normal model yielded the best fit with the lowest deviance of 139.41 (Table 2). The selectivity curves and deviance residual plots are shown in Figure 3 were drafted by bi-normal parameters via RStudio software (RStudio, 2016). The optimum length and spread values estimated by the bi-normal model per different mesh sizes of the net are shown in Table 3. Estimated model length for 4 cm mesh size was adapted as scale proportionally for all other mesh sizes (i.e. 5, 6, 7, 8 and 9 cm) (Park *et al.*, 2011). The *K-S* test showed differences between all paired comparisons of length frequency distributions per mesh sizes of gillnets (Table 4).





Figure 1. Study area and sampling locations.

Table 1. Descriptive statistics of the sampled individuals caught with the different mesh sizes of the nets

Length of mesh size (cm)	Number of fish caught (N)	Number of fish caught (%)	Average length (SD) (cm)	Minimum Length (cm)	Maximum Length (cm)
4	30	9.4	26.2(2.3)	23.9	34.2
5	76	23.8	29.1(4.1)	22.9	46.6
6	131	40.9	33.3(4.0)	23.8	52.9
7	47	14.7	37.7(3.6)	30.1	46.7
8	24	7.5	41.6(2.5)	34.7	46.1
9	9	2.8	43.9(4.4)	35.8	50.8

**Figure 2.** Length frequency distributions of *E. lucius*.**Table 2.** Selectivity model parameters of *E. lucius* and estimated selection curves for the 4 cm mesh size

Model	Parameters	Equal fishing power					Deviance	df
		Estimates	Mode 1	Spread 1	Mode 2	Spread 2		
Normal location	k	5.53(0.04)	22.15(0.16)	4.35(0.19)	-	-	199.21	153
	σ	4.35(0.19)						
Normal scale	k ₁	5.73(0.04)	22.94(0.18)	2.98(0.12)	-	-	221.52	153
	k ₂	0.55(0.04)						
Lognormal	μ_l	3.12(0.00)	22.36(0.16)	2.94(0.13)	-	-	183.86	153
	σ	0.12(0.00)						
Gamma	k	0.09(0.00)	22.54(0.17)	2.93(0.13)	-	-	194.37	153
	α	60.83(5.02)						
Bi-normal	k ₁	3.04	20.99(0.15)	1.11(0.17)	23.65(0.44)	3.77(0.31)	139.41	135
	k ₂	0.05						
	k ₃	3.18						
	k ₄	0.15						
	c	0.81						
Model	Parameters	Fishing power α mesh size					Deviance	df
		Estimates	Mode 1	Spread 1	Mode 2	Spread 2		
Normal location	k	5.63(0.04)	22.53(0.18)	4.42(0.20)	-	-	195.31	153
	σ	4.42(0.20)						
Normal scale	k ₁	5.83(0.04)	23.33(0.18)	2.95(0.12)	-	-	222.06	153
	k ₂	0.54(0.04)						
Lognormal	μ_l	3.14(0.00)	22.73(0.17)	2.99(0.14)	-	-	183.86	153
	σ	0.12(0.00)						
Gamma	k	0.09(0.00)	22.92(0.18)	2.96(0.13)	-	-	194.37	153
	α	61.83(5.02)						
Bi-normal	k ₁	3.04	21.05(0.15)	1.11(0.17)	24.21(0.49)	3.88(0.34)	139.41	135
	k ₂	0.05						
	k ₃	3.21						
	k ₄	0.15						
	c	0.68						

(Standard errors are in parentheses; bold value is lowest deviance).




Figure 3. Selectivity curves (A) and deviance residual plots (B) of gillnets for the *E. Lucius*.

Table 3. Optimum length and spread values of *E. lucius* according to the Bi-normal model

Mesh size (cm)	Model Length (cm)	Spread Value (cm)
4	20.99	1.11
5	26.24	1.39
6	31.49	1.67
7	36.73	1.94
8	41.98	2.22
9	47.23	2.50

Table 4. Results of the K-S test used to compare length frequency distributions between pairs of different mesh sizes of gillnets of *E. Lucius*

Net 1	Net 2	K-S Test	Decision	Net 1	Net 2	K-S Test	Decision
4	5	0.4792>0.2912	H ₀ Reject	5	9	0.9099>0.4082	H ₀ Reject
4	6	0.8296>0.2745	H ₀ Reject	6	7	0.6065>0.2245	H ₀ Reject
4	7	0.9471>0.3129	H ₀ Reject	6	8	0.8574>0.2832	H ₀ Reject
4	8	0.9667>0.3574	H ₀ Reject	6	9	0.8467>0.3955	H ₀ Reject
4	9	1.0000>0.4516	H ₀ Reject	7	8	0.6520>0.3247	H ₀ Reject
5	6	0.6472>0.1950	H ₀ Reject	7	9	0.6890>0.4262	H ₀ Reject
5	7	0.7766>0.2462	H ₀ Reject	8	9	0.4968>0.4683	H ₀ Reject
5	8	0.9380>0.3007	H ₀ Reject				

H₀: There are no significant differences in the length frequency distributions ($\alpha=0.05$; $k=1.36$).

Discussion

There is a circular regulating the commercial fishery in Turkey. According to this, legal minimum landing size of *E. lucius* is 40 cm total length for all inland waters of Turkey. Besides the sustainability of *E. lucius* commercial stocks, catch quotas and closed season regulations are issued by certain governmental organizations in Turkey. All of these regulations are

not only for *E. lucius* but also perform on other commercial inland species (*Cyprinus carpio*, *Astacus leptodactylus*, *Sunder lucioperca*, *Siluris glanis*, *Perca fluviatilis*, *Leuciscus cephalus* etc...). The regulations of minimum mesh size limitations are monitored by provincial directorates on behalf of Ministry of Food, Agriculture and Livestock Directorate.

Expectedly, the mean total lengths increased

with increasing mesh size. The most productive net was that of 6 cm mesh size, whereas the worst was that of 9 cm. The best suitable model was the bi-normal model in terms of the modal deviances (Table 2), which is considered as the most suitable in cases of capturing fish in mesh by jamming, wrapping and trammeling and large length frequency ranges in captured fish (Akamca, Kiyaga, & Özyurt, 2010; Holt, 1963; Hovgård, 1996). In this study, we observed many cases of tangled fish captured from sharp teeth of the pike. This result blend in Pierce et al. (1994), who reported that *E. lucius* captured by gill nets wedging or tangling.

There are very limited studies on *E. lucius* gillnet selectivity. One of them was conducted by Balık (2008) in Karamık Lake with the SELECT method and the authors reported as model lengths as 20.420, 22.689, 24.958, 28.362 and 34.034 cm (as fork length) for 3.6, 4, 4.4, 5 and 6 cm mesh size.

Reported optimum fork lengths by Balık (2008) for *E. lucius* in Lake Karamık are higher than those of the present study for the similar mesh sizes. One reason for this difference can be habitat and seasonal differences. Accordingly, selectivity of gillnets can vary for each fish species so selectivity should be determined separately for each species (Balık and Çubuk 2001). Another reason can be the model selected for the calculation. While the calculation was made according to the lognormal model in the former study (Balık, 2008), the bi-normal model was used in present study.

The estimation of selectivity parameters is needed to ensure a proper management of commercial gillnet fishery; protection of small fish thereby escape from fishing gear. Therefore, it is necessary to determine the optimum mesh size for sustainable production (Hamley, 1975; Sümer et al., 2010).

Taking into account the first maturity estimates reported by Apaydın Yağcı et al. (2009) and Balık et al. (2004) as well as the legal size limitation (40 cm), gillnet mesh sizes for catching *E. lucius* in Çivril Lake should not be less than 8 cm. Given also that, *E. lucius* is a very popular fish for anglers in Turkey, studies are required to determine the effects of hook selectivity and amateur fishing activities on the stocks.

Acknowledgements

This work was supported by the Republic of Turkey, Ministry of Food Agriculture and Livestock, General Directorate of Agricultural Research and Policy under grant TAGEM/HAYSÜD/2002/09/02/02 project number.

References

Akamca, E., Kiyaga, V. B., & Özyurt, C. E. (2010). The selectivity of monofilament trammel nets used in catch of gill head bream (*Sparus aurata*, Linnaeus,

1758) in İskenderun Bay-Turkey[in Turkish]. *Journal of FisheriesSciences.com*, 4(1), 28-37. doi: DOI: 10.3153/jfscom.2010004a

Anonymous, 2016. Turkish Statistical Institute. Fishery Statistics.

http://www.turkstat.gov.tr/PreTablo.do?alt_id=1005 (accessed Jenuary 13, 2014).

Apaydın Yağcı, M., Alp, A., Uysal, R., Yeğen, V., & Yağcı, A. (2009). Reproduction Properties of Pike (*Esox lucius* L.,1758) Population in the Işıklı Dam Lake (Çivril-Denizli/Turkey) [in Turkish]. *Journal of FisheriesSciences.com*, 3(3), 220-230. doi: 10.3153/jfscom.2009027

Balık, İ. (2008). Gillnet selectivity for pike, *Esox lucius*, in Lake Karamık, Turkey. *Eifac Symposium On Interactions Between Social, Economic And Ecological Objectives Of Inland Commercial And Recreational Fisheries And Aquaculture, Antalya, EIFAC Occasional*, 44, 93-99.

Balık, İ., & Çubuk, H. (2001). Selectivity of gillnets for catching rudd (*Scardinius erythrophthalmus* L. 1758) and white bream (*Blicca björkna* L., 1758) in Lake Ulubat (Apolyont,) [in Turkish]. XI. Ulusal Su Ürünleri Sempozyumu Bildirileri, Hatay, 04- 06 Eylül. Cilt :1 Mustafa Kemal Üniversitesi Su Ürünleri Fakültesi. 1-10.

Balık, İ., Çubuk, H., Özkök, R., & Uysal, R. (2004). Reproduction Properties of Pike (*Esox lucius* L., 1758) Population in Lake Karamık (Afyonkarahisar/Turkey). *Turkish Journal of Zoology*, 30, 27-30.

Hamley, J. M. (1975). Review of gillnet selectivity. *J. Fish. Board. Can*, 32, 1943-1969. doi: 10.1139/cjfas-2013-0401

Holt, S. J. (1963). A method for determining gear selectivity and its application. *ICNAF Special Publication*, 5, 106-115.

Hoşsucu, H. (2011). *Balıkçılık 1 (Avlama Araçları ve Teknolojisi) [in Turkish]*. Ege Üniversitesi Basımevi, İzmir: Ege Üniversitesi Yayınları Yayın No:55.

Hovgård, H. (1996). A two-step approach to estimating selectivity and fishing power of research gill nets used in Greenland waters. *Canadian Journal of Fisheries and Aquatic Sciences*, 53(5), 1007-1013. doi: 10.1139/f96-022

İlhan, A., & Balık, S. (2003). Investigation of the Bioecological Characteristics of Pike Population (*Esox lucius* Linnaeus, 1758) in Lake Işıklı (Çivril-Denizli/Turkey)[in Turkish]. *Süleyman Demirel Üniversitesi, Eğirdir Su Ürünleri Fakültesi Dergisi*, 1, 1-9.

Karakulak, F. S., & Erk, H. (2008). Gill net and trammel net selectivity in the northern Aegean Sea, Turkey. *Scientia Marina*, 72(3), 527-540.

Kiyaga, V. B. (2008). *The Investigation of the Selectivity of Monofilament Gill Nets Used in Catch of Pike Perch (Sander lucioperca Bogustkaya & Naseka, 1996) In Seyhan Dam Lake [in Turkish]*. (Thesis MSc), Çukuraova University, School of Natural and Applied Sciences, Adana.

Millar, R. B. (1992). Estimating the Size-Selectivity of Fishing Gear by Conditioning on the Total Catch. *Journal of the American Statistical Association*, 87(420), 962-968. doi: 10.2307/2290632

Millar, R. B., & Fryer, R. J. (1999). Estimating the size-selection curves of towed gears, traps, nets and hooks. *Reviews in Fish Biology and Fisheries*, 9(1), 89-116.

- doi: 10.1023/A:1008838220001
- Millar, R. B., & Holst, R. (1997). Estimation of gillnet and hook selectivity using log-linear models. *ICES Journal of Marine Science*, 54(3), 471-477. doi: 10.1006/jmsc.1996.0196
- Millar, R.B. (2010). R Code for fitting SELECT models to gillnet data. In: Department of Statistics, University of Auckland, New Zealand. Retrieved from <https://www.stat.auckland.ac.nz/~millar/selectware/R/Next/NextGeneration.R> (Accessed: November 9, 2016).
- Millar, R.B. (2011). R Code for fitting SELECT models to gillnet data. In: Department of Statistics, University of Auckland, New Zealand. Retrieved from <https://www.stat.auckland.ac.nz/~millar/selectware/R/gillnets/gillnetfunctions.R> (Accessed: November 9, 2016).
- Moslemi-Aqdam, M., Imanpour Namin, J., Sattari, M., Abdolmalaki, S., & Bani, A. (2014). Length- length, length -weight relationship and relative condition factor of pike, *Esox lucius* Linnaeus, 1758, in Anzali Wetland (Southwest of the Caspian Sea). *Caspian Journal of Environmental Sciences*, 12(1), 109-117.
- Özekinci, U. (1995). *Selectivity trial of gill net which have 18, 20, 22 mm mesh size [in Turkish]*. (Msc Thesis), Ege University, Grad. School of Natural and Applied Sciences, Bornova, İzmir.
- Park, H.-H., Millar, R. B., Bae, B.-S., An, H.-C., Chun, Y. Y., Yang, J. H., & Yoon, S. C. (2011). Size selectivity of Korean flounder (*Glyptocephalus stelleri*) by gillnets and trammel nets using an extension of SELECT for experiments with differing mesh sizes. *Fisheries Research*, 107(1-3), 196-200. doi: <http://dx.doi.org/10.1016/j.fishres.2010.10.020>
- Paukert, C. P., Stancill, W., DeBates, T. J., & Willis, D. W. (2003). Predatory Effects of Northern Pike and Largemouth Bass: Bioenergetic Modeling and Ten Years of Fish Community Sampling. *Journal of Freshwater Ecology*, 18(1), 13-24. doi: 10.1080/02705060.2003.9663947
- Pierce, R. B., Tomcko, C. M., & Kolander, T. D. (1994). Indirect and Direct Estimates of Gill-Net Size Selectivity for Northern Pike. *North American Journal of Fisheries Management*, 14(1), 170-177. doi: 10.1577/1548-8675(1994)014<0170:IADEOG>2.3.CO;2
- RStudio (2016). RStudio: Integrated development environment for R (Version 0.99.903) [Computer software]. Boston, MA. Retrieved November 10, 2016. Available from <https://www.rstudio.com/products/rstudio/download3/>
- Siegel, J., & Castellan, N. S. (1989). *Non parametric statistics for the behavioural sciences. Statistics Series, 2nd Edition*: Mc Graw Hill, New York.
- Sparre, P., & Venema, S. C. (1998). *Introduction to Tropical Fish Stock Assessment, Part 1-Manual* (Vol. 306/1): FAO Fisheries Technical Paper.
- Sümer, Ç., Özdemir, S., & Erdem, Y. (2010). Determination of Selectivity Monofilament and Multifilament Gillnets with Different Mesh Size for Horse Mackerel (*Trachurus trachurus* L., 1758) [in Turkish]. *Ege Üniversitesi Su Ürünleri Dergisi*, 27(3), 125-128.
- Yuksel, F., Gündüz, F., & Demiroglu, F. (2014). Gillnet selectivity for *Luciobarbus esocinus* (Heckel, 1843) in Keban Dam Lake, Elazığ, Turkey. *Indian Journal of Fisheries*, 61(2), 108-111.