



Photoperiodic Modulation on Growth and Behaviour of the Giant Gourami, *Trichogaster fasciata* (Bloch and Schneider, 1801)

Farah Bano¹, Mohammad Serajuddin^{1,*}

¹ University of Lucknow, Department of Zoology, Fish Biogenetics Research Lab, Lucknow-226007 (U.P.), INDIA.

² Address.

* Corresponding Author: Tel.: +91.941 5871277; Fax: ;
E-mail: lu.fisheries@gmail.com

Received 24 January 2017
Accepted 15 May 2017

Abstract

The influence of different photoperiods was studied on the growth and behaviour of the giant gourami, *Trichogaster fasciata*. Fish (weight=1.1g; length=4.36cm) were subjected to photoperiod regimes of 0 L: 24 D, 8 L: 16 D, 12 L: 12 D and 16 L: 8 D for 90 days in triplicates. The growth rate was highest (weight=4g) and lowest (weight=2.36g) in the groups subjected to 16L: 8D and 0 L: 24 D photoperiods respectively. The mean body weight and mean total length were significantly higher ($p < 0.05$) while standard length and caudal fin length ($p > 0.05$) were not significant in the 16 hours light. Mean values of food conversion ratio and condition factor were lowest while various growth parameters were found to be maximum in the 16 hours light. 16 L: 8 D photoperiod regime produced pronounced effects on the welfare of giant gourami as shown by the nonintrusive welfare indicators. The photoperiodic signals were found to be capable of modifying the behavioural activities where male aggression related to territory or defence and female selection were reduced in control group. The optimum photoperiod (16 L: 8 D) was an environmental cue for the better growth, welfare and behaviour.

Keywords: Aggression, female selection, growth, photoperiod, welfare indicators.

Introduction

Photoperiod acts as a zeitgeber which controls growth through endogenous rhythms and different circulating levels of growth hormones in body (Simensen et al., 2000). Specific photoreceptor cells of retina of the eyes and pineal gland produce melatonin in fish which modulates the production of growth hormone releasing factors (Moyle & Cech, 2000; Falcon et al., 2010; Reinecke, 2010). The appetite, food conversion and growth energy requirement are dependent on the secretion of growth hormones in fish (Donaldson et al., 1979; Bjornsson et al., 1989) and the feeding activity is regulated by the light: dark cycle (Boujard & Leatherland, 1992). Photoperiod was reported to increase the growth and survival of larvae (Boeuf & Baille, 1999) of fish particularly in temperate regions through photo stimulation which induced feed intake and improved the feed conversion efficiency (Woiwode & Adelman, 1991). Effects of different photoperiods on growth of various fishes have been reported by a number of workers, notable among them are Davies et al. (1986 a/ b); Imsland et al. (1995); Arvedlund et al. (2000); Purchase et al. (2000); Ergün et al. (2003); Rad et al. (2006); Taylor et al. (2006); Valenzuela et al. (2006);

Bonnet et al. (2007); Askarian and Kousha (2009) and Yager and Yigit (2009). The developmental and maturation process are governed by genetic and environmental factors in fish. Yamamoto et al. (2001) reported the effect of temperature and photoperiod on the growth and maturation in fish. The other factors affected growth of fish were found to be feeding rate, feed, water quality, stock density and size (Trzebiatowski et al., 1981). The optimum photoperiod can successfully be used to raise the yield of fish in aquaculture was concluded by many workers such as Saunders and Henderson (1970); Lundqvist (1980); Brauer (1982) and Saunders et al. (1985) who worked on relationship between photoperiod and growth rate in fishes. Ruchin (2004, 2006) investigated the influence of colour light on growth performance of *Carassius carassius*, and also on the effect of light on white blood cell count in *Cyprinus carpio*. Davies et al. (1986a/b) and (Davies & Hanyu, 1986) worked on the effect of temperature and photoperiod on sexual maturity and spawning of the common carp. Bhattacharyya et al. (2005) carried out the effect of photoperiod on gonadal activity of *Catla catla*. A reaction of environment determines the behaviour which acted as key element for fish welfare as reported by Martins et al. (2012). The welfare

indicators must provide information on potential problems and the causes of impaired welfare (Rousing et al., 2001). The welfare of fish denoting the fish being healthy, comfortable, well-nourished, safe and able to express their innate behaviour and not suffering from unpleasant states such as pain, fear or distress. Mustapha et al. (2014) reported that the welfare can be greatly influenced by photoperiod in *Oreochromis niloticus* and *Clarias gariepinus*. The influence of photoperiodism on reproductive physiology and behaviour was reported by Fiszbein et al. (2010) in the cichlid fish, *Cichlasoma dimerus*. The specific behavioural signals for an early assessment of fish welfare are considered to be important for both farmers and researchers in a developing aquaculture industry.

The studies showed that behavioural changes can be interpreted as either good or poor welfare depending on the fish species. A survey of literature showed that the information on the influence of light on growth and behaviour of giant gourami is scanty. Hence, the present study was undertaken to evaluate the effects of different photoperiod regimes on the welfare, survival, growth and behaviour in the giant gourami, *T. fasciata*.

Materials and Methods

A total of 200 samples of giant gourami, *T. fasciata* (weight: 0.9 to 1.7 g; length: 4.0 to 4.9 cm) were collected using cast and drag nets from river Gomti situated at Lucknow region (26° 56' N 80° 43' E) and acclimatized in the laboratory for a month. The experiment was conducted in glass aquaria (0.45×0.22×0.30 m) for the duration of 90 days where individuals of giant gourami were classified into four groups namely A, B, C and D in triplicates. Each group containing 10 samples of fish (length=1.11cm; weight=4.36g) were subjected to different photoperiod regimes of 0 L: 24 D as control and the other three were 8 L: 16 D, 12 L: 12 D and 16 L: 8 D respectively. All the aquaria were properly covered with black paper so that the light does not escape outside and also to avoid entry of external light. Artificial white fluorescent lamps of 40 watt and

intensity of light of 200 lux were installed at a height of 14 cm above the water level in each aquarium. The experiment was carried out in dark room to avoid hindrance of natural light. The physico-chemical conditions such as ambient temperature, dissolved O₂, pH and volume of water were maintained constant in aquaria throughout the experiment. Fish were fed with artificially prepared balanced food comprised of 35% fish meal, 28% mustard oil cake, 28% rice barn, 2% sunflower and cod liver oils each, 5% carboxy methyl cellulose and multivitamin and multimineral tablets ('Becozyme Forte' Glaxo India Ltd, 25 tablets/kg foods) throughout the investigation. The faecal matter was removed daily before feeding the artificial food to the fish. The amount of food offered to each aquarium was kept constant (5% of body weight of stocked fish).

The weight of the fish exposed to the four different photoperiod regimes were taken using an electronic balance sensitive up to 0.001 g in the beginning and at a regular interval of 15 days till the end of experiment. The total length (tip of the snout to the tip of the longest caudal fin ray), standard length (tip of the snout to the base of the caudal fin) and caudal fin length (terminal vertical fin of a fish) were measured. The various parameters of weight and length of fish samples exposed to the three light regimes and darkness are given in Table 1.

Growth Parameters

Feed Conversion Ratio (FCR), Specific Growth Rate (SGR), Relative Growth Rate (RGR), Absolute Growth Rate (AGR), Weight Gain were calculated using equations as per Degani et al. (1989).

FCR = Weight of Dry Feed used/ Wet Weight of Fish (g)

Percentage of SGR = [(Final Weight – Initial Weight)/ (Days)] × 100

Percentage of RGR = [(Final Mean Weight- Initial Mean Weight)/ Initial Mean Weight] × 100

Percentage of AGR = Final Mean Weight- Initial Mean Weight

Weight Gain (mg/g/day) = [(Final Mean Weight-Initial Mean Weight)/ (Initial Mean Weight / Days)] × 100

Table 1. The effects of different photoperiod regimes on body weight (g), total length (cm), standard length (cm) and caudal fin length (cm) in *T. fasciata* after.

Parameter	Photoperiod			
	Group A	Group B	Group C	Group D
Initial Body Weight	1.11±0.44	1.11±0.71	1.11±0.23	1.11±0.42
Final Body Weight	2.53±0.36	2.80±0.43	3.60±0.48	4.00±0.44
Initial Total Length	4.36±0.71	4.36±0.43	4.36±0.49	4.36±0.70
Final Total Length	4.63±0.74	5.10±1.07	5.99±0.88	7.02±1.66
Initial Standard Length	3.42±0.61	3.41±0.46	3.40±0.81	3.41±0.45
Final Standard Length	3.78±0.34	4.44±0.94	4.88±0.64	5.31±0.82
Initial Caudal Fin Length	0.90±0.79	0.90±0.59	0.92±0.72	0.94±0.60
Final Caudal Fin Length	1.20±0.62	1.66±0.96	2.00±0.24	2.47±0.60

Condition Factor

The study of well being and fatness of fish was calculated through the Fluton's Condition Factor (Ricker, 1973) using the formula: $K=W \times 100/L^3$

Where K is Fluton's condition factor, W is net wet weight (g) of fish and L is length in cm. The factor 100 is used to bring K close to unity.

Evaluation of Welfare of the Fish

The welfare of fish in each photoperiod tanks was conducted visually daily through the use of non-intrusive signs and danger signals. Behaviour, swimming activity, injuries and colour were studied during the experimental period as per Mustapha et al. (2014). Winner status was assigned to the fish that were dominated and loser (subordinate) were defeated, suppressed and also stressed.

Behavioral Study

Behavioural study on *T. fasciata* was carried out to observe the female choice and male aggressive behaviour of different experimental groups such as control group (A) vs B, C and D as per Fiszbein et al. (2010) on cichlid fish, *Cichlasoma dimerus*. Altogether twelve observations were made to study the behavioural activity (female selection and aggressive interactions) at an interval of five minutes for duration of 1 h.

(A) Effect of Photoperiod on Female Selection

After 90 days of experiment, male and female gouramis were selected from three experimental groups namely 'B' (8 L: 16 D), 'C' (12 L: 12 D) and 'D' (16 L: 8 D). The female gourami were isolated from control group 'A' and placed in a neutral tank with males and females from different groups of the experiment i.e., females (control) vs group B (male & female); females (control) vs group C (male & female) and females (control) vs group D (male & female). The observation was made to analyse the selection of females (control or experimental) by males. The number of attack activity by males and females of three experimental groups towards the female of control group was counted.

(B) Aggressive behaviour between males

The male gouramis were selected from control group and from different experimental groups such as B, C and D and were placed in three tanks separately with a big stone in the middle as: males (control) vs B, males (control) vs C and males (control) vs D. Winner and loser status was assigned to the fish that performed more aggression and received more aggressive contacts respectively. The number of these interactions was counted.

(C) Time spent

The time spent in each observation for female selection and aggressive behaviour was recorded.

Statistical Analysis

Mean body weight and mean total length were analysed using one-way ANOVA (Parametric test) while mean standard length and mean caudal fin length with Kruskal-Wallis one-way ANOVA (non parametric test). Statistical significances between behavioural activities (female selection and aggressive behaviour) and time spent in each activity were analysed using one-way ANOVA followed by post hoc Tukey's tests. Data are presented as mean±SD and a probability level of 5% was used as the minimal criterion of significance. The Graph Pad Prism (version 5.01) and Paleontological Statistics (PAST, version 3.12) software were used for analysis.

Results

Maximum growth was observed in the group 'D' subjected to continuous light of 16 hours followed by groups 'C' and 'B' while minimum growth was noted in the group 'A' subjected to continuous darkness. The mean body weight (4.00 ± 0.44), total length (7.02 ± 1.66), standard length (5.31 ± 0.82) and caudal fin length (2.47 ± 0.60) were found to be maximum in group 'D' as given in Table 1.

One-way ANOVA (Parametric) revealed that the total weight ($p = 0.0002$, $F = 6.614$), total length ($p = 0.0143$, $F = 4.322$) were found to be statistically significant whereas Kruskal-Wallis one-way ANOVA (non parametric test) depicted that the standard length ($p=0.0831$, $F=2.506$) and caudal fin length ($p=0.0951$, $F=2.376$) were not significant (Table 2).

Growth Parameters

The photoperiod significantly affected the fish growth parameters (Table 3). Maximum value of specific growth rate (3.22 ± 1.38), relative growth rate (24.10 ± 5.32), absolute growth rate (0.48 ± 0.21) and weight gain ($361.53 \pm 79.83\%$) was found to be in group 'D' subjected to photoperiod of 16 L: 8 D. Highest and lowest values of feed conversion ratio were observed in the groups subjected to continuous dark ($FCR = 0.30 \pm 0.20$) and continuous light of 16 hours ($FCR = 0.14 \pm 0.09$) in the present study. The survival rate was 100 percent in three groups (B, C and D) subjected of different photoperiod regimes while 20 percent mortality was observed in the group subjected to continuous darkness.

Condition Factor

Highest value of condition factor ($K = 1.97 \pm 0.93$) was observed in the group 'A' (control)

but lowest ($K = 1.96 \pm 0.82$) in the long light phase of 16 L: 8 D (Table 3).

Welfare Indicators of the Fish

The observations of non intrusive welfare indicators on *T. fasciata* in the four different photoperiod regimes are given in Table 4. The differences in behaviour including swimming activity and different extent of injuries were noted in the various groups of *T. fasciata* subjected to different photoperiod regimes but no any colour differences were observed. The better survival, active swimming and absence of injuries were noted in the specimens of *T. fasciata* exposed to 16 hours light duration.

Behavioral Study

(A) Effect of Photoperiod on Female Selection

Females from the control group were driven to the upper part of the tank by the attack of males and females from other groups (B, C and D) in all cases

within hours of beginning of the experiment. The attack was started by males and later on by females of treated groups of different photoperiod regimes over females of control group in the present study indicated that territory was established by males and females of treated groups (B, C, D), and defended by male of the respective groups. In all the three groups "selected" female were from photoperiod influenced group but not from the control group and found to be significantly different for group 'D' ($p = 0.0043$, $F = 6.697$) (Table 5). The non selected females from control group were found to be hidden among the plants.

(B) Effects of Photoperiod on Male Aggressive Behaviour

After 19 minutes of the experiment the fish showed physical interactions. Two kinds of contact aggression were observed: bites (one fish bites another one mainly in the fins and lateral parts of the body) and mouth holding (one fish lock jaws with another fish and push against one another). A

Table 2. The results of ANOVA for the effects of different photoperiod regimes on mean body weight, mean total length, mean standard length and mean caudal fin length in *T. fasciata*.

Parameter	'p' value	'F' value
Mean body weight**	0.0002	6.614
Mean Total Body Length *	0.0143	4.322
Mean Standard Length	0.0831	2.506
Mean Caudal Fin Length	0.0951	2.376

Values with different subscript in the same row are significant ($p < 0.05$).

Values significant at the level of *less than 0.05, ** less than 0.01 and *** less than 0.001.

Table 3. Survival rate, growth parameters including feed conversion ratio, specific growth rate, relative growth rate, absolute growth rate, weight gain percentage and condition factor after 90 days under different photoperiod regimes (Mean±standard deviation).

Growth parameters	Group A	Group B	Group C	Group D
Survival (%)	80%	100%	100%	100%
FCR	0.30±0.20	0.23±0.16	0.16±0.12	0.14±0.09
SGR	1.58±0.69	1.88±0.85	2.77±1.20	3.22±1.38*
RGR	14.95±4.15	16.91±4.09	21.95±5.58	24.10±5.32*
AGR	0.23±0.10	0.28±0.128	0.41±0.1811*	0.48±0.21*
Weight gain	224.3±62.27	253.66±61.35	329.33±83.77	361.53±79.85**
K	1.97±0.93	1.968±1.30	1.967±0.70	1.96±0.82*

Values with different subscript in the same row are significantly different ($p < 0.05$).

Significant level (* less than 0.05, ** less than 0.01 and *** less than 0.001).

Table 4. Welfare assessment of *T. fasciata* under different photoperiod regimes.

Welfare indicators	Group A	Group B	Group C	Group D
Injuries	Found on the body and fins	Found on the body	None	None
Colour	Normal	Normal	Normal	Normal
Swimming activity	Suppressed	Active	Active	Very active

Group A = 0 L: 16 D, Group B = 8 L: 16 D, Group C = 12 L: 12 D and Group D = 16 L: 8 D

photoperiod induced males from groups B, C and D dominated in all the three cases and performed the first attack biting the males of control group. A stronger territorial and aggressive behaviour was observed in males of treated groups towards control males ($p=0.0046$, $F=6.608$) (Table 5). Behavioural inhibition was observed including decreased locomotors activity in males of control group found to be hidden behind the stone or in the top surface of the aquaria. Number of observations in female attacks and aggressive interactions were counted and given in Figure 1.

(C) Time Spent

The time spent was found to be significantly different for female selection ($p = 0.0163$, $F = 4.811$) and male aggressive interactions ($p = 0.0433$, $F = 3.660$). The details are given in Figure 2 and Figure 3.

Discussion

The findings of the present study showed that the growth of giant gourami, *Trichogaster fasciata*

were affected by different photoperiod regimes, and maximum growth was recorded in the groups subjected to 16 L: 8 D photoperiod regime. The photoperiod regime of 16 L: 8 D hours was considered to be optimum for proper growth and survival in giant gourami because of increased feed intake and better feed conversion ratio. The present finding was similar to the work carried out by Tandler and Helps (1985); Folkvord and Ottera (1993); Imsland et al. (1995); Duncan et al. (1999); Ergün et al. (2003); Rad et al. (2006); Yager and Yigit (2009) and Kashyap et al. (2015) in various fish species, their reports also suggested that the fishes subjected to continuous photoperiod obtained highest growth. A number of workers such as Saunders and Henderson (1970); Lundqvist (1980); Brauer (1982); Saunders et al. (1985) reported that artificially increased day light is the good source for the enhancement of growth and development of fishes. In contrary to this, Arvedlund et al. (2000) reported slow growth under 24 L: 0 D as compared to 16 L: 8 D photoperiod regime in *Amphiprion melanopus* and emphasized that their developing juveniles had a period of inactivity during darkness and their growth was compromised. A

Table 5. Behavioural activity and time spent of *T. fasciata* between control vs different experimental trials. Values in the same row marked with different letters are significantly different (ANOVA results, Tukey's post hoc test, $p < 0.05$).

S.No	Behavioural activity	p value	F value
Exp.1	Female selection: Total number of attacks experienced by (control) female	0.0043 c***, b**	6.697
	Time spent in Female attacks	0.0163 c*	4.811
Exp.2	Total number of male aggressive interactions between photoperiod influenced groups towards the male (control)	0.0046 c*	6.608
	Time spent in male aggressive interactions	0.0433 c*** b**	3.660

(Group A = Control), (Group B = 8L: 16D), (Group C = 12L: 12D), (Group D = 16L: 8D)

a = (A vs B) vs (A vs C)

b = (A vs B) vs (A vs D)

c = (A vs C) vs (A vs D)

Values significant at the level of * less than 0.05, ** less than 0.01 and *** less than 0.001.




Figure 1. Total number of female attacks and aggressive interactions between control and three different experimental groups.




Figure 2. Time spent in each observation of female selection activity of 1 hour duration.




Figure 3. Time spent in each observation of male aggressive interactions in 1 hour duration.

numbers of workers such as Fuchs (1978) and Barahona-Fernandes (1979) reported different optimum photoperiod regimes for different fish species.

The requirement of photoperiod regime is considered to be species specific and play an important role for growth and survival of species (Britz & Pienaar, 1992; Silva-Garcia, 1996; Boeuf & Baille, 1999). The details of effect of photoperiod regimes on growth, survival and mortality are given in Figure 4. The less (2.80 g) and minimum (2.53 g) growth were recorded in the groups exposed to 8 L: 16 D hours photoperiod regime and complete darkness respectively in the current study. In contrary to this, the juvenile of *Clarius garpines* showed the high growth rate when reared under continuous darkness (Britz & Pienaar, 1992). Yamamoto et al. (2001) reported that shortening of light phase during the feeding trial affected feed intake of carp.

The differences in initial and pronounced growth were noticed after 45th and 60th day onwards of the experiment in the fish subjected to different photoperiod regimes. The same results were recorded in the case of AGR, SGR, RGR and weight gain

which suggested that the juveniles of *T. fasciata* required several weeks to acclimatize to new rearing conditions. The similar results were reported by Kashyap et al. (2015) in *Catla catla*. Minimum value of FCR (0.139 ± 0.090) was recorded in the group subjected to maximum duration of photoperiod which is similar to the finding of Rad et al. (2006) who pointed out that the duration of hours is inversely proportional to feed conversion (FCR) resulted into better growth. Sixteen hours duration of photoperiod was found to be suitable for the growth of giant gourami in the present study. Similar duration (16 L: 8 D) of photoperiod was reported by Gines et al. (2004) for maximum growth in Gilthead sea bream.

Condition factor is an important biological parameter which indicated the well being of fish in a specific water body where the fish grows. Moreover, it is an index which indicates the physiological state of the fish and shows population welfare during the various stages of the life cycle. The value of condition factor was found to be lowest ($K = 1.9635 \pm 0.8203$) in the group subjected to 16 L: 8 D photoperiod regime which was similar to the finding of Elsbaay (2013) who reported the lowest value of condition




Figure 4. The effects of different photoperiod regimes on growth, survival and mortality of fish.

factor ($K = 2.21$) in long light phase (24 L: 0 D) while estimated the effect of photoperiod on growth rate of Nile Tilapia.

Herbert and Steffensen (2005) pointed out that the status of activity can be used as an indicator of poor or good welfare in fishes. The group 'D' of *T. fasciata* in present study was found to be very active as indicated by the swimming activity of the specimens which reflected that how a fish is sensing and responding to its environment. Maximum injuries were noted in the control group while no injuries were found in the groups subjected to 12 and 16 hours light duration. Almaza'n-Rueda et al. (2004, 2005) also validated the use of skin lesion frequency to assess the welfare of African catfish and showed that feeding methods, photoperiod and light intensity affected skin lesions frequency as well as other welfare indicators such as swimming activity, growth, plasma cortisol and free fatty acids. Continuous darkness (0 L: 24D) was found to be adverse effect on the welfare of *T. fasciata* which had resulted into the mortality of two individuals of fish. Chronic stress may be responsible for welfare impairment while acute stress might be a reason for the mortality of two fish in the present study as reported by Pickering (1998) and Damsgard et al. (2006) in their research works. The unfavourable photoperiods could have affected the immune systems of the species resulted in the observed deviations in the normal welfare behaviour of the giant gourami as reported by Mustapha et al. (2014). Pottinger and Pickering (1992) and Burgos et al. (2004) reported that artificial photoperiods affected the immune system of rainbow trout leading to mortality. The normal welfare condition was found to be in the photoperiod regime of 16 L: 8 D in *T. fasciata* in the present study which could be due to their innate behaviour being displayed.

The females "selected" by males in the present study were from photoperiod treated groups (B, C and D) than control (group A). The selection of a territory was found to be a characteristic behaviour in giant gourami as shown by the aggressive interaction between males from control group with other groups

subjected to different photoperiod regimes led to the selection of a territory. Moreover, in aggressive interaction males of control were found to be hidden in order to escape themselves from the dominating ones. The present findings was found to be similar to that of the work reported by Fiszbein et al. (2010) where behavioural level of aggression related to territory selection and defence was reduced in control group, and females exposed to control were never chosen by males. Winners were those that keep most of the territory and subordinates (control) were exploited by the winners. A subordinate position within a group or social hierarchy may be a stressor (Schreck, 1981) which can be subjected to social stress resulting from the attacks of the more dominant individuals for access to resources (sexual partners and territory). Social stress caused to marked behavioural and physiological changes in subordinates, who often showed a general behavioural inhibition of food intake, locomotory activity and aggressiveness (Denight & Ward, 1982; Winberg & Nilsson, 1993a and O'Connor et al., 1999).

Acknowledgements

The authors thank the Head, Department of Zoology, University of Lucknow for providing facility and administrative support. One of the authors, Miss Farah Bano thanks the University Grants Commissions, New Delhi for the BSR fellowship, which was given to undertake advanced studies and research leading to Ph.D. degrees in sciences.

Funding: This work was carried out without any funding.

References

- Almaza'n-Rueda, P., Schrama, J.W., & Verreth, J.A.J. (2004). Behavioural Responses Under Different Feeding Methods and Light Regimes of the African Catfish (*Clarias gariepinus*) Juveniles. *Aquaculture*, 231, 347-359.

- <http://dx.doi.org/10.1016/j.aquaculture.2003.11.016>.
- Almaza'n-Rueda, P., Van-Helmond A.T.M., Verreth, J.A.J., & Schrama, J.W. (2005). Photoperiod Affects Growth, Behaviour and Stress Variables in *Clarias gariepinus*. *Journal of Fish Biology*, 67, 1029-1039. <http://dx.doi.org/10.1111/j.0022-1112.2005.00806.x>.
- Arvedlund, M., McCormick, M.I., & Ainsworth, T. (2000). Effects of Photoperiod on Growth of Larvae and Juveniles of the Anemonefish, *Amphiprion melanopus*. *Naga ICLARMQ*, 23, 18-23.
- Askarian, F., & Kousha, A. (2009). The Influence of Photoperiod on Farming of Beluga Sturgeon (*Huso huso*): Evaluation by Growth and Health Parameters in Serum. *Journal of Fisheries and Aquatic Sciences*, 4, 41-49. <http://dx.doi.org/10.3923/jfas.2009.41.49>.
- Barahona-Fernandes, M.H. (1979). Some Effects of Light Intensity and Photoperiod on the Seabass Larvae (*Dicentrarchus labrax*) Reared at the Centre Oceanologique de Bretagne. *Aquaculture*, 17, 311-321. [http://dx.doi.org/10.1016/0044-8486\(79\)900863](http://dx.doi.org/10.1016/0044-8486(79)900863).
- Bhattacharyya, S., Dey, R., & Maitra, S.K. (2005). Photoperiodic Regulation of Annual Testicular Events in the Indian Major Carp, *Catla catla*. *Acta Zoologica*, 86, 71-79. <http://dx.doi.org/10.1111/j.1463-6395.2005.00188.x>.
- Bjornsson, B.T., Thorarensen, H., Hirano, T., Ogasawara, T., & Kristinsson, J.B. (1989). Photoperiod and Temperature Affect Plasma Growth Hormone Levels, Growth, Condition Factor and Hypo Osmoregulatory Ability of Juvenile Atlantic Salmon (*Salmo salar*) during Parr-Smolt Transformation. *Aquaculture*, 82, 77-91. [http://dx.doi.org/10.1016/0044-8486\(89\)90397-9](http://dx.doi.org/10.1016/0044-8486(89)90397-9).
- Boeuf, G., & Baille, P.Y. (1999). Does Light have an Influence on Fish Growth? *Aquaculture*, 177, 129-152. [http://dx.doi.org/10.1016/S0044-8486\(99\)00074-5](http://dx.doi.org/10.1016/S0044-8486(99)00074-5).
- Bonnet, E., Montfort, J., Esquerre, D., Hugot, K., Fostier, A., & Bobe, J. (2007). Effect of Photoperiod Manipulation on Rainbow Trout (*Oncorhynchus mykiss*) Egg Quality: A Genomic Study. *Aquaculture*, 268, 13-22. <http://dx.doi.org/10.1016/j.aquaculture.2007.04.027>.
- Boujard, T., & Leatherland, J.F. (1992). Circadian Rhythms and Feeding Time in Fishes. *Environmental Biology of Fishes*, 35, 109-131. <http://dx.doi.org/10.1007/BF00002186>.
- Brauer, E.P. (1982). The Photoperiod Control of Coho Salmon Smoltification. *Aquaculture*, 28, 105-111. [http://dx.doi.org/10.1016/0044-8486\(82\)90013-8](http://dx.doi.org/10.1016/0044-8486(82)90013-8).
- Britz, G.P.J., & Pienaar, A. G. (1992). Laboratory Experiments on the Effect of Light and Cover on the Behavior and Growth of African Catfish, *Clarias gariepinus* (Pisces: Claridae). *Journal of Zoology*, 227, 43-62. <http://dx.doi.org/10.1023/A:1009299830102>.
- Burgos, A., Valenzuela, A., Gonzalez, M., & Klempau, A. (2004). Non-Specific Defence Mechanisms of Rainbow Trout (*Oncorhynchus mykiss*) during Artificial Photoperiod. *Bulletin of European Association Fish Pathology*, 24, 240-245.
- Damsgard, B., Juell, J.E., & Braastad, B.O. (2006). Fiskeriforskning. Welfare in Farmed Fish (Report No. 5). Bergen, Norway.
- Davies, P.R., & Hanyu, I. (1986). Effect of Temperature and Photoperiod on Sexual Maturation and Spawning of the Common Carp: I. Under Conditions of High Temperature. *Aquaculture*, 51, 277-288. [http://dx.doi.org/10.1016/0044-8486\(86\)90319-4](http://dx.doi.org/10.1016/0044-8486(86)90319-4).
- Davies, P.R., Hanyu, I., Furukawa, K., & Nomura, M. (1986a). Effect of Temperature and Photoperiod on Sexual Maturation and Spawning of the Common Carp: III. Induction of Spawning by Manipulating Photoperiod and Temperature. *Aquaculture*, 52, 137-144. [http://dx.doi.org/10.1016/0044-8486\(86\)90033-5](http://dx.doi.org/10.1016/0044-8486(86)90033-5).
- Davies, P.R., Hanyu, I., Furukawa, K., & Nomura, M. (1986b). Effect of Temperature and Photoperiod on Sexual Maturation and Spawning of the Common Carp: II. Under Conditions of Low Temperature. *Aquaculture*, 52, 51-58. [http://dx.doi.org/10.1016/0044-8486\(86\)90107-9](http://dx.doi.org/10.1016/0044-8486(86)90107-9).
- Degani, G., Ben-Zvi, Y., & Levanon, D. (1989). The Effect of Different Protein Levels and Temperatures on Feed Utilization, Growth and Body Composition of *Claris gariepinus* (Burchell, 1822). *Aquaculture*, 76, 293-301. [http://dx.doi.org/10.1016/0044-8486\(89\)90082-3](http://dx.doi.org/10.1016/0044-8486(89)90082-3).
- Denight, M.L., & Ward, J.A. (1982). Relationship of Chin Spot Size to Dominance in the black-Chinned Mouthbrooding Cichlid Fish (*Sarotherodon melanotheron*). *Animal Behaviour* 30, 1099-1104. [http://dx.doi.org/10.1016/S0003-3472\(82\)80200-5](http://dx.doi.org/10.1016/S0003-3472(82)80200-5).
- Donaldson, E.M., Fagerlund, H.M., Higgs, D.A., & MacBride, J.R. (1979). Fish Physiology. In Hoar, W.S., Randall, D.J., Brett, J.R. (Eds.), *Hormonal Enhancement of Growth*. New York Academic Press., 455 pp.
- Duncan, N., Mitchell, D., & Bromage, N.R. (1999). Post-smolts Growth and Maturation of Out-Season 0+ Atlantic Salmon (*Salmo salar*) Reared Under Different Photoperiods. *Aquaculture*, 177, 61-71. [http://dx.doi.org/10.1016/S0044-8486\(99\)00069-1](http://dx.doi.org/10.1016/S0044-8486(99)00069-1).
- Elsbaay, A.M. (2013). Effects of Photoperiod and Different Artificial Light Colors on Nile Tilapia Growth Rate. *Journal of Agriculture and Veterinary Science*, 3, 5-12.
- Ergün, S., Yigit, M., & Türker, A. (2003). Growth and Feed Consumption of Young Rainbow Trout (*Oncorhynchus mykiss*) Exposed to Different Photoperiods. *Israeli Journal of Aquaculture*, 55, 132-138. <http://dx.doi.org/35400011858355.0070>.
- Falcon, J., Migaud, H., Munoz-Cueto, J.A., & Carrillo, M. (2010). Current Knowledge on the Melatonin System in Teleost Fish. *General and Comparative Endocrinology*, 165, 469-482. <http://dx.doi.org/10.1016/j.ygcen.2009.04.026>.
- Fiszbein, A., Canepa, M., Vazquez, G.R., Maggese, C., & Pandolfi, M. (2010). Photoperiodic Modulation of Reproductive Physiology and Behaviour in the Cichlid Fish (*Cichlasoma dimerus*). *Physiology & Behaviour*, 99, 425-432. <http://dx.doi.org/10.1016/j.physbeh.2009.11.017>.
- Folkvord, A., & Ottera, H. (1993). Effects of Initial Size Distribution, Day Length and Feeding Frequency on Growth, Survival and Cannibalism in Juvenile Atlantic Cod (*Gadus morhua* L.). *Aquaculture*, 114, 243-260. <http://dx.doi.org/35400003539849.0060>.
- Fuchs, J. (1978). Effect of Photoperiod on Growth and Survival during Rearing of Larvae and Juveniles of Sole (*Solea solea*). *Aquaculture*, 15, 63-74. [http://dx.doi.org/10.1016/0044-8486\(78\)90072-8](http://dx.doi.org/10.1016/0044-8486(78)90072-8).
- Gines, R., Afonso, J.M., Arguello, A., Zamorano, M.J., & Lopez, J.L. (2004). The Effects of Long-Day

- Photoperiod on Growth, Body Composition and Skin Colour in Immature Gilthead Sea Bream (*Sparus aurata* L.). *Aquaculture Research*, 35, 1207-1212. <http://dx.doi.org/10.1111/j.1365-2109.2004.01126.x>
- Herbert, N.A., & Steffensen, J.F. (2005). The Response of Atlantic cod, *Gadus morhua*, to Progressive Hypoxia: Fish Swimming Speed and Physiological Stress. *Marine Biology*, 147, 1403-1412. <http://dx.doi.org/10.1007/s00227-005-0003-8>.
- Imsland, A., Folkvord, A.F., & Stefansson, S.O. (1995). Growth Oxygen Consumption and Activity of Juvenile Turbot *Scophthalmus maximus* (L.) reared Under Different Temperatures and Photoperiods. *Netherland Journal of Sea Research*, 34, 149-159. [http://dx.doi.org/10.1016/0077-7579\(95\)90023-3](http://dx.doi.org/10.1016/0077-7579(95)90023-3).
- Kashyap, A., Pathak, B.C., Awasthi, M., & Serajuddin, M. (2015). Effect of Different Photoperiods on the Growth and Survival of Juvenile of Indian Major Carp, *Catla catla*. *Iranian Journal of Fisheries Sciences*, 14, 946-955.
- Lundqvist, H. (1980). Influence of Photoperiod on Growth in Baltic Salmon Parr (*Salmo solar*) with Special Reference to the Effect of Precocious Sexual Maturation. *Canadian Journal of Zoology*, 58, 940-944. <http://dx.doi.org/10.1139/z80-132>.
- Martins, C.I.M., Galhardo, L., Noble, C., Damsgard, B., Spedicato, M.T., Zupa, W., Beauchaud, M., Kulczykowska, E., Massabuau, J.C., Carter, T., Planellas S.R., & Kristiansen, T. (2012). Behavioural Indicators of Welfare in Farmed Fish. *Fish Physiology and Biochemistry*, 38, 17-41. <http://dx.doi.org/10.1007/s10695-011-9518-8>.
- Moyle, P.B., & Cech, J.J. (2000). Feeding, Nutrition, Digestion and Excretion; Growth, In Ryu, T., Tarabokjia, L. (Eds), *Fishes: An Introduction to Ichthyology*. Prentice Hall, New Jersey, 97 pp.
- Mustapha, M.K., Oladokun, O.T., Salman, M.M., Adeniyi, I.A., & Ojo, D. (2014). Does Light Duration (Photoperiod) have an Effect on the Mortality and Welfare of Cultured *Oreochromis niloticus* and *Clarias gariepinus*? *Turkish Journal of Zoology*, 38, 466-470. <http://dx.doi.org/10.3906/zoo-1309-23>.
- O'Connor, K.I., Metcalfe, N.B., & Taylor, A.C. (1999). Does Darkening Signal Submission in Territorial Contests between Juvenile Atlantic Salmon, *Salmo salar*? *Animal Behaviour*, 58, 1269-1276. <http://dx.doi.org/10.1006/anbe.1999.1260>.
- Pickering, A.D. (1998). Stress Responses in Farmed Fish. In Black KD, Pickering AD (Eds.), *Biology of Farmed Fish*. Sheffield Academic Press, 222 pp. <http://dx.doi.org/10.1080/00288330.2013.808236>
- Pottinger, T.G., & Pickering, A.D. (1992). The Influence of Social Interaction on the Acclimation of Rainbow Trout, *Oncorhynchus mykiss* (Walbaum) to chronic stress. *Journal of Fish Biology*, 41, 435-447. <http://dx.doi.org/10.1111/j.1095-8649.1992.tb02672.x>.
- Purchase, C.F., Boyce, D.L., & Brown, J.A. (2000). Growth and Survival of Juvenile Yellowtail Flounder, *Pleuronectes ferrugineus* (Storer) Under Different Photoperiods. *Aquaculture Research*, 31, 547-552. <http://dx.doi.org/10.1046/j.1365-2109.2000.00480.x>.
- Rad, F., Bozaoğlu, S., Gözükar, S.E., Karahan, A., & Kurt, G. (2006). Effects of Different Long-Day Photoperiods on Somatic Growth and Gonadal Development in Nile Tilapia (*Oreochromis niloticus*). *Aquaculture*, 255, 292-300. <http://dx.doi.org/10.1016/j.aquaculture.2005.11.028>.
- Reinecke, M. (2010). Influences of the Environment on the Endocrine and Paracrine Fish Growth Hormone-Insulin-Like Growth Factor-I System. *Journal of Fish Biology*, 76, 1233-1254. <http://dx.doi.org/10.1111/j.1095-8649.2010.02605.x>.
- Ricker, W.E. (1973). Linear Regressions in Fishery Research. *Journal of the Fisheries Board of Canada*, 30(3), 409-434. <http://dx.doi.org/10.1139/f73-072>.
- Rousing, T., Bonde, M., & Sørensen, J.T. (2001). Aggregating Welfare Indicators into an Operational Welfare Assessment System: A Bottom Up Approach. *Acta Agriculturae Scandinavica, Section A- Animal Science*, 30, 53-58. <http://dx.doi.org/10.1080/090647001300004790>.
- Ruchin, A.B. (2004). Influence of Colored Light on Growth Rate of Juveniles of Fish. *Fish Physiology and Biochemistry*, 30, 175-178. <http://dx.doi.org/10.1007/s10695-005-1263-4>.
- Ruchin, A.B. (2006). Effect of Light on White Blood Cell Count in Carp (*Cyprinus carpio*). *Biology Bulletin*, 33, 517-520. <http://dx.doi.org/10.1134/S1062359006050153>.
- Saunders, R.L., & Henderson, E.B. (1970). Influence of Photoperiod on Smolt Development and Growth of Atlantic Salmon (*Salmo solar*). *Journal of Fisheries Research Board of Canada*, 27, 1295-1311. <http://dx.doi.org/10.1139/f70-151>.
- Saunders, R.L., Henderson, E.B., & Harmon, P.R. (1985). Effects of Photoperiod on Juvenile Growth and Smolting of Atlantic Salmon and Subsequent Survival and Growth in Sea Cages. *Aquaculture*, 45, 55-66. [http://dx.doi.org/10.1016/0044-8486\(85\)90257-1](http://dx.doi.org/10.1016/0044-8486(85)90257-1).
- Schreck, C.B. (1981). Stress and Compensation in Teleostean Fishes: Response to Social and Physical Factors. In Pickering AD (Eds.) *Stress and Fish*. Academic Press, New York, 294 pp.
- Silva-Garcia, A.J. (1996). Growth of Juvenile Gilthead Seabream (*Sparus aurata* L.) Reared Under Different Photoperiod Regimes. *Israeli Journal of Aquaculture*, 48, 84- 93.
- Simensen, L.M., Jonassen, T.M., Imsland, A.K., & Stefansson, S.O. (2000). Photoperiod Regulation of Growth Juvenile Atlantic Halibut (*Hippoglossus hoppeglossus* L.). *Aquaculture*, 190, 119-128. [http://dx.doi.org/10.1016/S0044-8486\(00\)00397-5](http://dx.doi.org/10.1016/S0044-8486(00)00397-5).
- Tandler, A., & Helps, S. (1985). The Effects of Photoperiod and Water Exchange Rate on Growth and Survival of Gilthead Sea Bream (*Sparus aurata*, L.) from Hatching to Metamorphosis in Mass Rearing Systems. *Aquaculture*, 48, 71-82. [http://dx.doi.org/10.1016/0044-8486\(85\)90053-5](http://dx.doi.org/10.1016/0044-8486(85)90053-5).
- Taylor, J.F., North, B.P., Porter, M.J.R., Bromage, N.R., & Migaud, H. (2006). Photoperiod Can Be Used to Enhance Growth and Improve Feeding Efficiency in Farmed Rainbow Trout, (*Oncorhynchus mykiss*). *Aquaculture*, 256, 216-234. <http://dx.doi.org/10.1016/j.aquaculture.2006.02.027>.
- Trzebiatowski, R., Filipiak, J., & Jakubowski, R. (1981). Effect of Stocking Density on Growth and Survival of Rainbow Trout (*Salmo gairdneri* Rich.). *Aquaculture*, 22, 289-295. [http://dx.doi.org/10.1016/0044-8486\(81\)90155-1](http://dx.doi.org/10.1016/0044-8486(81)90155-1).
- Valenzuela, A.E., Silva, V.M., & Klempau, A.E. (2006). Qualitative and Quantitative Effects of Constant Light Photoperiod on Rainbow Trout (*Oncorhynchus mykiss*) Peripheral Blood Erythrocytes. *Aquaculture*, 255, 292-300.

- 251, 596-602.
<http://dx.doi.org/10.1016/j.aquaculture.2005.06.012>.
- Winberg, S., & Nilsson, G.E. (1993a). Roles of Brain Monoamine Neurotransmitters in Agonistic Behaviour and Stress Reactions, with Particular Reference to Fish. *Comparative Biochemistry and Physiology*, 106, 597-614.
[http://dx.doi.org/10.1016/0742-8413\(93\)90216-8](http://dx.doi.org/10.1016/0742-8413(93)90216-8).
- Woiwode, J.G., & Adelman, I.R. (1991). Effects of Temperature, Photoperiod, and Ration Size on Growth of Hybride Striped Bass X White Bass. *Transactions of the American Fisheries Society*, 120, 217-222.
[http://dx.doi.org/10.1577/15488659\(1991\)120<0217:EOTPAR>2.3.CO](http://dx.doi.org/10.1577/15488659(1991)120<0217:EOTPAR>2.3.CO).
- Yager, D.D., & Yigit, M. (2009). Influence of Increased Photoperiods on Growth, Feed Consumption and Survival of Juvenile Mirror Carp (*Cyprinus carpio* L., 1758). *Journal of Fisheries Sciences*, 3, 146-152.
<http://dx.doi.org/10.3153/jfscm.2009018>.
- Yamamoto, T., Shima, T., Furuita, H., Shiraishi, M., Sánchez-Vázquez, F.J., & Tabata, M., (2001). Influence of Decreasing Water Temperature and Shortening of the Light Phase on Macronutrient Self-Selection by Rainbow Trout (*Oncorhynchus mykiss*) and Common Carp (*Cyprinus carpio*). *Fisheries Science*, 67, 420-429.
<http://dx.doi.org/10.1046/j.1444-2906.2001.00260.x>.