



Metazoan Parasite Fauna of the Red Mullet, *Mullus barbatus ponticus* Essipov, 1927 in the Sinop Coasts of the Black Sea

Türkay Öztürk^{1,*}, Ali Yesil¹

¹ Sinop University, Faculty of Fisheries and Aquatic Sciences, Sinop, Turkey.

* Corresponding Author: Tel.: +90.368 2876265; Fax: +90.368 2876268;
E-mail: turkay.ozturk@gmail.com

Received 12 April 2017
Accepted 29 May 2017

Abstract

In the present study, a total of 330 specimens the red mullet, *Mullus barbatus ponticus* caught in the Black Sea coasts near Sinop, Turkey were investigated throughout a 1-year period for their metazoan parasite fauna. Thirteen metazoan parasite species, *Loma* sp., *Proctotrema bacillioatum*, *Galactosomum lacteum* (metacercariae), *Progrillotia dasyatidis*, *Nybelinia* sp., *Scolex pleuronectis* (pleurocercoid), *Hysterothylacium fabri*, *H. aduncum*, *Ascarophis valentina*, *Capillaria* sp. *Contraecum* sp., unidentified larval nematode (L2) and *Acanthocephaloides irregularis* were identified. The overall infection prevalence and mean intensity values were 97.88% and 34.12 ± 5.23 parasites per infected fish, respectively. *Proctotrema bacillioatum* was the core species with infection prevalence of 70.61% and mean intensity value of 29.09 ± 7.01 parasites per infected fish. The infection indice values for each parasite species were also determined and discussed according to season, length classes and sex of fish. This paper is the first report on metazoan parasite fauna of the red mullet in Turkey.

Keywords: *Mullus barbatus ponticus*, metazoan parasites, Black Sea.

Introduction

The family Mullidae is a commercially important fish group which can be seen throughout the world's seas. They are demersal fishes in coastal water, including estuaries, over sandy or muddy bottoms of temperate and tropical region. The genus *Mullus* is represented by two species (*M. barbatus* L., 1758 and *M. surmuletus* L., 1758) and one subspecies (*M. barbatus ponticus* Essipov, 1927). *Mullus barbatus* and *M. surmuletus* are distributed in the Mediterranean Sea, the eastern North Atlantic Ocean and the Black Sea. *Mullus barbatus ponticus* Essipov, 1927 is taxonomically classified as a subspecies of *Mullus barbatus* in the Black Sea (Hureau, 1986; Sahin & Akbulut, 1997; Turan, 2006). Since the red mullet is both economically valuable and an export product of Turkish fisheries, it is among the important demersal fish of the Black Sea. In spite of their ecological and economical importance in the region, the research on the parasitic fauna and parasite ecology of the fish are still not enough. There are many parasitological studies on *Mullus barbatus* from the Mediterranean and Atlantic Seas (Essafi, Raibaut & Boudaoud-Krissat, 1983; Abo-Esa, 2007; Paradiznik & Radujkovic, 2007; Ramdane & Trilles,

2010; Carreras-Aubets, Repulles-Albelda, Kostadinova & Carrassón, 2011; Carreras-Aubets, Montero, Kostadinova & Carrassón, 2012; Debenedetti, Madrid & Fuentes, 2013; Radujkovic & Sundic, 2014). On the other hand, studies on the parasite fauna of *Mullus barbatus* or *M. barbatus ponticus* in Black Sea are limited (Ovcharenko & Yurahno, 2006; Gaevskaya, 2012).

The aim of the present study is to reveal the parasite fauna of the red mullet, their infection prevalences and intensities in relation to season and length classes as well as sex of fish. This study is the first to provide data on metazoan parasites and their infection indices in red mullet collected from the Turkish Black Sea coast.

Materials and Methods

Fish specimens were collected from commercial fishing vessels off Sinop coast in the Black Sea in Turkey (42°00'05" N, 35°08'59" E). Samplings were carried out monthly throughout a one-year period. A total of 330 fish specimens were investigated for parasites. At necropsy, total length and sex were recorded for each individual. The length of fish varied between 8.0 and 21.0 cm during whole sampling

period. Seasonally, the average length of the fish was recorded as Lsp: 14.2 cm in spring, Lsu: 13.5 cm in summer, Lau: 13.9 cm in autumn, Lwi: 14.5 in winter. In addition fish were grouped according to the following size classes after total lengths had been measured: < 13 cm (up to 13.0), 13-15 cm (13.0-14.9), and > 15 cm (15.0-21.0). The three size classes were chosen according to the approximate age of the species suggested by Aydın and Karadurmuş (2013). External (gills, skin, fins, eyes) and internal organs (heart, liver, kidney, gall bladder, muscle) of fishes were examined for parasites and the number of parasites was counted individually and the site of infection was recorded.

Fish were then examined for parasites using standard methods. Digenean parasites were fixed in Bouin's solution, nematodes were fixed in hot (70-80 °C) 70% ethanol, cleared in Amann lactophenol, acanthocephalans and cestode were fixed in alcohol 70% and studied by direct examination between slide and cover slip. The parasites were identified based on morphological criteria. Parasite preparations were conducted according to methods indicated by Bray, Gibson and Jones (2008) for digeneans, Petter, Lebre and Radujkovic (1984); Ferrer *et al.* (2005) for nematodes, Beveridge, Neifar and Euzet (2004) for cestodes, Amin, Oğuz, Heckmann, Tepe and Kvach (2011) for acanthocephala. Morphological diagnostic features of parasite specimens were identified using light microscope (Olympus BX53) fitted with digital camera attachment (DP50).

The prevalence and mean abundance values of *Hysterothylacium aduncum* and *H. fabri* were given as *Hysterothylacium* spp. for pooled data rather than by each *Hysterothylacium* species.

The prevalence, mean intensity, abundance and intensity range (minimum and maximum parasites number) were determined according to Bush, Lafferty, Lotz and Shostak (1997). The standard error (SE) for mean intensity and abundance is given. While the differences in the mean intensities values of each

parasite species for length classes of fish and the seasons was tested by the Kruskal-Wallis test (nonparametric ANOVA), the difference between parasite loads in male and female fish was tested by the Mann-Whitney U test. Chi square was used on the prevalence of parasites. The analyses were carried out using the computer programme GraphPad InStat 3.0 and "Minitab 16 statistical software. The significance level was $P < 0.05$.

Results

A total of 13 metazoan parasite species were found (Table 1). These are; *Loma* sp. (Figure 1), *Proctotrema bacilliovatum*, *Galactosomum lacteum* (metacercariae) (Figure 2), *Progrillotia dasytidis*, *Nybelinia* sp., *Scolex pleuronectis* (pleurocercoid) (Figure 3), *Hysterothylacium fabri*, *H. aduncum*, *A. valentina*, *Capillaria* sp., *Contracaecum* sp. (Figure 4), an unidentified larval nematode (L2) and *Acanthocephaloides irregularis* (Figure 5). Table 1 illustrates identified parasite list with indications of prevalence, mean intensity, site of infection (microhabitat) and parasite range. No parasite species was detected in the muscle of examined fish. The infection prevalence (%) and mean intensity values were 97.88% and 34.12 ± 5.23 parasites per infected fish, respectively. Parasite component communities in *M. barbatus ponticus* comprised 13 parasite species; of these, 6 were rare by having prevalence values less than 10% in the entire dataset (Table 1).

Seasonal prevalence and mean intensity of all parasite species are reported in Table 2. The overall mean intensity values of parasite species on *M. barbatus ponticus* varied significantly among the seasons ($P < 0.05$) and the highest mean intensity occurred in summer. *Proctotrema bacilliovatum*, *P. dasytidis*, *Hysterothylacium* spp. (*H. fabri*, *H. aduncum*), *A. valentina*, *Capillaria* sp., *Contracaecum* sp., and an unidentified larval nematode (L2) were recorded in fish at all seasons. The mean intensity




Figure 1. Photomicrographs of *Loma* sp. **A.** spores of *Loma* sp, fixed in Bouin's fixative; **B.** xenomas of *Loma* sp.; **C.** part of the intestine wall containing cysts of *Loma* sp. (young and mature xenomas), (original).




Figure 2. Photomicrographs of two digenean species. **A.** *Proctotrema bacilliovatum*, Odhner, 1911. **B.** *Galactosomum lacteum* (Jagerskiold, 1896) excysted metacercaria, (Os: oral sucker; p; pharynx, Ov, ovary, Vs: ventral sucker, t, testis, Cs; cirrus sac) (fresh material, original).




Figure 3. Photomicrographs of the three cestode species: **A.** *Progrillotia dasyatidis* Beveridge Neifar & Euzet, 2004; **B.** tentacles of *P. dasyatidis*, **C.** *Nybelinia* sp., **D.** tentacle of *Nybelinia* sp. **E.** *Scolex pleuronectis* (plerocercoid), **F.** bothrial margin of *S. pleuronectis* (original).

values of *P. bacilliovatum*, *P. dasyatidis*, *Hysterothylacium* spp., *Capillaria* sp., *Contracaecum* sp., and an unidentified larval nematode (L2) varied significantly depending on the seasons ($P < 0.05$, Table 2).

The infection parameters of parasite species in three length classes of fish are given in Table 3. *Loma* sp., *G. lacteum*, *Nybelinia* sp. and *A. irregularis* were

found only in some length classes of fish, *Loma* sp. in the middle and large and *Nybelinia* sp. only in the largest. Although large fish had higher mean intensity value than did small fish, the difference among the fish length classes was statistically insignificant ($P > 0.05$, Table 4).

Both overall infection prevalence and mean intensity values were higher in females than in males




Figure 4. Photomicrographs of the four nematode species. **A.** anterior end of *Hysterothylacium fabri* (Rudolphi, 1819), **B.** anterior end of *H. aduncum* Rudolphi, 1802, **C.** *Contracaecum* sp. (encysted), **D.** tail (female) of *Ascarophis valentina* Ferrer *et al.*, 2005, **E.** tail of (male) of *A. valentina*, **F.** gravid female of *A. valentina*, **G.** tail (female) of *Capillaria* sp., **H.** tail (male) of *Capillaria* sp., **I.** developed egg of *Capillaria* sp. (original).




Figure 5. Photomicrographs of *Acanthocephaloides irregularis* Amin, Oğuz, Heckmann, Tepe & Kvach, 2011. **A.** female specimen, **B.** male specimen, **C.** anterior regio, **D.** Proboskis, **E.** posterior regio (original).

(Table 4). However, the difference between the mean values was not statistically significant ($P>0.05$).

The only microsporidian parasite detected was *Loma* sp. in the rectum and intestine of fish with a prevalence of 0.91%. The digeneans, *Proctotrema bacilliovatum* and *Galactosomum lacteum* occurred in the digestive tract as adults and metacercaria, respectively. *Proctotrema bacilliovatum* was the most abundant and had 70.61% infection prevalence and mean intensity values of 29.09 ± 7.01 parasites per infected fish. On the contrary, *Galactosomum lacteum* was the least (Table 1). *Progrillotia dasyatidis*, *Nybelina* sp. and *Scolex pleuronectis* cestodes were found in the gall bladder, stomach and intestine.

Six nematode species, *Hysterothylacium aduncum*, *H. fabri*, *Ascarophis valentina*, *Capillaria* sp., *Contracaecum* sp. and unidentified larval nematode (L2), were identified. While *Ascarophis valentina* and *Capillaria* sp. were adults, *Contracaecum* sp. was third stage larvae (L3) and *H. aduncum* and *H. fabri* were at their third and fourth larval stage. Adult *Acanthocephaloides irregularis* specimens were isolated from the intestine at a low infection rate.

Discussion

The present study is the first report on the

Table 1. Infection site (microhabitat), prevalence (%), mean intensity (MI \pm SE:) and mean abundance (MA \pm SE) values of parasite species found in red mullet, *Mullus barbatus ponticus*

	Microhabitat	Pr (%)	MI \pm SE	MA \pm SE	Min-Max
<i>Loma</i> sp.	rectum	0.91	96.67 \pm 76.56	0.88 \pm 0.76	7-249
<i>Galactosomum lacteum</i>	gills	0.61	3.00 \pm 0.00	0.018 \pm 0.015	1-5
<i>Proctotrema bacilliovatum</i>	intestine and rectum	70.61	29.09 \pm 7.01	20.54 \pm 5.00	1-1187
<i>Progrillotia dasyatidis</i>	gall bladder, stomach	38.79	3.30 \pm 0.49	1.28 \pm 0.21	1-46
<i>Nybelinia</i> sp.	gall bladder, stomach	1.21	1.00 \pm 0.00	0.006 \pm 0.004	1-1
<i>Scolex pleuronectis</i>	intestine	2.12	1.57 \pm 0.30	0.033 \pm 0.014	1-3
<i>Hysterothylacium</i> spp.	intestine, body cavity, gall bladder, stomach	56.36	4.56 \pm 0.45	2.57 \pm 0.28	1-46
<i>Ascarophis valentina</i>	esophagus, stomach	3.94	3.31 \pm 1.21	0.13 \pm 0.06	1-16
<i>Capillaria</i> sp.	stomach	39.39	3.39 \pm 1.21	1.34 \pm 0.15	1-26
<i>Contracaecum</i> sp.	intestine, body cavity, gall bladder, stomach	40.0	4.77 \pm 0.51	1.91 \pm 0.24	1-45
unidentified larval nematode (L2)	Pyloric ceace	43.33	10.78 \pm 1.97	4.67 \pm 0.90	1-170
<i>Acanthocephaloides irregularis</i>	intestine	1.21	2.75 \pm 1.75	0.033 \pm 0.025	1-8
OVERALL		97.88	34.12 \pm 5.23	33.40 \pm 5.62	1-1187

Pr.: prevalence, SE: standard error

parasite fauna of the red mullet captured from the Black Sea coast of Turkey and provides some new data about its parasite fauna and their intensities in relation with season and some host factors.

So far, members of the genera *Aponurus*, *Holorchis*, *Derogenes*, *Lecithaster*, *Lasiotocus*, *Proctotrema* and *Anisocladium* have been reported in *Mullus barbatus* in the Mediterranean Sea (Bray, 1985; Bartoli & Bray, 1996; Bartoli, Gibson & Bray, 2005; Carreras-Aubets *et al.*, 2011; Carreras-Aubets *et al.*, 2012; Debenedetti *et al.*, 2013; Radujkovic & Sundic, 2014). Except *Proctotrema*, the absence of these digenean parasites in this study could be resulted from the lack of the final and intermediate hosts in our sampling area. *Proctotrema bacilliovatum* which was previously recorded in *Mullus surmuletus* in the Sea of Marmara and the Aegean Sea, had not been previously recorded in *M. barbatus ponticus* from the Black Sea coasts of Turkey. The mean intensity of *P. bacilliovatum* was higher in summer than in winter in the present study. Water temperature is one of the most important factors determining release of digenean cercariae from the first host snail. Moreover, transmission to the fish host is highly temperature-dependent. It's could be due to effect of seasonal temperature. To date, *Galactosomum lacteum* metacercaria has been reported in marine teleosts from the Black Sea and the Mediterranean Sea (Lushchina, 1985; Culurgioni, D'Amico Figus, 2007). It is noticeable that this digenean species identified here is reported for the first time in this fish species from Turkey.

Progrillotia dasyatidis has been recorded in gobiid fish in the Sea of Marmara (Oğuz & Bray, 2008), and in various teleosts including *M. barbatus* in the Black Sea coast of Turkey (Tepe, Oğuz & Heckmann, 2014). Marine cestodes such as *Progrillotia dasyatidis* use planktonic copepods as first and, sometimes, second intermediate hosts. Zander, Strohbach and Groenewold (1993) reported that, in benthic fish, larval cestodes are presumably acquired during the period of the preadult stages, as a result of the ingestion of planktonic crustaceans. The results of this study show that infection with *P. dasyatidis* increases with fish length, so that parasites

are still being acquired after the fish have apparently ceased to feed on copepods.

Nematoda was the most represented taxa with two adult (*Ascarophis valentina*, *Capillaria* sp.) and four larval (*Hysterothylacium fabri*, *H. aduncum*, *Contracaecum* sp. and unidentified larval nematode) parasites in *M. barbatus ponticus* in the present investigation. Among nematodes, *H. fabri* and *H. aduncum* showed consistently high prevalence as they were found commonly in *Mullus surmuletus* and *M. barbatus* in the Mediterranean Sea (Radujkovic & Raibaut, 1989; Le Pommelet, Bartoli & Silan, 1997; Carreras-Aubets *et al.*, 2012). Moreover, several fish species have also been reported to act as intermediate and/or paratenic host for *H. fabri* and *H. aduncum* in the Mediterranean and Black Sea (Carreras-Aubets *et al.*, 2012; Tepe & Oğuz, 2013). In the present study, seasonal changes in the prevalence of *Hysterothylacium* spp., *Capillaria* sp. and *Contracaecum* sp. and an unidentified larval nematode (L2) were determined. Statistical analysis has revealed the presence of significant changes among seasons for these parasite species (Table 3). Generally the highest values of the prevalence were recorded in spring and summer. Similarly, Gordon (1977) observed a decline in prevalence during cold months and the highest prevalence in July/August in whiting. Moreover, Ismen and Bingel (1999) found that the prevalence and intensity of *H. aduncum* were significantly higher in the warm season (July/August) than in the colder period (January/February). There is good agreement between the results obtained in the present study and those found by Gordon (1977) and Ismen and Bingel (1999).

Capillariid nematodes use usually oligochaetes and rarely fish as intermediate hosts. Especially, oligochaetes have been reported by Moravec, Prokopic and Shlikas (1987) as act as transport hosts and is important for the transmission of capillariids. *Mullus barbatus* is a benthic predator and mainly feeds on small crustaceans and polychaetes (Chérif *et al.*, 2011). It has been reported that the composition of the ingested prey varied with the size of predator fish and cephalopods occurred exclusively in the diet of

Table 2. Seasonal infection prevalence (%) and mean intensity (MI ± SE) values of parasites found in *M. barbatus ponticus* (n: number of examined fish, SE: standard error, Means followed by the same superscript letter are not significantly different (P < 0.05))

Parasites species	Prevalence (%)				Mean Intensity ± SE			
	Spring (n: 78)	Summer (n: 82)	Autumn (n: 80)	Winter (n: 90)	Spring (Lsp: 14.2 cm)	Summer (Lsu: 13.5 cm)	Autumn (Lau: 13.9 cm)	Winter (Lwi: 14.5 cm)
<i>Loma</i> sp.	1.28	0.0	0.0	2.22	7.00 ± 0.00	0.0	0.0	141.50 ± 107.50
<i>P. bacilliovatum</i>	51.28	79.27	62.50	86.67	11.90 ± 2.73 ^a	77.37 ± 23.98 ^b	10.60 ± 3.23 ^a	9.51 ± 1.37 ^a
<i>G. lacteum</i>	0.0	0.0	2.53	0.0	0.0	0.0	.00 ± 2.00	0.0
<i>P. dasyatidis</i>	37.18	36.59	28.75	51.11	4.03 ± 1.55	3.67 ± 1.11	3.48 ± 1.07	2.50 ± 0.36
<i>Nybelinia</i> sp.	0.0	4.88	0.0	0.0	0.0	1.00 ± 0.00	0.0	0.0
<i>S. pleuronectis</i>	5.13	2.44	0.0	0.0	1.25 ± 0.25	2.50 ± 0.50	0.0	0.0
<i>Hysterothylacium</i> spp.	53.85	41.46	70.00	60.00	2.74 ± 0.35 ^a	3.79 ± 0.85 ^a	7.68 ± 1.25 ^b	3.22 ± 0.41 ^a
<i>A. valentina</i>	1.28	6.09	3.75	4.44	1.00 ± 0.00	4.60 ± 2.87	3.67 ± 2.19	2.00 ± 1.00
<i>Capillaria</i> sp.	46.15	58.54	37.50	17.78	2.50 ± 0.33 ^{bc}	4.90 ± 0.71 ^{bc}	2.87 ± 0.43 ^{bc}	1.88 ± 0.30 ^a
<i>Contraceacum</i> sp.	66.67	28.05	3.75	53.33	3.17 ± 0.55 ^a	6.61 ± 2.01 ^{bc}	2.56 ± 0.97 ^a	6.02 ± 0.74 ^{bc}
larval nematod (L2)	33.33	53.66	52.50	34.44	4.58 ± 2.24	7.93 ± 1.83	14.67 ± 4.95	14.74 ± 5.12
<i>A. irregularis</i>	0.0	4.88	0.0	0.0	0.0	2.75 ± 1.75	0.0	0.0
TOTAL	96.15	97.56	67.50	100	14.60 ± 2.32^a	75.54 ± 19.57^b	19.30 ± 3.09^{bc}	23.32 ± 3.99^c

Lsp, Lsu; Lau, Lwi: seasonally, the average length of the fish

Table 3. Infection prevalence (%) and mean intensity (MI±SE) levels of parasites founded in *M. barbatus ponticus* according to fish lenght classes

Parasites species	Prevalence (%)			Mean Intensity±SE		
	< 13 cm (n: 60)	13-15 cm (n: 193)	> 15 cm (n: 77)	< 13 cm (n: 60)	13-15 cm (n: 193)	> 15 cm (n: 77)
<i>Loma</i> sp.	-	0.52	2.60	-	34.0 ± 0.00	128.0 ± 121.0
<i>P. bacilliovatum</i>	60	72.54	74.02	41.39 ± 20.52	31.97 ± 10.29	14.23 ± 3.61
<i>G. lacteum</i>	-	1.04	-	-	3.00 ± 2.00	-
<i>P. dasyatidis</i>	48.33	25.23	40.26	2.93 ± 0.82	3.25 ± 0.55	3.74 ± 1.45
<i>Nybelinia</i> sp.	-	-	5.19	-	-	1.00 ± 0.00
<i>S. pleuronectis</i>	3.33	1.04	3.89	1.00 ± 0.00	1.50 ± 0.50	1.33 ± 0.33
<i>Hysterothylacium</i> spp.	48.33	58.55	57.14	3.86 ± 0.72	4.49 ± 0.58	5.20 ± 1.12
<i>A. valentina</i>	3.33	3.63	5.19	1.00 ± 0.00	1.86 ± 0.55	7.00 ± 3.34
<i>Capillaria</i> sp.	51.67	38.86	31.17	3.13 ± 0.65	3.16 ± 0.31	4.46 ± 1.12
<i>Contraceacum</i> sp.	36.67	36.27	51.95	4.18 ± 1.05	5.06 ± 0.77	4.58 ± 0.82
unidentified larval nematode (L2)	48.33	46.11	32.41	8.38 ± 2.97	12.84 ± 2.92	6.20 ± 2.50
<i>A. irregularis</i>	3.33	1.04	-	1.00 ± 0.00	4.50 ± 3.50	-
TOTAL	98.33	97.41	98.07	36.05 ± 12.74	37.25 ± 7.81	24.87 ± 4.88

Table 4. Infection prevalence (%) and mean intensity (MI ± SE) values of parasites found in female and males of *M. barbatus ponticus*

Parasites species	Prevalence (%)		Mean Intensity ± SE	
	Female (n:266)	Male (n:64)	Female (Lf: 14.27)	Male (Lm: 13.11)
<i>Loma</i> sp.	0.75	1.56	128.0 ± 121.0	34.0 ± 0.00
<i>P. bacilliovatum</i>	72.18	64.06	30.11 ± 7.62	24.29 ± 17.85
<i>G. lacteum</i>	0.38	1.56	1.00 ± 0.00	5.00 ± 0.00
<i>P. dasyatidis</i>	40.6	31.25	3.57 ± 1.26	1.80 ± 0.35
<i>Nybelinia</i> sp.	1.13	1.56	1.00 ± 0.00	1.00 ± 0.00
<i>S. pleuronectis</i>	1.88	3.12	1.80 ± 0.74	1.00 ± 0.00
<i>H. aduncum</i> <i>H. fabri</i>	56.77	54.67	4.58 ± 0.53	4.46 ± 0.76
<i>A. valentina</i>	3.38	3.12	3.56 ± 1.62	2.75 ± 1.75
<i>Capillaria</i> sp.	40.6	34.38	3.59 ± 0.37	2.41 ± 0.34
<i>Contraceacum</i> sp.	42.11	31.25	4.71 ± 0.54	5.05 ± 1.42
unidentified larval nematode (L2)	42.11	48.44	7.34 ± 1.43	23.19 ± 7.13
<i>A. irregularis</i>	1.13	1.56	3.33 ± 2.33	1.00 ± 0.00
TOTAL	98.1	96.88	34.12 ± 5.78	34.11 ± 12.33

Lf: the average length of the female fish, Lm: the average length of the male fish

larger specimens than smaller ones, and diets varied seasonally; decapods were commonly consumed in summer, whereas amphipods were commonly consumed during winter and spring (Labropoulou, Machias, Tsimenides & Eleftheriou, 1997). The prevalence value of *Capillaria* sp. was higher in summer than that of in winter (Table 2). These values were also negatively correlated with increasing fish size (Table 3). Moreover, larger fish (Lwi: 14.5 cm) caught during winter had lower parasite intensity than smaller fish (Lsu: 13.5) caught during summer. Seasonal average length of fish in summer has higher and this clearly explained its lower presence in other seasons. Thus our results indicated that the prevalence of infection had a relation with host size.

Ascarophis valentina was first identified in *Mullus surmuletus* from the Mediterranean Sea (Ferrer *et al.*, 2005). So far, some species of *Ascarophis* genus have been described in mullid species; *Ascarophis mullusi* in *M. surmuletus* from the Aegean Sea and in *M. barbatus* from the Adriatic Sea. Indeed, *A. valentina* has not been recorded outside the Mediterranean Sea (Le Pommelet *et al.*, 1997; Ferrer *et al.*, 2005; Klimpel, Kleinertz & Palm, 2008) and this study is the first report of *A. valentina* in *M. barbatus ponticus* from the Black Sea.

In conclusion, the metazoan parasite fauna of *M. barbatus ponticus* in Turkish Black Sea coasts firstly reported with this study. We report *Proctotrema bacilliovatum* and *Ascarophis valentina* to be new records for Turkish parasite fauna. *Ascarophis valentina* is also a new record in Black Sea. Thus, we can say that the geographical distribution of this parasite has extended.

References

- Abo-Esa, J.F.K. (2007). Helminth parasites in Barbonny *Mullus barbatus* fish with reference to public health hazards. *Egyptian Journal of Aquatic Biology and Fisheries*, 11(3), 127-137.
- Amin, O.M., Oğuz, M.C., Heckmann, R.A., Tepe, Y., & Kvach Y. (2011). Acanthocephaloides irregularis n. Sp. (Acanthocephala: Arhythmacanthidae) from marine fishes off the Ukrainian Black Sea coast. *Systematic Parasitology*, 80(2), 125-135. <http://dx.doi.org/10.1007/s11230-011-9312-0>
- Aydın, M., & Karadurmuş, U. (2013). An investigation on age, growth and biological characteristics of red mullet (*Mullus barbatus ponticus*, Essipov, 1927) in the Eastern Black Sea. *Iranian Journal of Fisheries Sciences*, 12(2), 277-288.
- Bartoli, P., & Bray, R.A. (1996). Description of three species of *Holorchis* Stossich, 1901 (Digenea: Lepocreadiidae) from marine fishes off Corsica. *Systematic Parasitology*, 35(2), 133-143. <http://dx.doi.org/10.1007/BF00009822>
- Bartoli, P., Gibson, D.I., & Bray R.A. (2005). Digenean species diversity in teleost fish from a nature reserve off Corsica, France (Western Mediterranean), and a comparison with other Mediterranean regions. *Journal of Natural History*, 39(1), 47-70. <http://dx.doi.org/10.1080/00222930310001613557>
- Beveridge, I., Neifar, L., & Euzet, L. (2004). Review of the genus *Progrillotia* Dollfus, 1946 (Cestoda: Trypanorhyncha), with a redescription of *Progrillotia pastinacae* Dollfus, 1946 and description of *Progrillotia dasyatidis* sp. n. *Folia Parasitologica*, 51 (1), 33-44. <http://dx.doi.org/10.14411/fp.2004.005>
- Bush, A.O., Lafferty, K.D., Lotz, J.M., & Shostak, A.W. (1997). Parasitology meets ecology on its own terms: Margolis *et al.* revisited. *Journal of Parasitology*, 83, 575-583.
- Bray, R.A. (1985). Some helminth parasites of marine fishes of South Africa: Families Gorgoderidae, Zoogonidae, Cephaloporidae, Acanthocolpidae and Lepocreadiidae (Digenea). *Journal of Natural History*, 19(2), 377-405. <http://dx.doi.org/10.1080/00222938500770271>
- Bray, R.A., Gibson, D.I., & Jones, A. (2008). Keys to trematoda (Volume 3). CAB International, Wallingford, 848 pp.
- Carreras-Aubets, M., Ropullés-Albelda, A., Kostadinova, A., & Carrassón, M. (2011). A new cryptic species of

- Aponurus* Looss, 1907 (Digenea: Lecithasteridae) from Mediterranean goatfish (Teleostei: Mullidae). *Systematic Parasitology*, 79(2), 145–159. <http://dx.doi.org/10.1007/s11230-011-9297-8>
- Carreras-Aubets, M., Montero, F.E., Kostadinova, A., & Carrassón, M. (2012). Parasite communities in the red mullet, *Mullus barbatus* L., respond to small-scale variation in the levels of polychlorinated biphenyls in the Western. *Mediterranean Marine Pollution Bulletin*, 64(9), 1853–1860. <http://dx.doi.org/10.1016/j.marpolbul.2012.06.008>
- Chérif, M., Amor, M.M.B., Selmi, S., Gharbi, H., Missaoui, H., & Capapé, C. (2011). Food and feeding habits of the red mullet, *Mullus barbatus* (Actinopterygii: Perciformes: Mullidae), off the northern Tunisian coast (Central Mediterranean). *Acta Ichthyologica et Piscatoria*, 41(2), 109–116. <http://dx.doi.org/10.3750/AIP2011.41.2.06>
- Culurgioni, J., D'Amico, V., & Figus, V. (2007). Metacercariae of *Galactosomum lacteum* (Jagerskiöld, 1896) Looss, 1899 (Heterophyidae) from marine teleosts in the Gulf of Cagliari (southern Sardinia, Italy). *Journal of Helminthology*, 81(4), 409–413. <http://dx.doi.org/10.1017/S0022149X07862912>
- Debenedetti, A.L., Madrid, E., & Fuentes, M.V. (2013). Study of helminth parasites in the red mullet, *Mullus barbatus*, from the Mediterranean Sea and acquired in greater València, Spain. *Revista Ibero-Latinoamericana de Parasitología*, 72(2), 118–123.
- Essafi, K., Raibaut, A., & Boudaoud-Krissat, K. (1983). *Colobomatus steenstrupi* (Richiardi, 1876) and *Colobomatus mulli* n.sp. (Copepoda: Philichthyidae), parasitic on fish of the genus *Mullus* (Mullidae) in the western Mediterranean. *Systematic Parasitology*, 5(2), 135–142. <http://dx.doi.org/10.1007/BF00049240>
- Ferrer, E., Aznar, F.J., Balbuena, J.A., Kostadinova, A., Raga, J.A., & Moravec, F. (2005). A new cystidicolid nematode from *Mullus surmuletus* (Perciformes: Mullidae) from the western mediterranean. *Journal of Parasitology*, 91(2), 335–344. <http://dx.doi.org/10.1645/GE-366R>
- Gaevskaya, A.V. (2012). Parasites and diseases of the fishes of the Black and Azov Seas. Sevastopol: ECOSI-Gidrofizika. 380 pp. (in Russian)
- Gordon, J.D.M. (1977). The fish populations in inshore waters of the west coast of Scotland. The food and feeding of the whiting, *Merlangius merlangus*. *Journal of Fish Biology*, 11(6), 513–529. <http://dx.doi.org/10.1111/j.1095-8649.1977.tb05709.x>
- Hureau, J.C. (1986). Mullidae. In: Whitehead P.J.P., Bauchot M.L., Hureau J.C., Nielsen J. & Tortonese E. (eds) Fishes of the North-Eastern Atlantic and the Mediterranean. Paris, UNESCO, pp. 877–882.
- Ismen, A., & Bingel, F. (1999). Nematode infection in the whiting *Merlangius merlangus euxinus* off Turkish Coast of the Black Sea. *Fisheries Research*. 42(1-2). 183–189. [http://dx.doi.org/10.1016/S0165-7836\(99\)00022-3](http://dx.doi.org/10.1016/S0165-7836(99)00022-3)
- Klimpel, S., Kleinertz, S., & Palm, H.W. (2008). Distribution of parasites from red mullets (*Mullus surmuletus* L., Mullidae) in the North Sea and Mediterranean Sea. *Bulletin of Fish Biology*, 10(1-2), 1–14.
- Labropoulou, M., Machias, A., Tsimenides, N., & Eleftheriou, A. (1997). Feeding habits and ontogenetic diet shift of the striped red mullet, *Mullus surmuletus* Linnaeus, 1758. *Fisheries Research*, 31(3), 257–267. [https://doi.org/10.1016/S0165-7836\(97\)00017-9](https://doi.org/10.1016/S0165-7836(97)00017-9)
- Le Pommelet, E., Bartoli, P., & Silan, P. (1997). Biodiversité des digènes et autres helminthes intestinaux des Rougets: synthèse pour *Mullus surmuletus* (Linné 1758) et *M. barbatus* (L. 1758) dans le bassin méditerranéen. *Annales des Sciences Naturelles-Zoologie et Biologia Animale*, 18(1), 117–133.
- Lushchina, V.G. (1985) The helminth fauna of blennioid fish in the Black Sea. *Ekologiya Morya*, 20, 43–48.
- Moravec, F., Prokopic, J., & Shlikas, A.V. (1987). The biology of nematodes of the family Capillariidae Neveu-Lemaire, 1936. *Folia Parasitologica*, 34(1), 39–56.
- Oğuz, M.C., & Bray, R.A. (2008). Cestoda and Monogenea of some teleost fishes off the Mudanya Coast (Sea of Marmara, Turkey). *Helminthologia*, 45(4), 192–195. <https://doi.org/10.2478/s11687-008-0038-8>
- Ovcharenko, M.O., & Yurahno, V.M. (2006). First Registration of *Loma* (Microsporidia, Glugeidae) Belonging Microsporidia in Black Sea Fish. *Vestnik zoologii*, 40 (4), 291–296.
- Paradiznik, V., & Radujkovic, B. (2007). Digenea trematodes in fish of the North Adriatic Sea. *Acta Adriatica*, 48(2), 115–129.
- Petter, A.J., Lebre, C., & Radujkovic, B.M. (1984). Nematode parasites of osteichthyen fish of the southern Adriatic. *Acta Adriatica*, 25, 205–221.
- Radujkovic, B.M., & Raibaut, A. (1989). Parasites des poissons marins du Montenegro: liste des espèces de poissons avec leurs parasites. *Acta Adriatica*, 30(1/2), 307–320.
- Radujkovic, B.M., & Sundic, D. (2014). Parasitic flatworm (Platyhelminthes: Monogenea, Digenea, Cestoda) of fishes from the Adriatic Sea. *Nature Montenegrina, Podrogonica*, 13(1), 7–280.
- Ramdane, Z., & Trilles, J. (2010). New Algerian parasitic copepods. *Bulletin European Association of Fish Pathologists*, 30: 41–47.
- Sahin, T., & Akbulut, B. (1997). Some biological characteristics of *Mullus barbatus ponticus* Essipov, 1927 in the Eastern Black Sea coast of Turkey. *Turkish Journal of Zoology*, 21(2), 179–185.
- Tepe, Y., & Oğuz, M.C. (2013). Nematode and Acanthocephalan parasites of marine fish of eastern Black Sea coast of Turkey. *Turkish Journal of Zoology*, 37(6), 753–760. <http://dx.doi.org/10.3906/zoo-1206-18>
- Tepe, Y., Oğuz, M.C., & Heckmann, R.A. (2014). Digenean and cestode parasites of teleost fish from the Eastern Black Sea Region. *Turkish Journal of Zoology*, 38(2), 209–215. <http://dx.doi.org/10.3906/zoo-1205-5>
- Turan, C. (2006). Phylogenetic relationships of Mediterranean Mullidae species (Perciformes) inferred from genetic and morphologic data. *Scientia Marina*, 70(2), 311–318.
- Zander, C.D., Strohbach, U., & Groenewold, S. 1993. The importance of gobies (Gobiidae, Teleostei) as hosts and transmitters of parasites in the SW Baltic. *Helgol Wiss Meeresuntersuch*, 47(1), 81–111. <http://dx.doi.org/10.1007/BF02366186>