



An Evaluation of Optimal Dietary Protein Requirements of All Parts Chicken Meal for Indian Major Carps (*Catla catla*, *Labeo rohita* and *Cirrhinus mrigala*)

Aasia Karim^{1*} and Mohammad Shoaib²

¹ Federal Urdu University of Arts, Science and Technology, Department of Zoology, Karachi, Pakistan.

² University of Karachi, Department of Zoology, Karachi 75270, Pakistan.

Phone: # 0322-2139244

E-mail: aasiakarim@gmail.com

Abstract

A feeding experiment was conducted for one year with six formulated diets containing all parts chicken meal in three different inclusion levels (25%, 35% and 45%) as APCM I, APCM II, APCM III and Fish meal as Control I, Control II and Control III, to examine the potential of all parts chicken meal as a substitute of fish meal in the diet of Indian major carps in intensive polyculture system. Fingerlings of *Catla catla*, *Labeo rohita* and *Cirrhinus mrigala* were fed on experimental diets. It is evident by the results that growth is highly affected by dietary ingredients and its level of inclusion in diet. With increasing level of fishmeal, a noticeable decrease in growth was observed, as Control I, II and III produced 67.3 g, 50.5 g and 39.6 g mean weight gain respectively. While opposite trend of growth was observed with APCM based diet i.e., APCM III (45% inclusion of APCM) produced significantly higher growth with a decreased FCR. The two-way analysis of variance for weight, DFA, SGR and FCR against months showed highly significant relationship in all diets. Except moisture and ash, fish carcass composition was significantly affected by protein source and its inclusion level.

Keywords: Chicken meal, Carps, Polyculture

Introduction

Fish meal is used as a main dietary protein source in aquaculture industry in order to increase the energy content of feed by the expansion of intensive aquaculture (Tacon & Jackson, 1985; Kaushik, Medale, Fauconneau, & Blanc, 1989). However, about 35% of the total global fish catch is being used in the production of fish meal (Tacon & Dominy, 1999), as approximately 1 Kg dry fish meal is obtained from 4 kg of wet fish (Allan et al., 2000), but its cost is going to be continuously rising due to its high demand and therefore directly affecting total expenditure of aquaculture industry. The exploration for alternative to replace fish meal is global research precedence nowadays (Manzi, 1989; Hardy & Kissil, 1997). During recent years substantial research efforts have been made to replace fish meal from aquaculture feed with some other alternative ingredients as potential substitutes of fish meal (Tacon & Jackson, 1985). Aquaculture of carp's species has been expanded to a great extent during last ten years in Pakistan. The combination of all parts chicken meal with rice polish and corn gluten meal has been employed for the evaluation of successful



carps farming. This research trial has been made to complementing the efficacy of all parts chicken meal diets with fish meal based diets in an intensive polyculture of major carps i.e. *Catla catla*, *Labeo rohita* and *Cirrhinus mrigala*.

Materials and Methods

Experimental Design, Diets Formulation and Feeding Regime

Six experimental diets were formulated by using fishmeal and all parts chicken meal in three different inclusions i.e. 25%, 35% and 45%. Fishmeal based diets were named as control I, II, III for 25%, 35% and 45% inclusion level respectively, while diets with all parts chicken meal were labeled as APCM I, II and III for three inclusion levels in that order. All experimental diets were formulated by mixing of fish meal (only in controls) or all parts chicken meal with corn gluten meal, rice polish, starch and canola oil (Table 1). Ingredients were pulverized, mixed and then pelletized with the help of a pellet maker. Pellets were then dehydrated at room temperature for 24 hours and followed by storing in freezer.

Each experiment was performed in triplicates for the accuracy of data and thus eighteen re-circulatory concrete raceways of the dimension 22'×50' (W×L) were designed. The juveniles of *Catla catla*, *Labeo rohita* and *Cirrhinus mrigala* were reared in each raceway at a ratio of 33:33:34 respectively by following Wahab, Rahman and Milstein (2002) for a period of one year.

Water quality variables such as temperature of water, dissolved oxygen, pH and ammonia were recorded weekly. The water temperature ranged 10.1°C to 30.5°C, pH ranged 7.8 to 8.4 and dissolved oxygen was within the range of 5.1 to 8.4 mg/l during the whole experimental period.

The Daily feed allowance (DFA) was calculated at 3% body weight and feed was supplied manually twice daily.

Estimation of Feed Response

At the end of each month, a few fishes were randomly opted from each raceway, to measure monthly increase in weight and length and then released back into their respective raceway. The following growth parameters were investigated during whole research trial; Monthly weight gain, survival rate, feed conversion ratio (FCR), specific growth rate (SGR) and Protein efficiency ratio (PER) (Khan, Ahmed & Abidi, 2004).

Proximate Composition of Fish Meat

The proximate composition of under considered carps was evaluated in terms of crude protein, carbohydrate, fat, ash and moisture in accordance with Association of Official Analytical Chemist (A.O.A.C, 2005) to check the effectiveness of formulated diet on nutritional quality of fish after consuming the experimental diets.

Statistical Analysis of Experimental Data

After compilation, data was subjected to statistical analysis from Minitab release 16.1. Fisher's least-significant-difference (LSD) test and Two-way analyses of variance were executed to discover relationships among growth variables.



Results

The response (means of three replicate) of experimental carps against different inclusion levels i.e. 25%, 35% and 45% of fishmeal (Controls) and all parts chicken meal (APCM) is presented in Table 2. It is evident by the results that growth is highly affected by dietary ingredients and its level of inclusion in diet. In the present feeding trial, with increasing level of fishmeal, a noticeable decrease in growth was observed, as Control I, II and III produced 67.3 g, 50.5 g and 39.6 g mean weight gain respectively for all three fish species. While opposite trend of growth was observed with APCM based diet i.e., APCM III (45% inclusion of APCM) produced significantly higher growth with a decreased FCR. However, a considerably higher growth was obtained by APCM II and APCM III in comparison with Control II and Control III, but APCM I couldn't proved itself better than Control I (25% inclusion level of fishmeal) with respect to weight gain although no differences were found in the values of FCR (3.5). 45% fish meal based diet showed minimal weight gain (39.6 g) among all levels of Control and APCM.

However no significant differences were found in the means of DFA and SGR among all treatments by LSD Test, but the values of FCR were differed significantly in APCM II and APCM III (3.33 and 2.42 respectively). FCR gradually decreased as inclusion of APCM was increased while in case of fish meal (Control) the result was vice versa. Highest production was obtained by APCM III as 75.08 Kg/treatment followed by Control I which yielded 66.56 Kg/treatment (Table 3). Percent individual contribution indicated a remarkably higher contribution of *Cirrhinus mrigala* in all three levels of All parts chicken meal based diets and Controls than other two species i.e. *Labeo rohita* and *Catla catla*. Survival rate was 80% after one year trial in all treatments. Water quality was not the actual reason of mortalities. Except moisture and ash, fish carcass and fillet nutrient composition was significantly affected by protein source and inclusion level (Table 5).

The two-way analysis of variance for weight, DFA, SGR and FCR against months showed highly significant relationship in all diets (Table 4).

Discussion

A number of studies have been carried out to evaluate the potential of chicken meal as fish meal replacers without any significant reduction in growth of the fish (Fowler, 1991; El Sayed, 1998). In the present feed trial 100% substitution of fishmeal was made by poultry by product meal. It was evident by comparison of FM (Control) with APCM that all parts chicken meal was not so much effective at lower concentrations (25%) as compare to control I, but a noticeable increase in growth was evident in comparison with control II and III. Due to the higher digestibility rate (Sugiura, Dong, Rathbone & Hardy, 1998) APCM produced higher final body weight in APCM II and APCM III than that of the Controls, thereby boosting growth. The acceptance could be basically attributed to improved manufacturing practices, enhancing the quality of the feed (Bureau, Harris & Cho, 1999 & Bureau et al., 2000). In addition, chicken meals are locally available and economically more viable than fish meal even at total replacement levels (Rodriguez-Serna, Olvera-Novoa & Carmona-Osalde, 1996).

In most of the treatments *C. mrigala* showed an exceptional growth among all three species, as it attained maximum feed being a bottom feeder. Apparently, replacement of fish meal with APCM did not influence growth, SGR, or FCR



of major carps (*Catla catla*, *Labeo rohita* and *Cirrhinus mrigala*) and thus confirmed the capability of major carps to utilize poultry by-product meal as a better protein than fish meal.

Except moisture and ash, all body compositions were significantly affected ($P \leq 0.05$) by protein source and its inclusion level. Maximum carcass protein was observed by APCM I and control II (FM 35%). In the present feed trial, fish growth was not adversely affected by increased fat content of APCM. This result is similar to the findings of El-Sayed (1998).

APCM showed its potential and could proficiently surrogate up to 100% fish meal protein in diet of major carps. Many researchers have been applied this for many species (Fowler, 1991; Sugiura et al., 1998; Steffens, 1994; Alexis, 1997). Several studies have been signified chicken meal for its potential to replace 1000 g per Kg of fish meal for many species (Steffens, 1994; Alexis, 1997; Nengas, Alexis & Davies, 1999; Kureshy, Davis & Arnold, 2000; Takagi, Hosokawa, Shimeno & Ukawa, 2000). A little attention has been paid for the evaluation of APCM on cyprinidae by researchers (Welcomme, 1988).

The results of this finding coincide with those of Alexis (1997), who substituted 100% fish meal by poultry by-product meal in practical diets for sea bass and sea bream devoid of any adverse effect on growth performance. Similar results were also observed in chinook salmon *Oncorhynchus tshawytscha* W. (Fowler, 1991) and gilthead sea bream (Nengas et al., 1999). Based on our results, major carps are capable to accept 100% replacement of fish meal with APCM in a higher proportion as compare to fish meal without decrease in growth performance.

References

- A.O.A.C. (2005). Official Methods of Analysis. (16th Ed.). Association of Official Analytical Chemists. Arlington, VA, USA. 1193 pp.
- Alexis, M. N. (1997). Fish meal and fish oil replacers in Mediterranean marine fish diets. In: Feeding Tomorrow's Fish. *Proceedings of the workshop of the CIHEAM network on technology of aquaculture in the Mediterranean* (Tacon, A. & Barsureo, B. eds), CIHEAM, Zaragoza, Spain, 1989, 183–204 pp.
- Allan, G. L., Parkinson, S., Booth, M. A., Stone, D. A., Rowland, S. J., Frances, J., & Warner-Smith, R. (2000). Replacement of fish meal in diets for Australian silver perch, *Bidyanus bidyanus*: I. Digestibility of alternative ingredients. *Aquaculture*, 186(3), 293-310.
- Bureau, D. P., Harris, A. M., & Cho, C. Y. (1999). Apparent digestibility of rendered animal protein ingredients for rainbow trout. *Aquaculture*, 180, 345–358.
- Bureau, D. P., Harris, A. M., Beven, D. J., Simmons, L. A., Azevedo, P. A., & Cho, C. Y. (2000). Feather meals and meat and bone meals from different origins as protein sources in rainbow trout (*Oncorhynchus mykiss*) diets. *Aquaculture*, 181, 281–291.
- El-Sayed, A. F. (1998). Total replacement of fish meal with animal protein sources in Nile tilapia, *Oreochromis niloticus* (L.), feeds. *Aquaculture Research*, 29(4), 275-280.
- Fowler, L. G. (1991). Poultry by-product meal as a dietary protein source in fall chinook salmon diets. *Aquaculture*, 99(3-4), 309-321.
- Hardy, R. W., & Kissil, G. W. M. (1997). Trends in aquaculture feeding. *Feed Mix* 5, 31–34.
- Kaushik, S. J., Medale, F., Fauconneau, B., & Blanc, D. (1989). Effect of digestible carbohydrates on protein/energy utilization and on glucose metabolism in rainbow trout (*Salmo gairdneri* R.). *Aquaculture*, 79(1-4), 63-74.
- Khan, M. A., Ahmed, I., and Abidi, S. F. (2004). Effect of ration size on growth, conversion efficiency and body composition of fingerling mrigala, *Cirrhinus mrigala* (Hamilton). *Aquaculture Nutrition*, 10, 47–53.
- Kureshy, N., Davis, D. A., & Arnold, C. R. (2000). Partial replacement of fish meal with meat and bone meal, flash-dried poultry by-product meal, and enzyme-digested poultry by-product meal in practical diets for juvenile red drum. *North American Journal of Aquaculture*, 62, 266–272.



- Manzi, J. J. (1989). Aquaculture research priorities for the 1990s. *World Aquaculture*, 20, 29–32.
- Nengas, I., Alexis, M.N. & Davies, S.J. (1999). High inclusion levels of poultry meals and related by products in diets for gilthead seabream *Sparus aurata* L. *Aquaculture*, 125, 119–129.
- Rodriguez-Serna M., Olvera-Novoa M. A., & Carmona-Osalde, C. (1996). Nutritional value of animal by-product meal in practical diets for Nile tilapia *Oreochromis niloticus* (L.) fry. *Aquaculture Research*, 27, 67–73.
- Steffens, W. (1994). Replacing fish meal with poultry by-product meal in diets for rainbow trout *Oncorhynchus mykiss*. *Aquaculture*, 124, 27–34.
- Sugiura, S. H., Dong, F. M., Rathbone, C. K., & Hardy, R. W. (1998). Apparent protein digestibility and mineral availabilities in various feed ingredients for salmonid feeds. *Aquaculture*, 159, 177–202.
- Tacon, A. G. J., & Jackson, A. J. (1985). Utilization of conventional and unconventional protein sources in practical fish feeds. In: Nutrition and Feeding of Fish (Cowey, C.B., Mackie, A. M., & Bell, J. G. eds), Academic Press, London, UK, 119–145 pp.
- Tacon, G. J., & Dominy, W. G. (1999). Overview of world aquaculture and aquafeed production. In: Book of Abstracts. World Aquaculture '99, 26 April–2 May 1999, Sydney, Australia. World Aquaculture Society, Baton Rouge, LA, 853 pp.
- Takagi, S. T., Hosokawa, H., Shimeno, S., & Ukawa, M. (2000). Utilization of poultry by-product meal in a diet for red sea bream *Pagrus major*. *Nippon Suisan Gakkaishi*, 66, 428–438.
- Wahab, M. A., Rahman, M. M., & Milstein, A. (2002). The effects of common carp, *Cyprinus carpio* (L.) and *Cirrhinus mrigala* (Hamilton) as bottom feeders in major Indian carp poly cultures. *Aquaculture Research*, 33, 547–556.
- Welcomme, R. L. (1988). International introductions of inland aquatic species. FAO Fisheries and aquaculture Technical Paper. No. 294, 318 pp.

Table 1. Percentages of ingredients, proximate values and energy contents per 100 g of experimental diets.

	Control I	*APCM I	Control II	APCM II	Control III	APCM III
Ingredients (%)						
Fish meal	25	-	35	-	45	-
All Parts Chicken meal		25	-	35	-	45
Corn gluten meal	1.16	1.74	16.69	16.90	30.98	32.06
Rice polish	63.83	63.24	38.30	38.08	14.00	12.93
Starch	5	5	5	5	5	5
Canola oil	4.5	4.5	4.5	4.5	4.5	4.5
Vitamins and mineral mixture	0.5	0.5	0.5	0.5	0.5	0.5
Proximate composition (%)						
Crude protein	24.99	24.98	35.2	34.98	44.98	44.99
Crude fat	15.19	16.85	12.74	15.12	10.42	13.39
Crude fiber	2.56	2.95	2.48	3.02	2.38	3.07
Ash	13.7	12.02	13.42	11.13	13.25	10.24
Nitrogen-free extract	42.99	42.63	35.6	35.17	28.39	27.74
DE (K cal/Kg)	3109.90	3231.26	3186.43	3356.82	3263.44	3482.67
GE (K cal/Kg)	4520.48	4655.95	4554.32	4824.17	4587.67	4832.79

*APCM = All Parts Chicken meal,

Table 2. Mean values of growth parameters of experimental carps fed with different levels of All Parts Chicken meal based diets.



	Weight (g)			Mean WG	ADG (g/day) ²	DFA (g) ³	SGR (%/day) ⁴	FCR ⁵	PER ⁶
	Initial	Final	Monthly WG ¹						
Control I									
<i>C. catla</i>	36.11	823.14	65.6 ±13.3abc		2.18abc		0.37 ±0.09a		8.4±0.20a
<i>L. rohita</i>	18.90	813.85	66.0 ±14.3abc	67.3	2.19abc	679.0±122a	0.34 ±0.10a	3.53±0.41a	8.4±0.20a
<i>C.mrigala</i>	44.25	889.14	70.4 ±15.6abc		2.34abc		0.34 ±0.04a		8.4±0.26a
*APCM I									
<i>C. catla</i>	28.6	615.5	48.90 ±7.66cde		1.63cde		0.37 ±0.04a		6.4±0.20c
<i>L. rohita</i>	37.5	621.2	48.64 ±6.25abcde	51.4	1.62abcde	629.8±117a	0.33±0.04a	3.51±0.31a	6.4±0.23c
<i>C.mrigala</i>	64.1	746.3	56.85 ±6.59a		1.89a		0.29±0.03a		7.2±0.25b
Control II									
<i>C. catla</i>	36.11	623.42	48.94 ±6.4cde		1.63cde		0.45 ±0.10a		4.5±0.20e
<i>L. rohita</i>	18.90	645.14	52.19 ±5.11abcde	50.5	1.73abcde	662.7±104a	0.42 ±0.15a	4.05±0.45a	4.8±0.40e
<i>C.mrigala</i>	44.25	648.71	50.37 ±6.7cde		1.67cde		0.31 ±0.03a		4.5±0.17e
APCM II									
<i>C. catla</i>	27.9	752.3	60.36 ±9.92cde		2.10cde		0.39 ±0.05a		6.0±0.10c
<i>L. rohita</i>	38.6	801.4	63.56 ±9.36abcd	62.4	2.11abcd	724.2±146a	0.36±0.04a	3.33±0.33a	6.0±0.23c
<i>C.mrigala</i>	61.7	822.0	63.3 ±7.76bcde		2.11cde		0.31±0.04a		6.0±0.00c
Control III									
<i>C. catla</i>	27.31	466.43	36.59 ±5.5e		1.21e		0.36 ±0.05a		2.6±0.30h
<i>L. rohita</i>	37.52	490.50	37.68 ±3.9de	39.6	1.25de	532.3±89a	0.32 ±0.07a	3.82±0.33a	2.6±0.17h
<i>C.mrigala</i>	62.51	599.30	44.73 ±4.1cde		1.49cde		0.27 ±0.03a		3.1±0.40g
APCM III									
<i>C. catla</i>	28.5	963.5	77.9 ±15.0bcde		2.59bcde		0.42 ±0.05a		5.5±0.10d
<i>L. rohita</i>	37.9	657.1	51.6 ±7.05abcd	68.3	1.72abcd	604.5±121a	0.34±0.06a	2.42±0.22b	3.7±0.40f
<i>C.mrigala</i>	64.5	972.0	75.6 ±9.86ab		2.52ab		0.32±0.04a		5.5±0.10d

*APCM = All Parts Chicken meal,

Values are means ± SE of three replicates.

Means in a column followed by different letter were significantly different from each other at $P = 0.05$ by the Fisher's least-significant-difference (LSD) test.

¹Monthly weight gain (WG) (g) = Final value of growth variable – Initial value of growth variable



² Average daily gain (ADG) (g/day) = weight gain/number of days

³ Daily Feed Allowance (DFA) (g) = Av body weight X Number of stocks X % Survival X Feeding rate

⁴ Specific growth rate (SGR) (%/day) = Log Fish final weight – Log Fish initial weight / Time X 100

⁵ Feed conversion ratio (FCR) = Weight of food presented/Weight of animal gained

⁶ Protein efficiency ratio (PER) = Weight gain/Crude protein intake

Table 3. Observed and computed fish biomass harvested against different inclusions.

	Control I	*APCM I	Control II	APCM II	Control III	APCM III
Total harvested weight (Kg per treatment)	66.56	52.30	50.49	62.58	41.06	75.08
Total fish production (Kg/hectare/year)	665.68	523.04	504.97	625.82	410.60	750.87
Percent individual contribution at harvest (%)						
<i>C. catla</i>	32.14	30.59	32.09	31.24	29.53	33.36
<i>L. rohita</i>	31.75	30.87	33.21	33.29	31.05	31.68
<i>C.mrigala</i>	36.06	38.52	34.68	35.46	39.40	34.95

*APCM = All Parts Chicken meal,

Table 4. Two-way analysis of variance of different variables against treatments and months.

Variables	Weight (g)			DFA (g)	SGR (%/day)			FCR
	<i>C. catla</i>	<i>L. rohita</i>	<i>C. mrigala</i>		<i>C. catla</i>	<i>L. rohita</i>	<i>C. mrigala</i>	
	Control							
Diet	0.000**	0.002**	0.189*	0.000**	0.782*	0.251*	0.054*	0.348*
Months	0.000**	0.000**	0.000**	0.000**	0.000**	0.000**	0.000**	0.000**
	*APCM							
Diet	0.001**	0.002**	0.000**	0.000**	0.005**	0.458*	0.054*	0.000**
Months	0.000**	0.000**	0.000**	0.000**	0.000**	0.000**	0.000**	0.000**

*APCM = All Parts Chicken meal, **=Significant; * = Non significant

Table 5. Comparison of means of the proximate values under different treatments.



Diet		Moisture %	Crude Protein %	Crude Fat %	Carbohydrate %	Total Ash %
Control I	<i>Catla catla</i>	78.23	16.50	1.94	1.52	1.81
	<i>Labeo rohita</i>	76.76	17.54	2.25	1.60	1.57
	<i>Cirhinus mrigala</i>	78.44	17.33	1.89	0.19	2.14
	Mean	77.81b	17.12ab	2.02a	1.10b	1.84ab
APCM I	<i>Catla catla</i>	79.01	16.61	2.02	0.42	1.93
	<i>Labeo rohita</i>	75.84	17.89	2.10	2.82	1.35
	<i>Cirhinus mrigala</i>	77.19	17.42	2.15	1.23	1.88
	Mean	77.34b	17.30a	2.09a	1.49b	1.72b
Control II	<i>Catla catla</i>	78.60	16.88	2.05	1.33	1.14
	<i>Labeo rohita</i>	76.18	17.82	1.95	2.56	1.42
	<i>Cirhinus mrigala</i>	77.20	17.54	2.01	1.42	1.70
	Mean	77.32b	17.41a	2.00a	1.77b	1.42b
APCM II	<i>Catla catla</i>	80.95	12.24	1.07	4.11	2.05
	<i>Labeo rohita</i>	80.15	13.25	1.25	3.62	1.94
	<i>Cirhinus mrigala</i>	80.04	11.83	1.50	4.59	2.58
	Mean	80.38a	12.44c	1.27b	4.10a	2.19a
Control III	<i>Catla catla</i>	76.52	17.23	2.65	1.84	1.71
	<i>Labeo rohita</i>	79.48	16.31	1.90	0.65	1.65
	<i>Cirhinus mrigala</i>	77.69	16.91	1.60	1.23	2.34
	Mean	77.89b	16.81ab	2.05a	1.24b	1.9ab
APCM III	<i>Catla catla</i>	79.25	15.72	1.70	1.61	1.72
	<i>Labeo rohita</i>	77.09	16.75	1.61	2.49	1.60
	<i>Cirhinus mrigala</i>	78.11	16.20	1.85	1.68	2.13
	Mean	78.15b	16.22b	1.72ab	1.92b	1.81ab
	P Value	0.070	0.000*	0.015*	0.004*	0.174

* Significant.

Means in a column followed by different letter were significantly different from each other at $P = 0.05$ by the Fisher's least-significant-difference (LSD) test.