



## Trophic Ecology of Eight Sympatric Nemipterid Fishes (Nemipteridae) in the Lower Part of the South China Sea

Mithun Paul<sup>1</sup>, Sukree Hajisamae<sup>2,\*</sup>, Siriporn Pradit<sup>1</sup>, Permsak Perngmark<sup>3</sup>, Rashedul Islam<sup>1</sup>

<sup>1</sup> Prince of Songkla University, Marine and Coastal Resources Institute (MACORIN), Hat Yai, Songkhla 90112, Thailand.

<sup>2</sup> Prince of Songkla University, Faculty of Science and Technology, Pattani 94000, Thailand.

<sup>3</sup> Southern Marine Fisheries and Development Center, Songkhla, Songkhla 90000, Thailand.

Phone: 66-73-313928

E-mail: hsukree@hotmail.com

### Abstract

Trophic ecology of eight species of nemipterid fish including *Nemipterus mesoprion*, *N. hexodon*, *N. nemurus*, *N. nematophorus*, *N. tambuloides*, *N. peronii*, *N. furcosus* and *Scolopsis taeniopterus* were studied to investigate their diet composition and trophic relationships between them. Our findings revealed that they were specific predators feeding mainly on shrimp and fish. *N. nemurus* and *N. tambuloides* were the most piscivorous species of them all, and *N. mesoprion* was a specific shrimp predator. In general, there were highly significant differences of stomach fullness (FL) and average number of food item (AF) among species ( $P < 0.01$ ). Size classes significantly affected FL of five species including *N. nematophorus*, *N. mesoprion*, *N. nemurus*, *N. tambuloides* and *S. taeniopterus* ( $P < 0.05$ ) and AF of four species including *N. nemurus*, *S. taeniopterus*, *N. hexodon* and *N. Peronii* ( $P < 0.05$ ). Sex was an influential factor on FL of *N. mesoprion* ( $P < 0.01$ ) and *N. peronii* ( $P < 0.05$ ) and AF of *N. hexodon* ( $P < 0.05$ ) and *N. peronii* ( $P < 0.05$ ). Maturity stages significantly affected FL of *N. mesoprion* ( $P < 0.01$ ) and *S. taeniopterus* ( $P < 0.05$ ), however, they had no impact on AF of any species ( $P > 0.05$ ). The co-existence of these species in the bottom waters of this habitat requires partitioning of available food resources.

**Keywords:** Feeding ecology, thread fin bream, demersal fish, Gulf of Thailand, fish biology

### Introduction

Trophic ecology knowledge is crucial to understand functional role of fish species in a particular ecosystem (Blaber, 1997; Wotton, 1998; Cruz-Escalona, Abitia-Cardenes, Campos-Davila, & Galvan-Magana, 2000; Linke, Platell, & Potter, 2001; Almeida, 2003; Hajisamae, Yeesin, & Ibrahim, 2006; Hajisamae, 2009). It is one of the most appropriate factors that provides help to define the success of a species in the habitat (Almeida, 2003) and understand food web dynamics (Trueman, Johnston, O'Hea, & MacKenzie, 2014; Espinoza, Samantha, Tayler, Aaron, & Ingo, 2015). Community structure of co-existing species can be described by the position of these species along different resource dimensions of space and time (Pianka, 1969; Ross, 1986; Bohorquez-Herrera, Cruz-Escalona, Adams, & Peterson, 2015). Understanding the factors affecting species co-existence is very important in ecological studies (Baeta & Ramon, 2013). In marine community, body sizes of both fish and its prey have been related directly to foraging success (Persson, 1990; Juanes, 1994; Hughes,



1997; Scharf et al., 2000; Karpouzi & Stergiou, 2003). Diets of most fishes change with growth, but the timing of changes varies from species to species and is often associated with changes in lifestyle, habitats (Blaber, 2000) and morphological characteristics (Labropoulou & Papadopoulou-Smith, 1999). The ultimate objective of dietary change is to maximize energy intake, enhance growth rate and minimize the risk of predation in competing for food with bigger predators (Brown, 1985). Many studies reported relationship between prey size and fish morphology or fish behavior including mouth dimension, visual acuity, digestive capacity and swimming performance (Keast & Webb, 1966; Galis, 1990; Kaiser & Hughes, 1993; Juanes, 1994; Juanes & Conover, 1994; Hart, 1997). Intra-specific and inter-specific food partitioning is a strategy for a survival of the species within the ecosystem and has been recognized as an important factor structuring fish community in a particular habitat (Carrete, 2010). Diet composition of a particular species has an important application to the sympatric and co-existing species. Nemipteridae or threadfin breams are important fish resources for both artisanal and commercial fisheries (Russell, 1990). They are extensively found in tropical indo-pacific regions (Russell, 1993), and have become important target species in many countries including India (Joshi, 2005), Japan, Taiwan (Russell, 1993), Indonesia, China (Ping, Rensie, & Jing, 2011) and others (ElHaweet, 2013). In Thailand, various species of threadfin fin breams contribute greatly to the trawl fisheries with the possible consequence of a depletion of resources (Stobutzki et al., 2006). Careful management is therefore highly required. However, feeding ecology and interspecific relationships among nemipterid species is poorly understood. This study therefore aims to investigate diet composition and trophic relationships of eight nemipterid species residing in the southern part of the South China Sea including *Nemipterus mesoprion* (Bleeker, 1853), *Nemipterus hexodon* (Quoy & Gaimard, 1824), *Nemipterus nemurus* (Bleeker, 1857), *Nemipterus nematophorus* (Bleeker, 1854), *Nemipterus tambuloides* (Bleeker, 1853), *Nemipterus peronii* (Valenciennes, 1830), *Nemipterus furcosus* (Valenciennes, 1830) and *Scolopsis taeniopterus* (Cuvier, 1830). This information will be useful for the understanding of natural phenomenon and may serve as useful data for future management of these fishery resources.

## Materials and Methods

### Study Site and Sample Collection

The study area was located between 6° 41' 42" to 9° 18' 10.8"N and 100° 2' 13.2" to 102° 3' 7.2"E in the lower part of the South China Sea (Fig. 1). A total of 22 sampling stations were set based on four depth contours (15-25 m, 25-35 m, 35-45 m, 45-55 m). Three replicated bottom trawling cruises were conducted at each station by the MV PRAMONG 9 research vessel during 22-30 April 2015, 26 May-4 June 2015 and 19-29 July 2015. The bottom-trawl net used was made by nylon with the headline of 39 m, mesh size of 40 mm and mesh size at the cod-end of 25 mm. The duration of trawl hauls was 60 minutes. Data such as longitude, latitude, depth of water, trawl duration and towing distance were recorded. From eight target fish species including *N. mesoprion*, *N. hexodon*, *N. nemurus*, *N. nematophorus*, *N. tambuloides*, *N. peronii*, *N. furcosus* and *S. taeniopterus*, a total of 476 specimens, were sorted from the whole catches, identified, counted, weighted, frozen and brought back to



laboratory for further analysis. Additional samples containing 430 fishes were collected from bottom gill net fisheries at five main fishing ports nearby trawling areas including Surattani, Nakornsritammarat, Songkhla, Pattani and Narathiwat provinces covering the lower part of the South China Sea.

### Laboratory Analysis

Frozen fish samples were thawed, length and weight were measured using a caliper and weight balance to the nearest centimeter and gram, respectively. The fish samples were classified into four different size classes (S1 =  $\leq 7.1$  cm, S2 = 7.1-14 cm, S3 = 14.1-21.0 cm, and S4  $> 21.0$  cm). They were gutted open by a surgical ocular scissors. Sexes were identified by gonad observation (Dan, 1977) and maturity stages (immature, maturing, early matured and fully matured stages) were recorded (modified from Raje, 1996 and Dan, 1977). Fish stomachs were removed and immediately preserved with 10% buffer formalin for a week, drenched overnight with freshwater and preserved in 70% ethanol separately (Hajisamae, 2009). Stomach was cut-open in a petri dish by a corneal scissor. An OLYMPUS SZ61 microscope was used to examine and identify food contents. Each dietary item was identified to the lowest taxonomic level as possible. Prior to stomach content analysis, stomach fullness (FL) was visually estimated at a scale of 0-10, where 0 was empty and 10 represented completely full with food (modified from Hajisamae, 2009). Prey item weight was taken by using a microbalance with an accuracy of 0.001 g (Sartorius TE 214S).

### Data Analysis

The percentage of index of relative importance (%IRI) was applied to analyze diet composition, attributes and overlap. %IRI was calculated by the following formula (Cortes 1997):

$$\%IRI = \frac{IRI}{\sum IRI} \times 100$$

The Index of Relative Importance (IRI) is calculated by the following formula (Hyslop, 1980):

$$IRI = (\%N + \%W) \times \%FO$$

Where %N is the percentage composition by number, %W is percentage composition by weight and %FO is percentage frequency of occurrence of each prey.

Vacuity index (VI) was calculated using the following formula:

$$VI = \frac{\text{Total number of empty stomach}}{\text{Total number of stomach examined}} \times 100$$

Average number of food items (AF) was an average number of prey items in each stomach.



**Diet breadth** ( $B_i$ ) was calculated using Levin's standardized index (Labropoulou & Papadopoulou-Smith, 1999).

$$B_i = \left( \frac{1}{n-1} \right) \left( \left( \frac{1}{\sum_{i,j=1}^n P_{ij}^2} \right) - 1 \right)$$

Where  $B_i$  = Levin's standardized index for predator, 'i'; ' $P_{ij}$ ' = proportion of diet of predator 'i' that is made up of prey 'j'; 'n' = number of prey categories

**Diet overlap** ( $C_H$ ) was calculated by Simplified Morisita index or Morisita-Horn index (Horn, 1966)

$$C_H = \frac{2(\sum p_{ij}p_{ik})}{\sum p_{ij}^2 + \sum p_{ik}^2}$$

Where  $C_H$  = Morisita-Horn index of overlap between species 'i' and 'k';  $p_{ij}$  = proportion food 'i' of the total food quantity by species 'j';  $p_{ik}$  = proportion food 'i' of the total food used by species 'k' and  $n$  = total number of food item. The rate of overlap was classified as low overlap = 0.0-0.29, moderately overlap = 0.30-0.59 and high overlap (to be biologically significant) = 0.60-1.00 (Langton, 1982).

### Statistical Analysis

One-way analysis of variance (ANOVA) was used to analyze stomach fullness (FL) and average number of food item (AF) for fish of different size classes, stages of maturity and sexes. Log (X+1) transformation of raw data was applied prior to statistical analysis to reduce non-normality. To assess inter-specific trophic relationship, cluster analysis was used. Prior to analysis, the dietary samples were square-rooted transformed. The Bray-Curtis similarity was constructed to form a cluster dendrogram by using a PRIMER statistical package 5.0 (Clarke & Gorley 2001). A one-way analysis of similarities (ANOSIM) was performed to test a significant difference of the grouping on the dendrogram. A similarity percentage (SIMPER) was applied to assess what dietary items make the greatest contribution to the grouping.

## Results

### Food and Dominant Food Items

Shrimp, fish, crab, echinoderm and mollusk were common food items for all species examined. Due to a great portion of shrimp and fish in the diet, they were considered the most important food for these fishes (Table 1). However, levels of contribution of each food items in the diet for each species were different. Shrimp largely dominated the diets of *N. mesoprion* (% IRI = 88.9) but fish highly contributed to the diets of *N. nemurus* (%)



IRI = 89.9%) and *N. tambuloides* (% IRI = 80.8%). Five species including *N. nametaophorus*, *N. peronii*, *S. taeniopterus*, *N. hexodon* and *N. furcosus* ingested almost equal proportion of shrimp and fish with slightly greater value in favour of shrimp. Very small contribution of shrimp was found in the diet of *N. nemurus* (% IRI = 2.8%).

### Dietary Attributes

Trophic attributes, including stomach fullness (FL), total number of food item (TLF), average number of food item (AF), vacuity index (VI) and diet breadth (Bi) are shown in Table 2. It was found that FL ranged from 3.34 for *N. nematophorus* to 8.08 for *N. tambuloides*. Analysis of variance (ANOVA) detected a highly significant difference between FL of eight fish species ( $P < 0.01$ ) (Table 3). TLF ranged from 11 items for *N. peronii* to 37 items for *S. taeniopterus*. AF was highest in the diet of *N. tambuloides* (4.87) and lowest in *N. nematopterus* (2.05). A highly significant difference between AF of the eight fish species was recorded ( $P < 0.01$ ). VI varied from 1.67 for *N. tambuloides* to 22.73 for *N. nemurus*. Bi ranged from 0.01 for *N. mesoprion* to 0.17 for *N. peronii*.

### Impacts of Size, sex and Maturity Stage on FL and Number of Food Items

It was found that FL varied significantly between different size classes of five species including *N. nematophorus*, *N. mesoprion*, *N. nemurus*, *N. tambuloides* and *S. taeniopterus* ( $P < 0.05$ ). Sexes of fish; identified as male, female and unidentified sex, significantly influenced FL of *N. mesoprion* ( $P < 0.01$ ) and *N. peronii* ( $P < 0.05$ ). Maturity stages significantly affected FL in the diets of two species, *N. mesoprion* ( $P < 0.01$ ) and *S. taeniopterus* ( $P < 0.05$ ). Fish size class significantly affected AF of four species including *N. nemurus*, *S. taeniopterus*, *N. hexodon* and *N. Peronii*. Sexes also showed significant impacts on AF of *N. hexodon* ( $P < 0.05$ ) and *N. peronii* ( $P < 0.05$ ). There was no impact of maturity stage on AF of all species ( $P > 0.05$ ). Details of statistical results are in Table 3.

### Diet Overlap and Inter-Specific Relationship

Out of 28 pairs of dietary overlap for eight fish species, 16 pairs or 57.5% had the values  $> 0.6$ , considered to be biologically significant (Table 4). Very low value of overlaps was found between *N. nemurus* with *N. mesoprion* and *N. tambuloides* with all other seven species. It was confirmed with a clustering dendrogram that *N. nemurus* and *N. tambuloides* formed a separate group (Figure 2). Four species formed the cluster G1 on the dendrogram including *N. hexodon*, *S. taeniopterus*, *N. mesoprion* and *N. nematophorous* indicating their preference over similar suite of food items. *N. peronii* and *N. furcosus* formed the cluster G2 on the dendrogram. Analysis of similarity (ANOSIM) indicated that classification significantly separated the dietary samples of these species into two distinct groups (Global  $R = 0.786$ ,  $P < 0.005$ ). Similarity percentage (SIMPER) showed that unidentified shrimp (26.4% contribution), unidentified fish (23.8%), mantis shrimp (11.2%) and *Leiognathus* spp. (9.1%) were the greatest contributors to the formation of cluster G1. Unidentified shrimp, unidentified fish



and crab were the major contributors to cluster G2 with the contributions of 29.6%, 21.3% and 12.6%, respectively. For dietary overlap between each size classes of eight species, out of 210 pairs examined, 79 pairs had the value  $> 0.6$ , indicating significant overlaps. Details of diet overlaps between fishes of different size classes of all species are in Table 5.

## Discussion

This is considered the first work dealing directly with feeding habits of eight nemipterid species residing in the same habitat. With the exception of *N. mesoprion* (Manojkumar, 2008) and *S. taeniopterus* (Hajisamae, 2009), no existing reference for other six species is available. Results from this study indicated that nemipterid fishes or thread fin breams feed on a wide range of food items but with high preference over fish and shrimp. Based on very low value of the overall  $B_i$  and feeding mainly on aquatic animals, they were considered as specific carnivorous fishes. It was also observed that many species were considered piscivorous, feeding on fishes, but the ratio of fishes contributing to the diet was different among species. *N. numurus* and *N. tambuloides* were the most piscivorous species as their diet largely dominated by fishes. *N. mesoprion* was the only species which can be defined as a specific crustacean predator as it fed mainly on mantis shrimp, shrimp and other crustaceans. This finding is coincident with those reported by Rao (1989), Joshi (2005) and Manojkumar (2008) that the main food of *N. mesoprion* was crustaceans. In coastal waters off Visakhapatnam, Rao (1989) reported that *N. mesoprion* fed mainly on crustaceans especially *Parapanope euagora* and teleosts especially *Leiognathus* sp. Joshi (2005) observed a large portion of *Stolephorus* spp. and *Leiognathus* spp. in the diet of *N. mesoprion* off Cochin, India. Raje (1996) noticed that *N. mesoprion* off Veraval, India was a carnivorous species, and its diet composed of crustaceans, fish, mollusks and annelids. Similar observations for the diet of this species were reported by Zacharia and Nataraja (2003) and Manojkumar (2008). Historically, feeding habits of *N. mesoprion* was long described by Krishnamoorthy (1971) that it was highly predaceous and possibly a sight feeder of crustaceans, mollusks, annelids and echinoderms. For *S. taeniopterus*, Hajisamae (2009) found that polychaete, crab and shrimp were its main food items. However, slightly different observation was made in this study where shrimp and fish were the major component with small contribution of crab. Additionally, some studies reported on diet composition of other nemipterid fishes. Boaden and Kingsford (2012) found a variety of benthic invertebrates in the diets of *S. bilineatus* and observed an ontogenetic shift in prey size. Studies on *N. japonicus* and *N. randalli* indicated that they were predators feeding on benthic organisms (Vinci, 1982 and Rao & Rao, 1991). Thangavelu et al. (2012) found that *N. japonicus* fed mainly on crustaceans including *Acetes* spp., penaeid prawns, crabs, squid, juvenile fishes such as flathead, lizard fish and some fish larvae. Gurlek et al. (2010) found that crustacean, fish, polychaete and mollusk were the main food for *N. randalli*. Apart from these reports, no reference was found on diets for other nemipterid species.

Generally, diets of most fishes change with sizes (Blaber, 1997). This study confirmed this assumption as it was found that FL of fish samples increased with the increasing of predator body size. Results from this study also indicated that larger fishes seemed to be more active and successful predators than the smaller of the same species. It is postulated that the discrete shift in diet with increasing body size is related to morphological



limitations. When nemipterid fishes reach an optimum body size, they may access to a higher trophic resources. There was no relationship between maturity stage and FL of fish, but a decrease of FL was observed at fully matured stage in this study. A small contribution of nylon and plastic materials in the diets of *N. nematophorus* and *N. nemurus* was probably due to incidental ingestion while capturing targeted preys rather than intentional selection.

Dietary overlap is a tool to compare partitioning of food resources consumed between animal species co-existing in the same habitat (Ross, 1986). According to the niche theory, when sympatric animals overlap in the use of shared resources along one dimension, they must differentiate along another resource to co-exist (Hutchinson 1959, MacArthur 1958). Food overlap among different species or different sizes of the same species is one of the important directions to understand fish community organization (Krebs 1989). Results from this study indicated that food overlap between species was high with the exception of the overlap between *N. tambuloides* with other co-existing species.

In conclusion, the eight nemipterid fish species from the lower part of the South China Sea are specific predators feeding mainly on shrimp and fish. The co-existence of these species in the bottom waters of this habitat requires partitioning of available food resources. This information is useful for an understanding of natural phenomena occurring in the ocean and can be useful for future management of these fishery resources.

### Acknowledgement

Financial support for this study was granted by the Graduate School, Prince of Songkla University, Thailand and research vessel was provided by Southern Marine Fisheries & Development Center, Songkhla, Thailand. Authors would love to thank Mrs. Sriwita Chaibundit, Mr. Thanate Sritakon and all crew members of MV Pramong 9 for helping in research vessel and laboratory works. Special thanks to Dr. Fazrul Hisam, and Dr. Prawit Towattana.

### References

- Almeida, P.R. (2003). Feeding Ecology of *Liza ramada* (Risso, 1810) (Pisces, Mugilidae) in a south-western estuary of Portugal. *Estuarine, Coastal and Shelf Science*, 57(1), 313-323. [http://dx.doi: 10.1016/S0272-7714\(02\)00357-8](http://dx.doi: 10.1016/S0272-7714(02)00357-8)
- Baeta, M., & Ramon, M. (2013). Feeding Ecology of Three Species of *Astropecten* (Asteroidea) Coexisting on Shallow Sandy Bottoms of the Northwestern Mediterranean Sea. *Marine Biology*, 160(11), 2781-2795. <http://dx.doi: 10.1007/s00227-013-2270-0>
- Blaber, S.J.M. (1997). *Fish and Fisheries of Tropical Estuaries*, London, Chapman & Hall., 367 pp.
- Blaber, S. J. M. (2000). *Tropical Estuarine Fishes; Ecology, Exploitation and Conservation*, Oxford, London, Blackwell Science., 384 pp.
- Boaden, A.E., & Kingsford, M.J. (2012). Diel Behaviour and Trophic Ecology of *Scolopsis bilineatus* (Nemipteridae). *Coral Reefs*, 31(3), 871-883. <http://dx.doi:10.1007/s00338-012-0903-2>
- Bohórquez-Herrera, J., Cruz-Escalona, V.H., Adams, D.C., & Peterson, M.S. (2015). Feeding Eco-morphology of Seven Demersal Marine Fish Species in the Mexican Pacific Ocean. *Environmental Biology of Fish*, 98(5), 1459-1473. <http://dx.doi: 10.1007/s10641-014-0373-1>
- Brown, J. A. (1985). The Adaptive Significance of Behavioural Ontogeny in Some Centrarchid Fishes. *Environmental Biology of Fish*, 13, 25-34. doi: 10.1007/BF00004853



- Carrete, M., Lambertucci, S.A., Speziale, K., Ceballos, O., Travaini, A., Delibes, M., Hiraldo, F., & Donazar, J.A. (2010). Winners and Losers in Human-Made Habitats: Interspecific Composition Outcomes in Two Neotropical Vultures. *Animal Conservation*, 13(4), 390-398. <http://dx.doi: 10.1111/j.1469-1795.2010.00352.x>
- Clarke, K.R., & Gorley, R.N. (2001). PRIMER Version 6 User Manual/Tutorial., Plymouth, Primer-E Ltd.
- Cortés, E. (1997). A Critical Review of Methods of Studying Fish Feeding Based on Analysis of Stomach Contents: Application to Elasmobranch Fishes. *Canadian Journal of Fisheries and Aquatic Sciences*, 54, 726-738. doi: 10.1139/f96-316
- Cruz-Escalona, V.H., Abitia-Cardenas, L.A., Campos-Davila, L., & Galvan-Magana, F. (2000). Trophic Interrelation of the Three Most Abundant Fish Species from Laguna San Ignacio, Baja California Sur, Mexico. *Bulletin of Marine Science*, 66(2), 361-373.
- Dan, S.S. (1977). Intraovarian Studies and Fecundity in *Nemipterus japonicus* (Bloch). *Indian Journal of Fisheries*, 24(1&2), 48-55.
- ElHaweet, A.E.A. (2013). Biological Studies of the Invasive Species *Nemipterus japonicus* (Bloch, 1791) As a Red Sea Immigrant into the Mediterranean. *Egyptian Journal of Aquatic Research*, 39(2), 267-274. <http://dx.doi: 10.1016/j.ejar.2013.12.008>
- Espinoza, M., Samantha, E.M.M., Tayler, M.C., Aaron, T.F., & Ingo, S.W. (2015). Feeding Ecology of Common Demersal Elasmobranch Species in the Pacific Coast of Costa Rica Inferred from Stable Isotope and Stomach Content Analyses. *Journal of Experimental Biology and Ecology*, 470(2015), 12-25. <http://dx.doi: 10.1016/j.jembe.2015.04.021>
- Galis, F. (1990). Ecological and Morphological Aspects of Changes in Food Uptake through the Ontogeny of *Haplochromis piceatus*. In Hughes, R. N. (Ed.), *Behavioural mechanisms of food selection* (pp. 281-302). Berlin, Springer-Verlag., 886 pp.
- Gurlek, M., Erguden, S., Yaglioglu, D., Turan, F., Demirhan, S., Gurlek, M., Gungor, M., Ozbalcilar, B., & Ozcan, T. (2010). Feeding Habits of Indo-Pacific Species *Nemipterus randalli* Russel, 1986 (Nemipteridae) in Iskenderun Bay, Eastern Mediterranean Sea. *Commission Internationale pour l'Exploration Scientifique de la Méditerranée*, 39, 539.
- Hajisamae, S., Yeasin, P., & Ibrahim, S. (2006). Feeding Ecology of Two Sillaginid Fishes and Trophic Interrelations with Other Co-Existing Species in the Southern Part of South China Sea. *Environmental Biology of Fish*, 76(2), 167-176. <http://dx.doi: 10.1007/s10641-006-9018-3>
- Hajisamae, S. (2009). Trophic Ecology of Bottom Fishes Assemblage along Coastal Areas of Thailand. *Estuarine, Coastal and Shelf Science*, 82(3), 503-514. <http://dx.doi: 10.1016/j.ecss.2009.02.010>
- Hart, P.J.B. (1997). Foraging Tactics: In Godin, J. G. J. (Ed.), *Behavioral ecology of teleost fishes* (pp. 104-133). New York, Oxford University Press., 384 pp.
- Horn, H.S. (1966). Measurement of "Overlap" in Comparative Ecological Studies. *The American Naturalist*, 100, 419-424. <http://dx.doi: 10.1086/282436>
- Hughes, R.N. (1997). Diet Selection. In Godin, J. G. J. (ed), *Behavioral Ecology of Teleost Fishes* (pp. 134-162). New York, Oxford University Press., 384 pp.
- Hutchinson, G.E. (1959). Omage to Santa Rosalia or Why are There So Many Kinds of Animals? *The American Naturalist*, 93, 145-159. <http://dx.doi: 10.1086/282070>
- Hyslop, E. (1980). Stomach contents analysis - A Review of Methods and Their Application. *Journal of Fish Biology*, 17(4), 411-429. <http://dx.doi:10.1111/j.1095-8649.1980.tb02775.x>
- Joshi, K.K. (2005). Biology and Population Dynamics of *Nemipterus mesoprion* (Bleeker) off Cochin. *Indian Journal of Fisheries*, 52(3), 315-322.
- Juanes, F. (1994). What Determines Prey Size Selectivity in Piscivorous fishes? In Stouder, D. J., Fresh, K. L. and R. J. Feller (Eds), *Theory and application in fish feeding ecology* (pp. 79-100). University of South Carolina Press., 390 pp.
- Juanes, F., & Conover, D.O. (1994). Piscivory and Prey Size Selection in Young-of-the-Year Blue Fish: Predator Preference or Size-Dependent Capture Success? *Marine Ecology Progress Series*, 114(1-2), 59-69. <http://dx.doi: 10.3354/meps114059>
- Kaiser, M.J., & Hughes, R.N. (1993). Factors Affecting the Behavioural Mechanisms of Diet Selection in Fishes. *Marine Behaviour and Physiology*, 23(1-4), 105-118. <http://dx.doi: 10.1080/10236249309378860>
- Karpouzi, V.S., & Stergiou, K.I. (2003). The Relationship Between Mouth Size and Shape and Body Length for 18 Species of Marine Fishes and Their Trophic Implications. *Journal of Fish Biology*, 62(6), 1353-1365. <http://dx.doi: 10.1046/j.1095-8649.2003.00118.x>
- Keast, A., & Webb, D. (1966). Mouth and Body Form Relative to Feeding Ecology in the Fish Fauna of a Small Lake, Lake Opinicon, Ontario. *Journal of the Fisheries Research Board of Canada*, 23(12), 1845-1874. <http://dx.doi: 10.1139/f66-175>
- Krebs, C. (1989). Niche Overlaps and Diet Analysis. In Krebs, C.J. (Ed.), *Ecological methodology*. New York, Harper and Row., 620 pp.
- Krishnamoorthy, B. (1971). Biology of Threadfin Bream, *Nemipterus japonicus* (Bloch). *Indian Journal of Fisheries*, 18, 1-12.
- Langton, R.W. (1982). Diet Overlap Between the Atlantic cod *Gadus morhua*, Silver Hake, *Merluccius bilinearis* and Fifteen Other Northwest Atlantic Fin Fish. *Fishery Bulletin*, 80, 745-759.
- Labropoulou, M., & Papadopoulou-Smith, K.N. (1999). Foraging Behavior Patterns of Four Sympatric Demersal Fishes. *Estuarine, Coastal and Shelf Science*, 49 (Suppl. A), 99-108. [http://dx.doi: 10.1016/S0272-7714\(99\)80014-6](http://dx.doi: 10.1016/S0272-7714(99)80014-6)





- Linke, T.E., Platell, M.E., & Potter, I.C. (2001). Factors Influencing the Partitioning of Food Resources Among Six Fish Species in a Large Embayment with Juxtaposing Bare Sand and Sea Grass Habitats. *Journal of Experimental Marine Biology and Ecology*, 266(2), 193-217. [http://dx.doi: 10.1016/S0022-0981\(01\)00356-2](http://dx.doi: 10.1016/S0022-0981(01)00356-2)
- MacArthur, R. (1958). Population Ecology of Some Warblers of Northeastern cCniferous Forests. *Ecology*, 39(4), 599-619. <http://dx.doi:10.2307/1931600>
- Manojkumar, P.P. (2008). Observations on the Food of *Nemipterus mesoprion* (Bleeker, 1853) from Malabar Coast. *Journal of Marine Biology Association of India*, 50(1), 52-56.
- Pianka, E. R. (1969). Sympatry of Desert Lizards (Ctenotus) in Western Australia. *Ecology*, 50, 1012-1030. <http://dx.doi: 10.2307/1936893>
- Persson, L. (1990). Predicting Ontogenetic Niche Shifts in the Field: What Can be Gained by Foraging Theory? In Hughes, R. N., (Ed.), *Behavioural mechanisms of food selection* (pp. 303-321), Berlin: Springer-Verlag., 886 pp.
- Ping, N., Rensie, W.U., & Jing, L.I.U. (2011). First Record of Red Filament Threadfin Bream, *Nemipterus marginatus* (Valenciennes, 1830) (Perciformes, Nemipteridae), from Chinese Waters. *Chinese Journal of Oceanology and Limnology*, 9(6), 1306-1308. <http://dx.doi: 10.1007/s00343-011-0312-3>
- Rao, T.A. (1989). Fishery of Threadfin Breams at Waltair with Notes on Some Aspects of Biology of *Nemipterus mesoprion* (Bleeker). *Journal Marine Biological Association of India*, 31(1&2), 103-19.
- Rao, D.M., & Rao, K.S. (1991). Food and Feeding Behavior of *Nemipterus japonicus* (Bloch) Populations of Visakhapatnam, South India. *Journal Marine Biological Association of India*, 33(1&2), 335-345.
- Raje, S.G. (1996). Some Observations on the Biology of *Nemipterus mesoprion* (Bleeker) from Verval (Gujrat). *Indian Journal of Fisheries*, 43(2), 163-170.
- Ross, S.T. (1986). Resource Partitioning in Fish Assemblages: A Review of Field Studies. *Copeia*, 1986(2), 352-388. <http://dx.doi: 10.2307/1444996>
- Russell, B.C. (1990). FAO Species Catalogue Vol. 12, Nemipterid Fishes of the World. (Threadfin breams, Whiptail breams, Monocle breams, Dwarf monocle breams, and Coral breams). Rome, Food and Agriculture organization of the United Nations., 149 pp.
- Russell, B.C. (1993). A Review of the Threadfin Breams of Genus *Nemipterus* (Nemipteridae) from Japan to Taiwan, with description of a new species. *Japanese Journal of Ichthyology*, 39(4), 295-310. <http://dx.doi: 10.1007/BF02905130>
- Scharf, F.S., Juanes, F., & Rountree, R.A. (2000). Predator Size-prey Size Relationships of Marine Fish Predators: Interspecific Variation and Effects of Ontogeny and Body Size on Trophic Niche Breadth. *Marine Ecology Progress Series*, 208, 229-248. <http://dx.doi: 10.3354/meps208229>
- Stobutzki, I.C., Silvestre, G.T., Talib, A.A., Krongprom, A., Supongpan, M., Khemakorn, P., Armada, N., & Garces, L.R. (2006). Decline of Demersal Fisheries Resources in Three Developing Asian Countries. *Fisheries Research*, 78(2), 130-142. <http://dx.doi:10.1016/j.fishres.2006.02.004>
- Thangavelu, R., Anbarasu, M., Zala, M.S., Mohamed, K.K., Sreenath, K.R., Suresh, K.M., & Shiju, P. (2012). Food and Feeding Habits of Commercially Important Demersal Finfishes off Verval Coast. *Indian Journal of Fisheries*, 59(4), 77-87.
- Trueman, C.N., Johnston, G., O'Hea, B., & MacKenzie, K.M. (2014). Trophic Interactions of Fish Communities at Midwater Depths Enhance Long-Term Carbon Storage and Benthic Production on Continental Slopes. *Proceeding of the Royal Society B, Biological Sciences*, 281(1787), 1-10. <http://dx.doi: 10.1098/rspb.2014.0669>
- Vinci, G.K. (1982). Threadfin Breams (Nemipteridae) Resources along the Kerala Coast with Notes on the Biology of *Nemipterus japonicus*. *Indian Journal of Fisheries*, 29, 37-49.
- Wotton, R.J. (1998). Ecology of Teleost Fishes. Kluwer, Dordrecht, Boston, London, Academic Publishers., 386 pp.
- Zacharia, P.U., & Nataraja, G.D. (2003). Fishery and Biology of Threadfin Bream, *Nemipterus mesoprion* from Mangalore-Malpe. *Indian Journal of Fisheries*, 50(1), 1-10.

**Table 1.** Diet composition of eight nemipterid fishes collected from the lower part of the South China Sea based on percentage by number (%N), percentage by weight (%W), percentage frequency of occurrence (%FO) and percentage of index of relative importance (%IRI)

Food items	<i>N. nematophorus</i>				<i>N. mesoprion</i>				<i>N. nemurus</i>				<i>N. peronii</i>			
	%N	%FO	%W	%IRI	%N	%FO	%W	%IRI	%N	%FO	%W	%IRI	%N	%FO	%W	%IRI
<b>Crab</b>	<b>9.9</b>	<b>17.4</b>	<b>25.0</b>	<b>3.4</b>	<b>11.1</b>	<b>16.5</b>	<b>5.7</b>	<b>2.2</b>	<b>5.3</b>	<b>25.8</b>	<b>4.6</b>	<b>2.5</b>	<b>5.3</b>	<b>19.0</b>	<b>7.4</b>	<b>2.2</b>
Unidentified crab	6.4	8.9	5.5	2.5	5.8	12.0	4.1	1.8	4	19.4	2.9	2.2	1.3	4.8	0.4	0.2
<i>Charybdis</i> sp.	0.2	0.6	0.1	0.0	-	-	-	0.0	-	-	-	0.0	-	-	-	-
<i>Charybdis truncata</i>	0.5	1.3	1	0.0	-	-	-	0.0	-	-	-	0.0	-	-	-	-
<i>Charybdis anisodon</i>	0.2	0.6	0.6	0.0	-	-	-	0.0	-	-	-	0.0	-	-	-	-
<i>Parapanope</i> sp.	0.2	0.6	0.8	0.0	-	-	-	0.0	-	-	-	0.0	-	-	-	-
Portunidae	0.5	1.3	0.5	0.0	4.8	3.8	1.4	0.4	-	-	-	0.0	2.7	9.5	4.7	1.6
<i>Portunus pelagicus</i>	-	-	-	-	-	-	-	-	-	-	-	-	1.3	4.8	2.2	0.4
<i>Thenus orientalis</i>	0.7	1.9	12.3	0.6	-	-	-	-	-	-	-	-	-	-	-	-
<b>Shrimp</b>	<b>36.5</b>	<b>51.9</b>	<b>31.9</b>	<b>54.7</b>	<b>57</b>	<b>54.1</b>	<b>68.9</b>	<b>88.9</b>	<b>9.3</b>	<b>25.8</b>	<b>4</b>	<b>2.8</b>	<b>45.3</b>	<b>57.1</b>	<b>27.2</b>	<b>54.7</b>
Mantis shrimp	1.7	3.9	6.2	0.7	1.3	3.0	2.7	0.2	4	13.0	2.2	1.3	2.7	9.5	8.7	2.5
<i>Penaeus</i> sp.	1.2	2.5	4.1	0.3	-	-	-	-	-	-	-	-	-	-	-	-
<i>Penaeus indicus</i>	-	-	-	-	-	-	-	-	-	-	-	-	1.3	4.7	8.6	1.1
Other shrimps	30.9	43.6	21.6	53.7	54.4	48.9	65.2	88.7	5.3	13.0	1.7	1.5	41.3	42.8	9.9	51.1
Other crustaceans	2.7	1.9	-	0.0	1.3	2.3	1	0.1	-	-	-	-	-	-	-	-
<b>Fish</b>	<b>44.4</b>	<b>35.3</b>	<b>33</b>	<b>40.7</b>	<b>26</b>	<b>27.1</b>	<b>18.7</b>	<b>8.6</b>	<b>80</b>	<b>135.9</b>	<b>68.8</b>	<b>89.9</b>	<b>40</b>	<b>42.8</b>	<b>29.1</b>	<b>32.9</b>
Carangidae	0.5	0.6	0.9	0.0	0.5	0.7	1.8	0.0	16	13.0	15.8	6.8	1.3	4.8	1.8	0.3
Engraulidae	-	-	-	0.0	-	-	-	-	1.3	6.4	5.5	0.7	-	-	-	-
<i>Gazza minuta</i>	0.2	0.6	0.4	0.0	-	-	-	0.0	1.3	6.4	1.1	0.3	-	-	-	-
<i>Ilisha malastoma</i>	-	-	-	-	-	-	-	-	1.3	6.4	7.4	0.9	-	-	-	-
Leiognathidae	0.5	0.6	7.2	0.1	3.2	3.0	4.9	0.4	-	-	-	-	10.7	4.8	5.3	1.8
<i>Leiognathus</i> sp.	1.5	1.9	5.6	0.3	2.1	2.3	4.7	0.2	4	6.4	2.2	0.6	-	-	-	-
<i>Kurtus</i> sp.	-	-	-	-	0.3	0.7	1.3	0.0	2.7	6.4	0.9	0.4	-	-	-	-
Pomacanthidae	0.2	0.6	0.8	0.0	-	-	-	-	-	-	-	0.0	-	-	-	-
<i>Saurida tumbil</i>	0.5	0.6	0.3	0.0	-	-	-	-	4	19.4	9.7	4.4	-	-	-	-
<i>Saurida undosquamis</i>	0.2	0.6	0.4	0.0	-	-	-	-	1.3	6.4	2.8	0.4	-	-	-	-
Other fishes	40.7	29.5	17.4	40.2	19.9	20.3	6	8.0	48	64.6	23.3	75.4	26.7	28.5	19.6	30.8
<b>Mollusk</b>	<b>8.6</b>	<b>14.7</b>	<b>10.9</b>	<b>1.2</b>	<b>5.8</b>	<b>6.8</b>	<b>6.7</b>	<b>0.3</b>	<b>4</b>	<b>19.4</b>	<b>22.6</b>	<b>4.8</b>	<b>9.3</b>	<b>9.5</b>	<b>36.3</b>	<b>10.1</b>
Bivalvia	3.5	5.8	0.7	0.6	1.6	1.5	0.1	0.0	-	-	-	-	-	-	-	-
Loliginidae	0.5	1.3	0.8	0.0	-	-	-	-	-	-	-	-	9.3	9.5	36.3	10.1
<i>Uroteuthis (P) chinensis</i>	0.5	1.3	1.3	0.1	-	-	-	-	-	-	-	-	-	-	-	-
<i>Uroteuthis (P) duvaucelii</i>	-	-	-	-	0.3	0.7	0.9	0.0	-	-	-	-	-	-	-	-
<i>Sepia</i> sp.	-	-	-	-	-	-	-	-	1.3	6.4	8	1.0	-	-	-	-
Teuthidae	1	1.3	7.1	0.3	0.8	0.7	0.1	0.0	2.7	13.0	14.6	3.7	-	-	-	-
Veneridae	2.5	3.3	0.3	0.2	2.4	2.3	0.1	0.1	-	-	-	-	-	-	-	-
Other mollusks	0.7	1.9	0.8	0.1	0.8	1.5	5.5	0.1	-	-	-	-	-	-	-	-
Nylon	0.5	1.3	-	0.0	-	-	-	-	1.3	6.4	0.1	0.1	-	-	-	-



Table 1 (continued).

Food items	<i>S. taeniopterus</i>				<i>N. hexodon</i>				<i>N. fuscus</i>				<i>N. tambuloides</i>			
	%N	%FO	%W	%IRI	%N	%FO	%W	%IRI	%N	%FO	%W	%IRI	%N	%FO	%W	%IRI
<b>Crab</b>	<b>3.0</b>	<b>10.8</b>	<b>2.7</b>	<b>0.9</b>	<b>17.4</b>	<b>27.0</b>	<b>27.7</b>	<b>8.7</b>	<b>5.7</b>	<b>14.9</b>	<b>7.2</b>	<b>1.6</b>	<b>18.9</b>	<b>38.3</b>	<b>6.8</b>	<b>5.1</b>
Unidentified crab	1.9	7.8	1.2	0.6	2.8	8.1	2.4	1.0	2.8	7.5	1.9	0.6	7.7	13.3	2.2	2.9
<i>Charybdis</i> sp.	-	-	-	-	-	-	-	-	-	-	-	-	2	1.7	1.6	0.1
<i>Charybdis anisodon</i>	-	-	-	-	-	-	-	-	-	-	-	-	1.3	5.1	0.5	0.2
<i>Parapanope</i> sp.	-	-	-	-	-	-	-	-	-	-	-	-	1	5.1	0.8	0.2
<i>Portunus pelagicus</i>	1.1	3.1	1.5	0.2	12.4	10.8	13.1	6.3	2.8	7.5	5.2	1.0	6.1	10.0	0.4	1.4
<i>Thenus orientalis</i>	-	-	-	-	1.1	2.8	3	0.3	-	-	-	-	-	-	-	-
<b>Shrimp</b>	<b>37.8</b>	<b>55.2</b>	<b>24.8</b>	<b>50.8</b>	<b>42.1</b>	<b>51.4</b>	<b>25.8</b>	<b>43.1</b>	<b>51.9</b>	<b>55.0</b>	<b>26.3</b>	<b>47.9</b>	<b>22.2</b>	<b>31.7</b>	<b>4</b>	<b>13.0</b>
Mantis shrimp	0.9	3.7	1.8	0.3	3.9	5.4	1.2	0.6	1.9	5.0	6.6	0.7	0.3	1.7	0.2	0.0
<i>Oratosquilla sollicitans</i>	0.3	1.0	0.8	0.0	0.6	2.8	0.4	0.1	-	-	-	-	-	-	-	-
<i>Penaeus indicus</i>	0.7	2.4	4.5	0.3	0.6	2.8	10.4	0.7	1.9	5.0	8	0.9	1	5.1	2	0.3
Other shrimps	28.5	39.9	17.1	48.4	36.5	37.9	11.5	41.5	48.1	45.0	11.8	46.4	20.9	25.0	1.9	12.7
Other crustaceans	7.4	8.2	0.7	1.8	0.6	2.8	2.3	0.2	-	-	-	-	-	-	-	-
<b>Echinoderm</b>	<b>1.3</b>	<b>4.5</b>	<b>7.1</b>	<b>0.6</b>	-	-	-	-	-	-	-	-	-	-	-	-
<i>Holothuria fuscogilva</i>	0.2	0.6	0.6	0.0	-	-	-	-	-	-	-	-	-	-	-	-
<i>Holothuria scabra</i>	0.9	2.7	6.4	0.5	-	-	-	-	-	-	-	-	-	-	-	-
Starfish	0.2	1.0	0.1	0.0	-	-	-	-	-	-	-	-	-	-	-	-
<b>Fish</b>	<b>49.5</b>	<b>39.6</b>	<b>57.9</b>	<b>46.8</b>	<b>37.6</b>	<b>45.9</b>	<b>28.3</b>	<b>46.9</b>	<b>34.9</b>	<b>45.0</b>	<b>38.8</b>	<b>47.4</b>	<b>53.5</b>	<b>68.4</b>	<b>81.7</b>	<b>80.8</b>
<i>Caranx sexfasciatus</i>	-	-	-	-	-	-	-	-	-	-	-	-	13.8	11.7	11.9	6.7
Carangidae	11.5	8.2	36.8	10.5	1.7	5.4	2.1	0.5	0.9	2.5	1.3	0.1	31	38.3	48.7	67.9
<i>Kurtus</i> sp.	0.2	0.6	0.6	0.0	1.1	5.4	4	0.6	0.9	2.5	1.9	0.1	-	-	-	-
<i>Selaroides leptolepsis</i>	1.9	1.4	8.7	0.4	-	-	-	-	-	-	-	-	5.7	10.0	16.2	4.8
Other fishes	35.7	28.5	11.7	35.9	34.8	35.1	22.3	45.8	33.0	40.1	35.5	47.2	3	8.3	4.8	1.4
<b>Mollusk</b>	<b>7.1</b>	<b>12.5</b>	<b>3.5</b>	<b>0.8</b>	<b>2.2</b>	<b>10.8</b>	<b>14.3</b>	<b>1.1</b>	<b>7.5</b>	<b>7.5</b>	<b>27.8</b>	<b>3.0</b>	<b>2.7</b>	<b>8.3</b>	<b>5.8</b>	<b>0.5</b>
Bivalvia	3.8	4.7	0.3	0.5	0.6	2.8	0.2	0.1	-	-	-	-	-	-	-	-
<i>Charonia variegata</i>	-	-	-	-	0.6	2.8	1.8	0.2	-	-	-	-	0.3	1.7	0.1	0.0
Loliginidae	0.2	0.4	0.3	0.0	-	-	-	-	6.6	5.0	27.3	2.9	1.0	1.7	-	0.0
<i>Monoplex intermedius</i>	-	-	-	-	-	-	-	-	0.9	2.5	0.5	0.1	-	-	-	-
<i>Uroteuthis (P) chinensis</i>	0.5	1.0	1.2	0.0	-	-	-	-	-	-	-	-	0.7	3.4	4	0.4
<i>Uroteuthis (P) duvaucelii</i>	-	-	-	-	-	-	-	-	-	-	-	-	0.7	1.7	1.7	0.1
<i>Sepioteuthis lessoniana</i>	-	-	-	-	0.6	2.8	10.5	0.7	-	-	-	-	-	-	-	-
<i>Sepia</i> sp.	0.1	0.4	1.4	0.0	-	-	-	0.0	-	-	-	-	-	-	-	-
Veneridae	1	2.0	0.1	0.1	-	-	-	0.0	-	-	-	-	-	-	-	-
Other mollusks	1.4	4.1	0.3	0.2	0.6	2.8	1.8	0.2	-	-	-	-	-	-	-	-
Urechidae	1.5	1.2	3.9	0.2	0.6	2.8	3.9	0.3	-	-	-	-	2.7	6.6	1.7	0.6



**Table 2.** Trophic attribute of eight nemipterid fishes of different ecological and biological conditions collected from the lower part of the South China Sea. (n = number of stomach analyzed; FL= fullness index; SD= standard deviation; TLF = total number of food item; AF= average number of food item; VI = vacuity index; Bi= diet breadth; S1 = <7cm, S2 = 7.1-14.0cm, S3 = 14.1-21.0cm, S4 = >21cm.)

Species	n	Size Range (cm)	FL± SD	TLF	AF	VI	Bi
<i>N. nematophorus</i>	198	All	4.24±2.98	32	2.05	21.21	0.04
	44	S1	3.36±3.29	8	1.16	36.36	0.21
	148	S2	4.35±2.78	29	2.28	17.57	0.04
	6	S3	8.00±2.45	5	2.67	0.00	0.60
<i>N. mesoprion</i>	171	All	3.34±2.73	20	2.20	22.22	0.01
	7	S1	0.86±1.57	-	-	71.43	D
	127	S2	3.46±2.73	15	1.96	21.26	0.05
	37	S3	3.41±2.74	11	3.30	16.22	0.01
<i>N. nemurus</i>	22	All	5.55±3.95	18	3.41	22.73	0.07
	9	S2	5.11±3.89	8	1.67	22.22	0.68
	13	S3	5.85±4.12	16	4.62	23.08	0.05
<i>N. peronei</i>	23	All	4.39±2.79	11	3.26	8.70	0.17
	13	S2	4.92±2.10	7	2.92	0.00	0.25
	10	S3	3.70±3.50	7	3.70	20.00	0.23
<i>S. taeniopterus</i>	345	All	4.09±2.75	37	3.65	14.49	0.04
	39	S2	4.08±2.25	14	4.97	10.26	0.09
	253	S3	3.47±2.38	28	2.91	17.39	0.03
	53	S4	7.06±2.84	21	6.17	3.77	0.04
<i>N. hexodon</i>	42	All	6.00±3.71	22.00	4.24	11.90	0.07
	8	S2	3.63±2.83	4	2.50	25.00	0.51
	27	S3	6.11±3.79	17	5.44	11.11	0.09
	5	S4	10.00±0.00	9	2.40	0.00	0.85
<i>N. furcosus</i>	44	All	4.89±2.92	14.00	2.41	9.09	0.09
	11	S2	5.82±3.06	6	1.91	9.09	0.13
	33	S3	4.58±2.85	13	2.58	9.09	0.13
<i>N. tambuloides</i>	61	All	8.08±2.48	20.00	4.87	1.67	0.05
	6	S2	4.33±2.66	5	1.67	16.67	0.47
	36	S3	8.64±1.96	15	5.56	0.00	0.08
	19	S4	8.21±2.33	10	4.58	0.00	0.10

**Table 3.** Statistical results for the effects size, sex and maturity stage on stomach fullness index and number of food items in the stomachs of eight nemipterid fishes collected from the lower part of the South China Sea

Factors	Species	No. of sample	df	Fullness index	No. of food items
Species	All	906	7	P<0.01	P<0.01
Size	<i>N. nematophorus</i>	198	2	P<0.01	P>0.05
	<i>N. mesoprion</i>	171	2	P<0.05	P>0.05
	<i>N. nemurus</i>	22	-	P<0.05	P<0.05
	<i>N. tambuloides</i>	61	2	P<0.01	P<0.05
	<i>S. taeniopterus</i>	345	2	P<0.01	P>0.05
	<i>N. hexodon</i>	42	2	P>0.05	P>0.05
	<i>N. furcosus</i>	44	-	P>0.05	P<0.05
	<i>N. peronii</i>	23	-	P>0.05	P<0.05
Sex	<i>N. nematophorus</i>	198	2	P>0.05	P>0.05
	<i>N. mesoprion</i>	171	2	P<0.01	P>0.05
	<i>N. nemurus</i>	22	2	P>0.05	P>0.05
	<i>N. tambuloides</i>	61	-	P>0.05	P>0.05
	<i>S. taeniopterus</i>	345	-	P>0.05	P>0.05
	<i>N. hexodon</i>	42	2	P>0.05	P<0.05
	<i>N. furcosus</i>	44	2	P>0.05	P>0.05
	<i>N. peronii</i>	23	-	P<0.05	P<0.05
Maturity stage	<i>N. nematophorus</i>	198	3	P>0.05	P>0.05
	<i>N. mesoprion</i>	171	3	P<0.01	P>0.05
	<i>N. nemurus</i>	22	3	P>0.05	P>0.05
	<i>N. tambuloides</i>	61	3	P>0.05	P>0.05
	<i>S. taeniopterus</i>	345	3	P<0.05	P>0.05
	<i>N. hexodon</i>	42	3	P>0.05	P>0.05
	<i>N. furcosus</i>	44	2	P>0.05	P>0.05
	<i>N. peronii</i>	23	2	P>0.05	P>0.05

**Table 4.** Morisita-Horn indices for the diets of different nemipterid fishes collected from the lower part of the South China Sea. The value >0.60 are highlighted in bold indicating significant overlap

Species	Species							
	Sp1	Sp2	Sp3	Sp4	Sp5	Sp6	Sp7	Sp8
<i>N. nematophorus</i> (Sp1)	-							
<i>N. mesoprion</i> (Sp2)	0.75	-						
<i>N. nemurus</i> (Sp3)	0.59	0.11	-					
<i>N. tambuloides</i> (Sp4)	0.15	0.18	0.14	-				
<i>S. taeniopterus</i> (Sp5)	0.98	0.83	0.54	0.33	-			
<i>N. hexodon</i> (Sp6)	1.00	0.79	0.63	0.16	0.98	-		
<i>N. furcosus</i> (Sp7)	0.99	0.88	0.48	0.17	0.98	0.98	-	
<i>N. peronii</i> (Sp8)	0.98	0.82	0.52	0.16	0.97	0.98	0.99	-



**Table 5.** Morisita-Horn indices for the diets of different size class of nemipterid fishes collected from the lower part of the South China Sea. (S1 = <7cm, S2 = 7.1-14.0cm, S3 = 14.1-21.0cm, S4 = >21cm. The value >0.60 are highlighted in bold indicating significant overlap


Species	Size	<i>N. nematophorus</i>			<i>N. mesoprion</i>		<i>N. nemurus</i>		<i>N. tambuloides</i>			<i>S. taeniopterus</i>			<i>N. hexodon</i>				<i>N. furcosus</i>		<i>N. peronii</i>		
		S1	S2	S3	S2	S3	S2	S3	S2	S3	S4	S2	S3	S4	S1	S2	S3	S4	S2	S3	S2	S3	
<i>N. nematophorus</i>	S1	-																					
	S2	1.0	-																				
	S3	0.4	0.4	-																			
<i>N. mesoprion</i>	S2	0.9	0.9	0.4	-																		
	S3	0.8	0.8	0.3	0.2	-																	
<i>N. nemurus</i>	S2	0.4	0.3	0.1	0.2	0.1	-																
	S3	0.5	0.6	0.2	0.3	0.0	0.4	-															
<i>N. tambuloides</i>	S2	0.7	0.7	0.3	0.5	0.3	0.4	0.9															
	S3	0.1	0.1	0.1	0.2	0.1	0.2	0.1	0.0	-													
<i>S. taeniopterus</i>	S4	0.1	0.1	0.1	0.2	0.2	0.2	0.1	0.0	1.0	-												
	S2	0.9	0.9	0.3	0.7	0.5	0.4	0.8	0.9	0.1	0.1	-											
	S3	1.0	1.0	0.4	1.0	0.9	0.2	0.4	0.6	0.2	0.2	0.8	-										
<i>N. hexodon</i>	S4	0.2	0.2	0.1	0.2	0.1	0.3	0.2	0.1	1.0	1.0	0.2	0.2	-									
	S1	0.2	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-								
	S2	0.9	1.0	0.4	0.8	0.6	0.4	0.7	0.8	0.3	0.3	1.0	0.9	0.3	0.0	-							
<i>N. furcosus</i>	S3	1.0	1.0	0.4	0.9	0.7	0.3	0.6	0.8	0.1	0.1	0.9	0.9	0.2	0.0	1.0	-						
	S4	0.2	0.2	0.3	0.2	0.1	0.1	0.2	0.2	0.0	0.0	0.2	0.2	0.0	0.0	0.2	0.2	-					
	S2	0.9	0.9	0.3	1.0	1.0	0.1	0.2	0.4	0.2	0.2	0.7	1.0	0.1	0.0	0.8	0.9	0.2	-				
<i>N. peronii</i>	S3	1.0	1.0	0.4	0.9	0.8	0.3	0.6	0.8	0.1	0.1	0.9	1.0	0.2	0.0	1.0	1.0	0.2	0.9	-			
	S2	0.8	0.9	0.3	0.7	0.4	0.4	0.9	0.9	0.1	0.1	1.0	0.7	0.2	0.0	0.9	0.9	0.2	0.6	0.9	-		
	S3	0.8	0.8	0.4	0.9	0.9	0.1	0.1	0.4	0.1	0.1	0.6	0.9	0.1	0.0	0.7	0.8	0.2	0.9	0.8	0.5	-	



Figure 1. Study area along the lower part of the South China Sea.

Accepted

27



**Figure 2.** Cluster dendrogram indicating inter-specific relationship between eight sympatric nemipterid fish species collected from the lower part of the South China Sea. (G1 = group 1 and G2 = group 2)

Accepted Manuscript