



Species Identification Using DNA Barcoding on Processed Panga Catfish Products in Viet Nam Revealed Important Mislabeling

Tran Thi Thuy Ha^{1,*}, Nguyen Thi Huong, Nguyen Phuc Hung, Yann Guiguen

¹Research Institute for Aquaculture No1, Centre for Aquaculture Biotechnology, Bac Ninh, Viet Nam

E-mail: thuyha@ria1.org

Abstract

Nucleotide sequences of the Cytochrome oxidase subunit I gene and Cytochrome b were analyzed for 10 processed fish products collected from supermarkets in Hanoi, Viet Nam. The similarity between our results and published data from the NCBI and BOLD was compared to identify species. This molecular analysis showed that the common names of only 4 of these products matched with their corresponding scientific names. The other six were mislabeled with an important mislabeling from *P. hypophthalmus* into *P. boucourti*. Although no commercial frauds were found in these mislabelled products, the correct scientific names of fish species should be labelled for the processed products as they are in supermarkets.

Keywords: Catfish, processed products, species identification

Introduction

Mislabeling seafood products can represent an economic fraud and may lead to potential health risks for consumers (<http://www.cbc.ca/news/mislabelling-means-rare-fish-sold-marketplace-1.919822>). A commonly economic fraud is for instance the commercialization of a low-priced species using the name of a high-priced species (Hellberg and Morrissey, 2011). The use of a false or misleading name may affect the ability of processors and consumers to make accurate assessments of the potential safety hazards associated with seafood. In fact, hazards such as allergenic proteins and scombrototoxin formation are associated with some species but not others, presenting potential food safety risks if the food is not accurately labeled (<http://www.foodsafetynews.com/2012/12/seafood-fraud-public-health-threat-or-economic-trick>). It has been reported that seafood products were abundantly mislabeled in the world (Carvalho et al., 2010; Filonzi et al., 2010; Barbuto et al., 2010) but this occurrence of seafood mislabeling has not been studied in Viet Nam to our knowledge.

Fish species can be identified from unprocessed products based on external morphological characteristics, such as body shape, number of fins or scales, texture or filet colors. However, it is often difficult for consumers to accurately determine fish species from processed products since the morphological features changed after processing. In addition, the identification of fish species based on DNA analysis usually encounters some difficulties due to a large size of genome or genetic variation. To date, many molecular markers have been studied and effectively used as a tool for identification of fish species (Ward *et al.*, 2009). Such markers should be ideally highly conserved in the same species within different populations and should be also clearly different among species. DNA barcoding is a



potential approach for such an identification of fish species as it is widely used for studies of the genetic diversity and the classification of species characterized by morphological similarities (Hebert *et al.* 2003). The core assumption of DNA barcoding is that the nucleotide sequence similarities are lower within a species than between different species (Meyer and Paulay, 2005). Typically, mitochondrial genome genes are often used for DNA barcoding. The mitochondrial genome has a higher rate of mutations compared to the nuclear genome, it is maternally inherited with less hybridization and a higher copy number, facilitating PCR amplification and sequence recovery from degraded tissue (Saccone *et al.*, 1999; Hebert *et al.*, 2004). Furthermore, mitochondrial genome genes lack introns, pseudogenes and repetitive sequences facilitating sequence alignments of the amplified genes (Lin *et al.*, 2005). Finally, many complete mitochondrial DNA genome sequences are publicly available and primers can therefore be designed to amplify and sequence mitochondrial genes in any species with a published mtDNA genome (Folmer *et al.*, 1994).

DNA barcoding could be used to monitor the illegal trade of wildlife, such as protected or endangered species as well as identify the species origin of commercially processed food (Dawnay *et al.*, 2007; Marko *et al.*, 2004). Until now, a number of studies have shown the applicability of DNA barcoding for accurate identification of a wide range of fish species (Ward *et al.*, 2005; Ward *et al.*, 2009; Hubert *et al.*, 2008). Besides, the combination of some genes for identified species becomes the preferred method. Nicole *et al.* (2012) identified species from fish processed products based on three distinct mitochondrial genes (16S-rDNA, COI and Cytb). Among the classical DNA barcoding mitochondrial markers used for animal species, the COI gene, which encodes for the cytochrome oxidase subunit I, was considered as a suitable marker because its mutation rate is often fast enough to distinguish closely related species and also its sequence is conserved among conspecifics. The *cyt b* sequence encoding Cytochrome b, shows considerable variation and allows for the differentiation of even closely related species due to its relatively high interspecies variation and low intraspecies variation (Mackie and others 1999; Aranishi and others 2005a).

The present study aimed at using COI and Cytb genes to identify fish species in some processed seafood products available in markets in Vietnam. The DNA of 37 seafood products, including raw processed products or frozen and fillet products were extracted successfully. Genes were amplified with an average length of 700 bp for COI; 500 bp for 16S-rDNA and 850 bp for the Cytb. Results showed that 16 seafood products were correctly labeled based on the results obtained with these three gene markers, 12 samples were correctly labeled with two gene markers and 5 samples were correctly labeled with only one gene marker.

Materials and Methods

Processed Fish Products

Twenty samples from ten processed fish products were collected from two well-known supermarkets, namely Big C Long Bien and Aeon Mall Long Bien in Hanoi, Viet nam. These products were labeled in the supermarkets as being processed from the following species: *Pangasius hypophthalmus*, *Pangasius bocourti*,



Prionace glauca and *Oncorhynchus mykiss* (Table 1). The processed products, which had been frozen at -20°C at the supermarkets, were labelled with both common and scientific names. Details of names of these products, their processing method and origin is presented in Table 1.

DNA Extraction

Total genomic DNA was extracted from the 20 samples, using the DNeasy mericon Food Kit (Qiagen, Germany) according to the manufacturer's instructions. After extraction, the quantity and quality of DNA was determined using 0.8% agarose gel and Nanodrop 2000C (Thermo Scientific, Country).

Amplification of COI and Cytb Genes

To amplify the COI genes, we used MAB primers (MAB F and MAB R) and FISH primers as described in Badhul Haq *et al.* (2012) and Ward *et al.* (2005) and the primers designed by Russell *et al.* (2000) and Wolf *et al.* (2000) were used for the Cytb genes (Table 2)

PCR reactions were carried out with a total volume of 25 μl , including 3 μl of DNA (100 ng l^{-1}), 0.5 μl of 100 mM Tris HCl (pH 8.3), 0.5 μl of 500 mM KCl (pH 8.3), 2.5 μl of MgCl_2 (25 mM), 1.0 μl of dNTPs (5 mM), 0.5 μl of each forward and reverse primers ($10\text{ pm } \mu\text{l}^{-1}$ per primer), and 1 μl of Taq Polymerase. The cycling conditions were: 94°C for 2 min; 35 cycles at 94°C for 50 s; $50\text{-}56^{\circ}\text{C}$ for 50 s; 72°C for 1 min ; 72°C for 10 min and kept at 4°C . Amplification reaction was performed on PCR Mastercycler Pro S (Manufacturer, Place, Country).

Sequences of COI and Cytb Genes

PCR products were analyzed on 2% agarose gels by electrophoresis. Before the sequencing step, all products were purified by using ExpinTM PCR SV kit (GeneAll). Then, the qualified PCR products were sequenced using the BigDye Terminator v3.1 Cycle Sequencing kit at First BASE Laboratories, Malaysia. The solution (10 μl) for reaction contained 4.94 μl of pure water, 1.94 μl of BigDye buffer $5 \times$ (400mM Tris HCl pH 9.0 and 10 mM MgCl_2), 0.12 μl of BigDye Terminator and 1 μl of an ExoSAP product. Then, the bidirectional analysis of sequence on Applied Biosystems machine was carried out. Genomelab software was used to analyse DNA sequences and gene sequences were checked using TV Finch 1.4.0 software (<http://www.geospiza.com>). ClustalW program implemented in BioEdit was used to compare the sequences.

Identification of the Fish Species

Species identification was based on sequence comparisons with reference gene sequences in the GenBank database using BLASTn program. Homologous sequences were compared with reference sequences using the following parameters: Coverage (coverage of the length of the comparing sequence), E-value (reliability of the homology) and Identity (homologous comparison between two sequences). In additional, we also used the BOLD



database (The Barcode of Life Data System) to determine the accuracy for identifying species using as threshold for non-similarity nucleotide sequences, which were less than 1% (Ratnasingham *et al.*, 2007).

Results

PCR

PCR products showed that the bands were relatively clear and sharp (Fig. 1 & 2). Moreover, the bands of all samples were in the range of 500 bp and 700 bp as described by Ward *et al.* (2005) and Badhul Haq *et al.* (2012).

Identification of the Fish Species

A broad species identification of the studied processed products was developed based on BOLD and NCBI databases. Most of the identified fishes could be verified from the present database. The summarised form of the fish identification of cytochrome oxidase I and cytochrome b gene sequences of the 20 different products is shown (Table 3).

Discussion

Our results showed that 20 samples from 10 different processed fish products have a great similarity when compared on the NCBI and BOLD. These products include ground and roasted *P. hypophthalmus* with cinnamon (CQ2), dry fillet of *P. hypophthalmus* (KC2), ball of ground and roasted *P. bocourti* (CCV1, CCV2), ball of ground and roasted *P. bocourti* with dill (TL2), ball of ground *P. bocourti* (VB1), fillet of *P. hypophthalmus* rolled with flour (LB1, LB2), ball of ground *P. bocourti* (FB1, FB2), roll of roasted *Oncorhynchus mykiss* (CH1, CH2), fillet of *Prionace glauca* (CM1, CM2). Both samples of ground and grilled *P. bocourti* (CC1, CC2) were not found on the BOLD database, however, the analysis on the NCBI of these samples showed a relatively high similarity (99%). In addition, the sequences of Cytb of 20 samples confirmed our results with the COI gene. Hence, the accuracy of these analysis results could be acceptable to identify the species.

In the present study, 7 samples (KC2, LB1, LB2, CH1, CH2, CM1, CM2) from 4 processed fish products were labeled correctly based on the high similarity to GeneBank reference COI and Cytb (99% to 100% when compared on NCBI and 100% when compared on BOLD). However, 12 samples (CC1, CC2, CQ1, CQ2, CCV1, CCV2, VB1, VB2, TL1, TL2, FB1, FB2) from 6 processed fish products presented differences with their GenBank references suggesting that these products were mislabeled. Even if our sampling size (20 samples) is not large enough to account for significant percentage this represent an overall percentage of 60% of mislabeled products suggesting a very large scale mislabeling of these fish products.

Incorrect labeling were mostly detected within the *Pangasius* catfish family and most mislabelling were found in *P. hypophthalmus* products mislabeled as *P. bocourti*. In fact, the price of *P. bocourti* catfish is cheaper



than that of *P. hypophthalmus* catfish. Therefore, commercial fraud issues for processed fish products can be eliminated in the present study. The mislabeling of these products could be due instead to simple confusion between names of these two fish species. Producers might not accurately check the scientific name of the raw fish materials before processing them and unintentionally confuse their names. Mislabeling has been reported in many countries and markets. Carvalho *et al.* (2010) revealed that 80% of the surubim (*Pseudoplatystoma spp.*) whole fish fillet products sold in the Brazilian market was mislabeled. To solve that problem this country had published a formal list of common names of fish species used in processed products (Carvalho *et al.*, 2011).

DNA barcodes have been used to identify fish species in numerous products made from different species, such as tuna (Terol *et al.*, 2002), cod (Espineira *et al.*, 2008), anchovy (Jérôme *et al.*, 2008) and shark (Barbuto *et al.*, 2010). Filonzi *et al.* (2010) have used DNA barcode for COI and cytb genes in some catfish products and showed that 32% of these fish products were mislabeled, including 26% of errors with closely related species. Their conclusion was that the commercial fraud on these products was high (~ 42%).

Conclusion

Analysis and comparison of sequence of COI and Cytb gene fragments with the NCBI gene references and the BOLD databases showed that 40% of processed fish products were correctly labeled while 60% of the total processed fish products have been recorded as mislabeled. Most of the mislabeling products were due to confusion of scientific names between *P. bocourti* and *P. hypophthalmus* and no commercial frauds were found. These findings suggest that it is necessary to better determine fish species in the processed products and to help transformers and producers to correctly label their fish-processed products.

Acknowledgement

This study was conducted with the financial support from the national project “Research on the development and application of DNA barcoding on catfish (*Pangasianodon hypophthalmus*)” funded by Ministry of Agriculture and Rural Development, Viet Nam.

References

- Badhul, H.M.A., Azarudeen, M. D., Vignesh, R., Kumar, A. T.T. & Srinivasan, M. (2012). Identification and intra species delineation of ornamental silver pompano *Trachinotus blochii* (Lacépède, 1801) with DNA barcodes. International Research Journal of Biochemistry and Bioinformatics 3: 26-36. <http://www.interestjournals.org/IRJBB>
- Barbuto, M., Galimberti, A., Ferri, E., Labra, M., Malandra, R., & Galli, P. (2010). DNA barcoding reveals fraudulent substitutions in shark seafood products: The Italian case of “palombo” (*Mustelus spp.*). Food Research International, 43, 376-381. <https://doi.org/10.1016/j.foodres.2009.10.009>
- Carvalho, D.C., Neto, D.A., Brasil, B.S., & Oliveira, D.A. (2011). DNA barcoding unveils a high rate of mislabeling in a commercial freshwater catfish from Brazil. Mitochondrial DNA 22(1): 97-105. <http://doi.org/10.3109/19401736.2011.588219>
- Dawnay, N., Ogden R., McEwing R., Carvalho G.R., & Thorpe R.S., (2007). Validation of the barcoding gene COI for use in forensic genetic species identification. Forensic Sci. Intl. 173, 1-6. <http://dx.doi.org/10.1016/j.forsciint.2006.09.013>
- Espineira, M., Nerea González-Lavín, N., Vieites, J.M., & Santaclara, F. (2008). Development of a method for the genetic identification of flatfish species on the basis of mitochondrial DNA sequences. Journal of Agriculture and Food Chemistry,



- 56, 8954-8961. <http://doi.org/10.1021/jf800570r>
- Filonzi, L., Chiesa, S., Marina V., & Francesco N.M. (2010). Molecular barcoding reveals mislabeling of commercial fish products in Italy. *Food Research International* Volume 43, Issue 5, June 2010, Pages 1383-1388. <https://doi.org/10.1016/j.foodres.2010.04.016>
- Folmer, O., Black, M., Hoeth, W., Lutz R., & Vrijenhoek, R. (1994). DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology Journal* 3, 294-299. <https://vi.scribd.com/document/261688092/>
- Hebert, P.D.N., Cywinska, A., Ball, S.L., & de Waard, J.R. (2003). Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London. Series B: Biological Sciences* 270, 313-321. <https://doi.org/10.1098/rspb.2002.2218>
- Hebert, P.D.N., Stoeckle, M.Y., Zemlak, T.S., & Francis, C.M. (2004). Identification of birds through DNA barcodes. *PLOS Biology*, 2, 1657-1663. <https://doi.org/10.1371/journal.pbio.0020312>
- Hellberg, R.S., & Morrissey, M.T. (2011). Advances in DNA-based techniques for the detection of seafood species substitution on the commercial market Part B: *Biological Sciences* 270, 313-322. <https://doi.org/10.1016/j.jala.2010.07.004>
- Hubert, N., Hanner, R., Holm, E., Mandrak, N.E., Taylor, E., Burrige, M., ... Bernatchez, L. (2008). Identifying Canadian freshwater fishes through DNA barcodes. *PLOS One* 3 (6), 2490. <https://doi.org/10.1371/journal.pone.0002490>
- Jérôme, M., Martinsohn, J. T., Ortega, D., Carreau, P., Verrez -Bagnis, V., & Mouchel, O. (2008). Toward fish and seafood traceability: Anchovy species determination in fish products by molecular markers and support through a public domain database. *Journal of Agriculture and Food Chemistry*, 56, 3460-3469. <https://doi.org/10.1021/jf703704m>
- Lin, W.F., Shiau, C.Y., & Hwang, D.F. (2005). Identification of four *Thunnus* tuna species using mitochondrial cytochrome b gene sequence and PCR-RFLP analysis. *Journal of Food Drug Analysis*, 13, 382-387. <https://www.researchgate.net/publication/282736365>
- Marko, P.B., Lee, S.C., Rice, A.M., Gramling, J.M., Fitzhenry, T.M., McAlister, J.S., Harper, G.R., & Moran, A.L. (2004). Mislabeling of a depleted reef fish. *Nature*, 430, 309-310. <https://doi.org/10.1038/430309b>
- Meyer, C.P., & Paulay G. (2005). DNA barcoding: error rates based on comprehensive sampling. *PLOS Biology*, 3, 422. <https://doi.org/10.1371/journal.pbio.0030422>
- Nicole, S., Negrisolo, E., Eccher, G., Mantovani, R., Patarnello, T., Erickson, D.L., Kress, W.J., & Barcaccia, G. (2012). DNA Barcoding as a Reliable Method for the Authentication of Commercial Seafood Products. *Smithsonian Institution*, P.O. Box 37012, Washington, DC 20013-7012, USA. <http://www.academia.edu/23602217/>
- Ratnasingham, S., & Hebert, P.D.N. (2007). BOLD: The Barcode of Life Data system. *Molecular Ecology Notes*, 7, 355-36. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1890991/>
- Saccone, C., De Carla, G., Gissi, C., Pesole, G., & Reyes, A. (1999). Evolutionary genomics in Metazoa: the mitochondrial DNA as a model system. *Gene*, 238, 195-210. [https://doi.org/10.1016/S0378-1119\(99\)00270-X](https://doi.org/10.1016/S0378-1119(99)00270-X)
- Terol, J., Mascarell, R., Fernandez-Pedrosa, V., & Perez-Alonso, M. (2002). Statistical validation of the identification of Tuna species: Bootstrap analysis of mitochondrial DNA sequences. *Journal of Agriculture and Food Chemistry*, 50, 963-969. <https://doi.org/10.1021/jf011032o>
- Wallace, L.J., Boilard, S.M.A.L., Eagle, S.H.C., Spall, J.L., Shokralla, S., & Hajibabaei, M. (2012). DNA barcodes for everyday life: routine authentication of Natural Health Products. *Food Research International* 49(1), 446-452. <https://doi.org/10.1016/j.foodres.2012.07.048>
- Ward, R.D., Hanner, R., & Hebert, P.D.N. (2009). The campaign to DNA barcode all fishes, FISH-BOL. *Journal of Fish Biology*, 74, 329-356. <https://doi.org/10.1111/j.1095-8649.2008.02080.x>
- Ward, R.D., Zemlak, T. S., Innes, B. H., Last, P. R. & Hebert, P. D. (2005). DNA barcoding Australia's fish species. *Philosophical Transactions of the Royal Society B*, 360, 1847-1857. <https://doi.org/10.1098/rstb.2005.1716>
- Woese, C.R., & Fox, G.E. (1977). Phylogenetic structure of the prokaryotic domain: the primary kingdoms. *Proceedings of the National Academy of Sciences of the United States of America*, 97, 8392-8396. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC432104/>

**Table 1.** Main information of 20 samples from 10 processed fish products collected in supermarkets in Viet Nam

No	Code of sample	Name of product	Scientific name of fish	Processing method	Origin
1	CCV1, CCV2	Ball of ground and roasted <i>Pangasius bocourti</i>	<i>Pangasius bocourti</i>	Fish ground and roasted	Vietnam
2	TL1, TL2	Ball of ground and Roasted <i>Pangasius bocourti</i> with dill	<i>Pangasius bocourti</i>	Fish ground and Roasted with dill	Vietnam
3	CQ1, CQ2	Ground and roasted <i>Pangasius hypophthalmus</i> with cinnamon	<i>Pangasius hypophthalmus</i>	Fish ground and Roasted with cinnamon	Vietnam
4	KC1, KC2	Dry fillet of <i>Pangasius hypophthalmus</i>	<i>Pangasius hypophthalmus</i>	Fish filleted and dry	Vietnam
5	CC1, CC2	Ground and grilled <i>Pangasius bocourti</i>	<i>Pangasius bocourti</i>	Fish ground and grilled	Vietnam
6	LB1, LB2	Fillet of <i>Pangasius hypophthalmus</i> rolled with flour	<i>Pangasius hypophthalmus</i>	Fish filleted and rolled with flour	Vietnam
7	VB1, VB2	Ball of ground <i>Pangasius bocourti</i>	<i>Pangasius bocourti</i>	Fish ground and boiled	Vietnam
8	FB1, FB2	Burger of <i>Pangasius bocourti</i>	<i>Pangasius bocourti</i>	Fish ground and roasted	Vietnam
9	CH1, CH2	Roll of roasted <i>Oncorhynchus mykiss</i>	<i>Oncorhynchus mykiss</i>	Fish filleted, rolled and roasted	Vietnam
10	CM1, CM2	Fillet of <i>Prionace glauca</i>	<i>Prionace glauca</i>	Fish filleted	Vietnam

Table 2. Characteristics of primers

Primers	Sequence (5'-3')	Annealing temperature	size of the amplified fragments	Amplified products in this study
MAB	F:TCAACCAACCACAAAGACATTGG CAC R:TAGACTTCTGGGTGGCCAAAGAA TCA	50 °C	680bp	CM1, CM2, CH1, CH2
Fish	F:CGACTAATCATAAAGATATCGGCA C R:TTCAGGGTGACCGAAGAATCAGA A	56°C	680bp	CCV1, CCV2, TL1, TL2, CQ1, CQ2, KC1, KC2, CC1, CC2, LB1, LB2, VB1, VB2, FB1, FB2
Cytb	AAAAACCACCGTTGTTATTCAACTA GCICCTCARAATGAYATTTGTCCT CA	53°C	430bp	CCV1, CCV2, TL1, TL2, CQ1, CQ2, KC1, KC2, CC1, CC2, LB1, LB2, VB1, VB2, FB1, FB2, CM1, CM2, CH1, CH2,

**Table 3.** Identification of fish species by the analysis of COI and Cytb genes and comparison the similarity to NCBI and BOLD databases

Code of sample	Name of product	COI				Cyt b		
		Species similarity	Similarity %		Note	Species similarity NCBI	Similarity % NCBI	Note
			NCBI	BOLD				
CC1	Ground and grilled <i>Pangasius bocourti</i>	<i>P. hypophthalmus</i>	99	No	×	<i>P. hypophthalmus</i>	99	×
CC2	Ground and grilled <i>Pangasius bocourti</i>	<i>P. hypophthalmus</i>	99	No	×	<i>P. hypophthalmus</i>	99	×
CQ1	Ground and roasted <i>Pangasius hypophthalmus</i> with cinnamon	<i>P. hypophthalmus</i>	95	No	×	<i>P. hypophthalmus</i>	99	×
CQ2	Ground and roasted <i>Pangasius hypophthalmus</i> with cinnamon	<i>P. hypophthalmus</i>	99	100	×	<i>P. hypophthalmus</i>	99	×
KC1	Dry fillet of <i>Pangasius hypophthalmus</i>	<i>P. hypophthalmus</i>	95	No	✓	<i>P. hypophthalmus</i>	99	✓
KC2	Dry fillet of <i>Pangasius hypophthalmus</i>	<i>P. hypophthalmus</i>	99	100	✓	<i>P. hypophthalmus</i>	99	✓
CCV1	Ball of ground and roasted <i>Pangasius bocourti</i>	<i>P. hypophthalmus</i>	99	100	×	<i>P. hypophthalmus</i>	99	×
CCV2	Ball of ground and roasted <i>Pangasius bocourti</i>	<i>P. hypophthalmus</i>	99	100	×	<i>P. hypophthalmus</i>	99	×
TL1	Ball of ground and Roasted <i>Pangasius bocourti</i> with dill	<i>P. hypophthalmus</i>	97	No	×	<i>P. hypophthalmus</i>	99	×
TL2	Ball of ground and Roasted <i>Pangasius bocourti</i> with dill	<i>P. hypophthalmus</i>	99	100	×	<i>P. hypophthalmus</i>	99	×
CV1	Ball of ground <i>Pangasius bocourti</i>	<i>P. hypophthalmus</i>	99	100	×	<i>P. hypophthalmus</i>	99	×



CV2	Ball of ground <i>Pangasius bocourti</i>	<i>P. hypophthalmus</i>	97	No	×	<i>P. hypophthalmus</i>	99	×
LB1	Fillet of <i>Pangasius hypophthalmus</i> rolled with flour	<i>P. hypophthalmus</i>	100	100	✓	<i>P. hypophthalmus</i>	99	✓
LB2	Fillet of <i>Pangasius hypophthalmus</i> rolled with flour	<i>P. hypophthalmus</i>	100	100	✓	<i>P. hypophthalmus</i>	99	✓
FB1	Burger of <i>Pangasius bocourti</i>	<i>P. hypophthalmus</i>	100	100	×	<i>P. hypophthalmus</i>	99	×
FB2	Burger of <i>Pangasius bocourti</i>	<i>P. hypophthalmus</i>	100	100	×	<i>P. hypophthalmus</i>	99	×
CH1	Roll of roasted <i>Oncorhynchus mykiss</i>	<i>Oncorhynchus mykiss</i>	99	100	✓	<i>Oncorhynchus mykiss</i>	99	✓
CH2	Roll of roasted <i>Oncorhynchus mykiss</i>	<i>Oncorhynchus mykiss</i>	99	100	✓	<i>Oncorhynchus mykiss</i>	99	✓
CM1	Fillet of <i>Prionace glauca</i>	<i>Prionace glauca</i>	99	100	✓	<i>Prionace glauca</i>	99	✓
CM2	Fillet of <i>Prionace glauca</i>	<i>Prionace glauca</i>	99	100	✓	<i>Prionace glauca</i>	99	✓

“✓” shows the same species between analysis results and products.

“×” represents the difference in species between analysis results and products.

“No” means no species name found in BOLD.




Figure 1. PCR products using Fish primer (left) and MAB primer (right)

* Number 1 to 10: PCR products of samples CCV1, CCV2, TL1, TL2, CQ1, CQ2, KC1, KC2, CC1, CC2, respectively.

Number 11 to 20: products of samples LB1, LB2, VB1, VB2, FB1, FB2, CH1, CH2, CM1 and CM2, respectively.

M: Marker, 100bp




Figure 2. PCR products using Cytb primers

Number 1 → 10: PCR products of samples CCV1, CCV2, TL1, TL2, CQ1, CQ2, KC1, KC2, CC1 and CC2, respectively.

Number 11 → 20: PCR products of samples LB1, LB2, VB1, VB2, FB1, FB2, CH1, CH2, CM1, CM2, respectively.

M: Marker, 100bp.

Accepted