



Northernmost Record of *Triaenodon obesus* (Rüppell, 1837) in the Eastern Tropical Pacific

Edgar Eduardo Becerril-García¹, Tania Cecilia Cota-Lucero², Felipe Galván-Magaña^{1,*}

¹ National Polytechnic Institute, Interdisciplinary Center of Marine Sciences, 23096 La Paz, Baja California Sur, Mexico.

² Autonomous University of Baja California Sur, Program of Marine Botany, 23080, La Paz, Baja California Sur, Mexico.

* Corresponding Author: Tel.: +52.612 1226001 ;
E-mail: galvan.felipe@gmail.com

Received 08 May 2017
Accepted 28 September 2017

Abstract

This is the northernmost record of a whitetip reef shark, *Triaenodon obesus*, in the Eastern Tropical Pacific. The shark was observed in Magdalena Bay, Mexico, at a depth of 4 m and had a total length of 1.2–1.5 m. The habitat where the record was made is consistent with the conditions for the distribution of species. The presence of *T. obesus* in the area could be related to the protection of near natural areas and the seasonal bans on shark and ray fishing implemented in Mexico since 2012.

Keywords: Elasmobranchs, Carcharhinidae, Mexico, Eastern Pacific, new record.

Introduction

The whitetip reef shark, *Triaenodon obesus* (Rüppell, 1837), is a small shark species with an extremely short and rounded snout, oval eyes, and bright white tips on the first dorsal fin and upper caudal fin lobe (Fischer, Krupp, Schneider, Sommer, Carpenter, & Niem, 1995). This mostly nocturnal predator feeds on benthic organisms like fishes, octopuses, lobsters, and crabs in coral and rocky reefs, and usually inhabits a depth range of 8–40 m (Randall, 1977). *T. obesus* is aplacental viviparous, giving birth to litters of 1–5 pups (Compagno, Dando, & Fowler, 2005). It is categorized as “Near Threatened” by the IUCN due to its biological characteristics, restricted habitat type, and distribution in coastal zones and islands, which makes it vulnerable to overexploitation (Smale, 2005).

In the Eastern Tropical Pacific, the whitetip reef shark is distributed along the coast of Panama, Costa Rica, Colombia, and Ecuador (Fischer et al., 1995). In Mexico, the records of this species are limited to tropical areas like the Revillagigedo Archipelago (Ochoa-López, Villavicencio-Garayzar, & Ruiz-Gaytan, 1997), the coast of Jalisco (Corgos & Rosende-Pereiro, 2016), Nayarit (Pérez-Jiménez et al., 2005; Erisman et al., 2011), and the mouth of the Gulf of California (GBIF, 2017; OBIS, 2017; Robertson & Allen, 2017), with no consistent records of its presence far north, or in temperate waters of this

country.

Materials and Methods

The record was made in July 2015 while monitoring marine phanerogams on a rocky reef off the town of El Barrote (24°40'17.9050"N, 112°09'32.6640"W), within the Magdalena Bay–Almejas lagoon complex in the western coast of the Baja California Peninsula (Figure 1). This bay is subjected to upwelling effects that lead to high primary productivity (Zaytsev, Cervantes-Duarte, Montante, & Gallegos-García, 2003). The input of nutrients by mangroves and sea grasses translates into a high availability of food within the bay, which is recognized as an area of high ecological importance (Acosta-Velázquez & Ruíz-Luna, 2007). During the monitoring, the sea surface temperature and salinity were measured with a YSI CastAway™ CTD, while the photographic record was performed with a Canon Powershot G15 camera. The taxonomic identity for the shark was determined by the description of Fischer et al. (1995).

Results and Discussion

This study constitutes the northernmost record of *T. obesus* in the Eastern Tropical Pacific. The sea surface temperature and salinity showed respective values of 23 °C and 34 psu. The presence of the shark

was recorded at approximately 50 m off the coast at a depth of 4 m. Its total length was estimated to be 1.2–1.5 m and it was identified as *T. obesus* by noting the presence of two white tips on the first dorsal fin and caudal fin, oval eyes, as well as the elongated and slender shape typical of this species (Figure 2). The sex of the specimen, however, could not be determined.

The average annual sea surface temperature in Magdalena Bay-Almejas Bay ranged from 18–24°C (Lluch-Belda, Hernández-Rivas, Saldierna-Martínez, & Guerrero-Caballero, 2000), so the presence of rocky reefs, the availability of prey as well as the environmental conditions observed when recording the shark are consistent with the habitat previously

described for *T. obesus* (Randall, 1977; Fischer et al., 1995). Although El Niño-Southern Oscillation could have an indirect effect on the presence of the organism in the bay, related to an increase of water temperature during July 2015 (Cervantes-Duarte & García-Romero, 2016), the record itself and the related conclusions to El Niño effect are beyond the scope of this study.

Sightings of some species like the bull shark *Carcharhinus leucas*, lemon shark *Negaprion brevirostris*, tiger shark *Galeocerdo cuvier*, and whitetip reef shark, have increased at some diving sites of the southern Baja California Peninsula coast over the last years (Reyes-Bonilla et al., 2016; OBIS, 2017). These sightings may be related to protected




Figure 1. Previous sightings and new record of *Trienodon obesus* in Baja California Sur, Mexico.



Figure 2. Whitetip reef shark in a rocky reef at El Barrote, BCS, Mexico.

areas like Cabo Pulmo National Park and Cabo San Lucas Flora and Fauna Protection Area, as well as the seasonal bans on shark and ray fishing implemented in the Mexican Pacific since 2012 (Martínez-Cruz & Oviedo-Pérez, 2014; Reyes-Bonilla *et al.*, 2016).

Acknowledgments

This work is dedicated to the memory of Dr. Rafael Ríosmena-Rodríguez. Our thanks also to the people who assisted in sampling, especially Karla Pedraza and Atzcalli Ehecatl Hernández-Cisneros. We thank Pelagios Kakunjá, A.C. and Consejo Nacional de Ciencia y Tecnología (CONACyT) for the scholarships provided and the Instituto Politécnico Nacional for funding through grants from Comisión de Operación y Fomento de Actividades Académicas and the Estímulo al Desempeño de los Investigadores (EDI).

References

- Acosta-Velázquez, J., & Ruiz-Luna, A. (2007). Variación en la cobertura, distribución y estructura de los manglares del complejo lagunar Bahía Magdalena-Bahía Almejas (1990-2005). In R. Funes-Rodríguez, J. Gómez-Gutiérrez & R. Palomares-García (Eds.), *Estudios ecológicos en Bahía Magdalena* (pp. 127-141). La Paz, México, Instituto Politécnico Nacional., 311 pp.
- Cervantes-Duarte, R. & García-Romero, F. de J. (2016). Hydrographic characteristics off the coast of Bahía Magdalena, BCS, Mexico during El Niño 2015. *CICIMAR Oceanides*, 31(2), 9-19.
- Compagno, L.J.V., Dando, M., & Fowler, S.L. (2005). *Sharks of the World*. New Jersey, USA, Princeton University Press., 496 pp.
- Corgos, A., & Rosende-Pereiro, A. (2016). First record of the whitetip reef shark, *Triaenodon obesus* from the coast of Jalisco, western Mexico mainland. *Marine Biodiversity Records*, 9(1), 66. <https://doi.org/10.1186/s41200-016-0069-7>
- Erisman, B.E., Galland, G.R., Mascarenas, I., Moxley, J., Walker, H., Aburto-Oropeza, O., ... Ezcurra, E. (2011). List of coastal fishes of Islas Marias archipelago, Mexico, with comments on taxonomic composition, biogeography, and abundance. *Zootaxa*, 2985, 26-40.
- Fischer, W., Krupp, F., Schneider, W., Sommer, C., Carpenter, K.E., & Niem, V.H. (1995). Guía FAO para la identificación de especies para los fines de la pesca. Pacífico centro - oriental. Volumen II. Vertebrados-Parte 1. Rome, Italy, FAO., 563 pp.
- GBIF. (2017). Global Diversity Information Facility. Retrieved from <http://www.gbif.org/>
- Lluch-Belda, D., Hernández-Rivas, M.E., Saldierna-Martínez, R., & Guerrero-Caballero, R. (2000). Sea surface temperature variability at Magdalena Bay, Baja California Sur, México. *CICIMAR Oceanides*, 15(1), 1-23.
- Martínez-Cruz, L., & Oviedo-Pérez, J.L. (2014). Temporada de veda para la captura de tiburones en el Golfo de California y Mar Caribe. Ciudad del Carmen, México, Instituto Nacional de Pesca., 25 pp.
- Ochoa-López, E., Villavicencio-Garayzar, C., & Ruiz-Gaytan, A. (1997). First record of the whitetip reef shark, *Triaenodon obesus* in the west coast of México (Revillagigedo Archipelago). *Revista de Biología Tropical*, 45(3), 687-688.
- OBIS. (2017). Global biodiversity indices from the Ocean Biogeographic Information System. Intergovernmental Oceanographic Commission of UNESCO. Retrieved from <http://www.iobis.org>
- Pérez-Jiménez, J.C., Sosa-Nishizaki, O., Furlong-Estrada, E., Corro-Espinosa, D., Venegas-Herrera, A., & Barragán-Cuencas, O.V. (2005). Artisanal shark fishery at "Tres Marias" islands and Isabel Island in the central Mexican Pacific. *Journal of Northwest Atlantic Fishery Science*, 35, 333-343.
- Randall, J.E. (1977). Contribution to the biology of the whitetip reef shark (*Triaenodon obesus*). *Pacific Science*, 31(2), 143-64.
- Reyes-Bonilla, H., Ayala-Bocos, A., Fernández-Rivera Melo, F.J., Zepeta-Vilchis, R., Asúnsulo-Rivera, A., & Ketchum, J.A. (2016). A bibliographic and field record chronology of sharks at Cabo Pulmo National Park, Gulf of California. *CICIMAR Oceanides*, 31(1), 55-57.
- Robertson, D., & Allen, G. (2017). Shorefishes of the Tropical Eastern Pacific: online information system. Retrieved from <http://biogeodb.stri.si.edu/sftep/es/pages>
- Smale, M.J. (2005). *Triaenodon obesus*. The IUCN Red List of Threatened Species 2005: e.T39384A10188990. Retrieved from <http://www.iucnredlist.org>
- Zaytsev, O., Cervantes-Duarte, R., Montante, O., & Gallegos-García, A. (2003). Coastal upwelling activity on the Pacific shelf of the Baja California Peninsula. *Journal of Oceanography*, 59(4), 489-502. <https://doi.org/10.1023/A:1025544700632>