



Length-Weight Relationships of Five Elasmobranch Species from the Pacific Coast of Mexico

José De La Cruz-Agüero*, Francisco Javier García-Rodríguez, Víctor Manuel Cota-Gómez

Instituto Politécnico Nacional, Centro Interdisciplinario de Ciencias Marinas, Colección Ictiológica. Avenida IPN s/n, Colonia Playa Palo de Santa Rita, La Paz, Baja California Sur 23096, México.

Tel: 52 612 122 5344-81600; Fax: 52 612 122 5344

E-mail: jacruz@ipn.mx

Abstract

Length-weight relationships (LWR) were estimated for five elasmobranch species found in the Eastern Pacific: *Heterodontus francisci* (Girard, 1855), *Urobatis halleri* (Copper, 1863), *Urobatis maculatus* (Garman, 1913), *Diplobatis ommata* (Jordan and Gilbert, 1890), and *Rhinobatos productus* (Ayres, 1854). Species were selected because none had previously published data on LWR. The specimens were sampled along the Pacific west coast of Mexico, including the Gulf of California, between June 2009 and May 2013, using different fishing gear. The values of the exponent “b” of the LWR ranged from 2.52 to 3.06. This study provides the first reference on the LWR for these elasmobranch species.

Introduction

The analyses that involve length-weight relationships (LWR) of fish species have increased because such data are useful, for example, for the conversion of growth-in-length equations to grow-in-weight. The parameters of LWR also have been applied to determining stock structure as well as for estimates of the fish condition and other fisheries applications (Mendes, Fonseca, & Campos, 2004). Nevertheless, available information on LWR of sharks and rays are not often reported or is very scarce (e.g. de Loyola-Fernández et al., 2017; Teixeira, Silva, Fabr e, & Batista, 2017; Ismen, Yigin, Altinagac, & Ayaz, 2009; Yigin & Ismen, 2009; Yeldan & Avsar, 2007). In the west coast of Mexico, elasmobranchs are commonly fished by the artisanal fisheries as well as bycatch by shrimp and gillnet fisheries (Ehemann et al., 2017). However, shark and ray populations have fallen drastically mainly due to its overexploitation. According to recent assessments, 36 species of elasmobranchs inhabiting Mexican waters are listed with some conservation status or control regime for their international trade (i.e. threatened, near threatened, critically endangered, and vulnerable) (Del Moral-Flores, Morrone, Alcocer-Durand, Espinosa-P erez, & Ponce de Le on, 2015). An initial step to mitigate adverse impacts on natural resources and to support the development of proactive resolution responses is the generation and diffusion of primary biological data. To our knowledge, this study provides the first reference of LWR for five elasmobranch species that inhabit the Eastern Pacific, where they are endemic. Therefore, the primary objective of this study was to provide the specific information to compensate for this lack of knowledge.

Material and Methods

Samples were collected along the northwest coast of Mexico (22°50'28.7"N - 110°12'12.8"W and 27°44' 40.6"N - 114°10' 7.1"W), including the Gulf of California (24°7'48"N - 109°53' 7.9"W and 26°58' 26.2"N - 111°56'54.7"W) within a multi-year collection program (June 2009- May 2013) for the study and evaluation of marine resources. The fish catches were made using a commercial bottom trawl net and long-line, and some were taken in local fishing grounds. Of all the species sampled, only those for which there was no published data for the LWR were selected for this study. In the laboratory, the fish were identified based on specific keys and all scientific names, authors, years and family assignments were checked against FishBase (Froese & Pauly, 2017). Each fish was measured to the nearest 0.1 mm (total length) and weighed with a digital balance, to the nearest 1 g (total weight). All specimen were firstly maintained in 10% buffered formalin solution and then preserved in 95% ethyl alcohol for subsequent deposit in the fish collection (CI) of the Centro Interdisciplinario de Ciencias Marinas in La Paz, Baja California Sur, México. The length and weight data were log-transformed and plotted for visual inspection of outliers; extreme outliers were omitted from the analyses (Froese, Tsikliras, & Stergiou, 2011). Parameters of the LWR were calculated for males and females combined, according to the formula: $\log W = \log "a" + "b" \log TL$, where (W) is the total weight (g), (TL) is the total length (cm), " a " is the intercept, and " b " is the slope of the LWR (which indicates isometric growth in body proportions if $b \sim 3$) (Ricker, 1973; Froese, 2006). If a fish grows without changing its shape or its density, then the fish is said to exhibit isometric growth. In this case, the volume of the fish is proportional to any linear measure of its size. If a fish changes shape or density as it grows, then " b " is significantly different from 3, and the fish is said to exhibit allometric growth (Froese, 2006). The 95% confidence interval (CI) for parameters " a " and " b " was calculated according to Cohen, Cohen, West, and Aiken (2003). The coefficient of determination (R^2) was used to evaluate the correlation between W and L . A test (Student's t -test; $H_0: b = 3; P < 0.05$) of whether the elasmobranchs studied exhibit isometric growth or not was applied, using the FishR Vignette by D. Ogle (<http://derekogle.com/fishR/>) implemented in the statistical software R. 3.3.1 (R Development Core Team, 2016). Ray individuals with broken or incomplete tails were not considered to avoid unreasonable parameter values. The LWR was analysed by software XLSTAT Pro© version 2010.

Results

In this LWR study, 244 elasmobranch specimens belonging to five species and four families were examined (other 23 were discarded as outliers). All samples sizes (N), minimum (Min) and maximum (Max) length and weight, parameters of LWR (" a " and " b "), 95% confidence intervals (CI) for these parameters, the coefficient of determination (R^2) by species, and the growth type, are presented in Table 1. All regressions were highly significant ($P < 0.01$), with the coefficient of determination equal or greater than 0.92 for all species studied. The estimated b values oscillated from 2.52 for the Shovelnose Guitarfish *R. productus* and 3.06 for the Horn Shark *H. francisci*. The species *D. ommata* and *R. productus* showed allometric growth, while the rest of species showed isometric growth.

Discussion

The five studied species had no previous LWR data in FishBase (Froese & Pauly, 2017) and therefore our results contribute to the knowledge of these elasmobranch species. Present values of " b " range between 2.52 and 3.06, falling



within the expected range of 2.5–3.5, and all calculated values for “a”, were within the range of 0.001 and 0.05, which validated these length-weight relationships estimates, as proposed by Froese (2006). However, our data are representative of a particular size or growth stage, and it is known that the LWR in fishes is affected by several factors (Froese, 2006) such as (1) environmental (seasonal variation, habitat type, geographic region), (2) biological (population, gonad maturity, sex, growth phase, diet, degree of stomach fullness, health and general fish condition), and (3) artifactual (preservation techniques, number of specimens examined, size range covered and type of length used). Thus, estimated parameters in the present work should be treated with caution or limited to similar fishery conditions and size ranges even though current parameters overlapped with the Bayesian confidence limits calculated interactively in FishBase (Froese, Thorson, & Reyes, 2014). It is recommended increasing information within the entire size range of each species studied. Finally, although there are no published records regarding LWRs of these species that would be comparable to the present results, they can provide a baseline for future studies concerning these elasmobranchs, some of which can be threatened by overfishing.

Acknowledgements

The authors would want to thank the support to the Ichthyology Collection (CI) at CICIMAR-IPN (<http://coleccion.cicimar.ipn.mx>) and SIP-IPN, CONABIO, and CONACyT. JDA and FJGR also appreciate the scholarly grants from SNI-CONACyT, COFAA-IPN, and EDI-IPN. Anonymous reviewers improved the manuscript with their comments and suggestions.

References

- Cohen, J., Cohen, P., West, S.G., & Aiken, L.S. (2003). Applied Multiple Regression/Correlation Analysis for the Behavioural Sciences (3rd edition). Mahwah, New Jersey, Lawrence Earlbaum Associates Publ., 1200 pp
- de Loyola- Fernández, I., Báez, J.C., García-Barcelona, S., Camiñas, J.A., Ortíz de Urbina, J.M., & Macías, D. (2017). Length-weight relationships of Kitefin Shark *Dalatias licha*, and Little Sleeper Shark *Somniosus rostratus* from the Western Mediterranean Sea, and Long Snouted Lancetfish *Alepisaurus ferox* from the Eastern North Atlantic Ocean. *Turkish Journal of Fisheries and Aquatic Sciences*, 17, 1073–1076. http://dx.doi.org/10.4194/1303-2712-v17_5_24
- Del Moral-Flores, L.F., Morrone, J.J., Alcocer Durand, J., Espinosa-Pérez, H., & Ponce de León, G.P. (2015). Lista patrón de los tiburones, rayas y quimeras (Chondrichthyes, Elasmobranchii, Holocephalii) de México. *Arxius de Miscellània Zoològica* 13: 47–163. <https://dx.doi.org/10.15470/hrllkv>
- Ehemann, N.R., Pérez-Palafox, X.A., Mora-Zamacona, P., Burgos-Vázquez, M. I., Navia, A. F., Mejía-Falla, P. A., & Cruz-Escalona, V. H. (2017). Size-weight relationships of batoids captured by artisanal fishery in the southern Gulf of California, Mexico. *Journal of Applied Ichthyology*, 33, 1051–1054. <https://doi.org/10.1111/jai.13421>
- Froese, R. (2006). Cube law, condition factor, and weight-length relationships: History, meta-analysis, and recommendations. *Journal of Applied Ichthyology*, 22, 241–253. <http://dx.doi.org/10.1111/j.1439-0426.2006.00805.x>
- Froese, R., Tsikliras, A.C., & Stergiou, K.I. (2011). Editorial note on weight-length relations of fishes. *Acta Ichthyologica Piscatoria*, 41, 261–263. <https://dx.doi.org/10.3750/AIP2011.41.4.01>
- Froese, R., Thorson, J.T., & Reyes, R.B. (2014). A Bayesian approach for estimating length-weight relationships in fishes. *Journal of Applied Ichthyology*, 30, 78–85. <https://dx.doi.org/10.1111/jai.12299>
- Froese, R., & Pauly, D. (2017). *FishBase*. World Wide Web Electronic Publication. Retrieved from <http://www.fishbase.org>
- Ismen, A., Yigin, C.C., Altinagac, U., & Ayaz, A. (2009). Length-weight relationships for ten shark species from Saros Bay (North Aegean Sea). *Journal of Applied Ichthyology*, 25, 109–112. <https://dx.doi.org/10.1111/j.1439-0426.2009.01263.x>
- Mendes, B., Fonseca, P., & Campos A. (2004). Weight-length relationships for 46 fish species of the Portuguese west coast. *Journal of Applied Ichthyology*, 20, 355–361. <https://dx.doi.org/10.1111/j.1439-0426.2004.00559.x>
- R Development Core Team (2016). R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. Version 3.3.1. Retrieved from <http://www.r-project.org/>.
- Ricker, W. E. (1973). Linear regressions in fishery research. *Journal of the Fisheries Research Board of Canada*, 30, 409–434. <https://doi.org/10.1139/f73-072>



- Teixeira, E.C., Silva, V.E.L., Fabr , N.N., & Batista, V.S. (2017). Length-weight relationships for four stingray species from the tropical Atlantic Ocean. *Journal of Applied Ichthyology*, 33, 594–596. <https://doi.org/10.1111/jai.13255>
- Yeldan, H., & Avsar, D. (2007). The length-weight relationship for five elasmobranch species from the Cilician Basin shelf waters (Northeastern Mediterranean). *Journal of Applied Ichthyology*, 23: 713–714. <https://dx.doi.org/10.1111/j.1439-0426.2007.00858.x>
- Yigin, C.C., & Ismen, A. (2009). Length-weight relationships for seven rays from Saros Bay (North Aegean Sea). *Journal of Applied Ichthyology*, 25, 106–108. <https://dx.doi.org/10.1111/j.1439-0426.2008.01161.x>

Accepted Manuscript



Table 1. Length-weight relationships for 5 elasmobranch species caught on the coast of Mexico. LWR parameters are shown with 95% confidence interval (CI). All regressions were significant to ($P < 0.01$).

TL, total length; N, sample size; Min-Max, minimum and maximum respectively (mm, millimeters; g, grams); “ a ” and “ b ”, parameters of the LWR, R^2 , the coefficient of determination.

Family/Species	N	Parameters of the			LWR		Length range TL (mm)			Weight range (g)			Growth Type /	P value
		a	95% CI of a	b	95% CI of b	R^2	Min	Max	Mean	Min	Max	Max		
Heterodontidae														
<i>Heterodontus francisci</i>	22	0.006	0.003-0.012	3.06	2.83-3.29	0.98	135	476	306	18.2	777.9	282.8	Isometric	0.7389
Narcinidae														
<i>Diplobatis ommata</i>	50	0.021	0.009-0.033	2.82	2.62-3.00	0.98	43	218	122	1.1	125.5	5.01	Allometric	0.0038
Rhinobatidae														
<i>Rhinobatos productus</i>	34	0.080	0.027-0.113	2.52	2.34-2.71	0.97	92	351	167	12.3	534.7	108.7	Allometric	7.3e-05
Urotrygonidae														
<i>Urobatis halleri</i>	62	0.065	0.039-0.091	2.99	2.85-3.13	0.99	57	192	114	12.5	470.6	143.6	Isometric	0.8081
<i>Urobatis maculatus</i>	76	0.058	0.008-0.108	3.04	2.71-3.35	0.92	42	174	111	7.1	390	117.8	Isometric	0.5498

Student's t-test for growth type; $H_0: b = 3$; $P < 0.05$