



Distribution of Ascidians with a New Record of the Non-Indigenous Species *Polyclinum constellatum* Savigny, 1816 from the Aegean Coast of Turkey

Sinem Aydin-Onen^{1,*}

¹ Dokuz Eylül University, Institute of Marine Sciences and Technology, Department of Marine Living Resources, 35340 Inciraltı, Izmir, Turkey.

* Corresponding Author: Tel.: +90.507 5936864 ;
E-mail: sinem.onen@deu.edu.tr

Received 14 August 2017
Accepted 20 November 2017

Abstract

The current study attempts to investigate the distribution of ascidian species collected from eight stations located along the eastern coasts of Aegean Sea, Turkey. For this purpose, field studies were conducted in July 2016. This study revealed the occurrence of ascidian species belonging to 7 families and 12 genera. The invasive species *Polyclinum constellatum* Savigny, 1816 is reported for the first time from Turkish waters and for the second time in the Mediterranean Sea. Furthermore, *Symplesma brakenhielmi* (Michaelsen, 1904) is found as a new record for the Turkish Aegean Sea fauna. Knowledge about the ascidian diversity of the Turkish Seas is still incomplete and large-scale surveys, additional taxonomic and molecular studies are needed.

Keywords: Tunicates, non-indigenous ascidians, invasive species, Aegean Sea, *Polyclinum constellatum*.

Introduction

In marine ecosystems, ascidians (Phylum: Chordata, Subphylum: Tunicata, Class: Ascidiacea) are commonly known as sea squirts and they comprise an important group of marine fouling organisms around the world (Pennai & Rothbacher, 2016). Moreover, a number of ascidians are successful invaders in marine ecosystems (Shenkar & Swalla, 2011; Zhan, Briski, Bock, Ghabooli, & MacIsaac, 2015).

On a global scale, great attention has been given to ascidians due to a) their nutritive value (Lambert, Karney, Rhee, & Carman, 2016), b) their ability to produce bioactive compounds, specifically substances which serve as cytotoxic, antibacterial, antipyretic, analgesic, and histamine (Gopalakrishnan, Meenakshi, & Shanmuga, 2011), and compounds against various solid-type tumors (Gab-Alla, 2008), c) and also their use in studies of evolutionary developmental biology (Lemaire, 2011). Furthermore, invasive ascidians have ability to a) change community structure (Tracy & Reynolds, 2014), b) they can lead to the loss of native genotypes (Katsanevakis *et al.*, 2014), c) they can modify habitats, d) they can affect ecosystem processes and trophic relationships in the food chain, e) they can economically affect fishing and aquaculture activities and also f) they can

introduce new diseases and pathogens or toxic substances (Marques *et al.*, 2013).

According to Shenkar *et al.* (2017), about 3,000 ascidian species have been described worldwide. In addition, knowledge on ascidian diversity across the Turkish coasts is relatively limited. Çınar (2014) published a comprehensive regional tunicate species checklist. According to the latter review the faunistic studies on the ascidians, thaliaceans, and appendicularians (or larvaceans) in the Turkish Seas began with Colombo (1885), who reported six species from the entrance of the Çanakkale Strait to the Aegean Sea. After then, Ostroumoff (1894) conducted a marine survey in the Sea of Marmara and reported two ascidian species. Moreover, Ostroumoff (1896) investigated plankton and benthos during the Selanik Expedition to the Sea of Marmara after the invitation of the Ottoman Sultan Abdulhamit II and he determined 17 tunicate species belonging to class Ascidiacea and one species belonging to class Thaliacea. Afterwards, ascidians have been documented in different studies from the Turkish coasts (Demir, 1952; Demir, 1954a, 1954b; Nalbantoğlu, 1955; Băcescu, 1961; Kiseleva, 1961; Ergen, 1967; Kiseleva, 1969; Geldiay & Kocataş, 1972; Gökalp, 1972; Pınar, 1974; Uysal, 1976; Kocataş, 1978; Mavili, 1987; Ergen, Kocataş, Katağan, & Cınar, 1994; Koçak, Ergen, & Çınar,

1999; Koçak & Küçüksezgin, 2000; Tarkan, 2000; Dinçaslan & Öber, 2004; Öztürk *et al.*, 2004; Aslan, 2006; Çınar, Bilecenoglu, Öztürk, & Can, 2006a; Çınar *et al.*, 2006b; Lakkis & Toklu, 2007; Okuş *et al.*, 2007; Çınar *et al.*, 2008; Mutlu & Ergev, 2008; Çınar *et al.*, 2011; Aslan-Cihangir, Izquierdo-Muñoz, Pancucci-Papadopoulou, & Can-Yılmaz, 2011; Gözcelioğlu, 2011; Ramos-Espla, Izquierdo, & Çınar, 2013; Artüz, Sönmez, & Kubanç, 2014).

In total 61 species belonging to the three tunicate classes Ascidiacea (50 species), Thaliacea (3 species), and Appendicularia (8 species) were reported from the Turkish coasts by Çınar (2014). Four of these species, namely *Symplesma brakenhielmi* (Michaelsen, 1904), *Microcosmus exasperatus* Heller, 1878, *Herdmania momus* (Savigny, 1816), and *Phallusia nigra* Savigny, 1816 are non-indigenous (NIS) (Çınar, Bilecenoglu, Öztürk, & Can, 2006a; Ramos-Espla, Izquierdo, & Çınar, 2013). According to this annotated checklist, the Aegean records include 29 species of Ascidiacea, 3 species of Thaliacea, and 7 species of Appendicularia. Two NIS, namely *M. exasperatus* and *P. nigra* were reported from this region (Çınar, 2014). After this publication, one more NIS, *Styela clava* Herdman, 1881 was recorded in one locality (Karamürsel, İzmit Bay) in the Sea of Marmara on rocks at 10 m depth (Çınar, 2016). Up to now, the addition of this new alien species brings the total number of identified tunicate species in Turkish coasts to 62.

The aim of the present work was to study

ascidian diversity and to detect the occurrence of alien and invasive ascidian species through field surveys in selected stations situated in the eastern Aegean Sea coasts.

Materials and Methods

Study Area

The study area is located along the coastline of the eastern Aegean Sea, Turkey (Figure 1). In this study, eight sampling stations were selected from potential places that are susceptible to introductions or the dispersal of ascidian species. Sampling was carried out in areas which have high numbers of local fishing boats, commercial fishing vessels and recreational boats (Figure 1). Station S1 is located in Aliğa Bay and Station S2 is placed in Yeni Foça located on the Aegean Coast. These stations host seasonal fishing and recreational vessels. Station S3 is situated in Güzelbahçe in İzmir Bay and it is subject to intense anthropogenic activities and urban stress. Station S4 is found in Özbek Bay (Urla, İzmir) and Station S5 is located on the coast of Özdere (İzmir). The number of fishing and recreational boats is high at these two stations, especially during summer months. Station S6 is placed in Kuşadası Bay (Aydın), which is a highly developed area with maritime traffic by fishing and recreational boats, including cruise vessels. Stations 7 and 8 are placed in Taşburun (Didim, Aydın) located on the Aegean




Figure 1. Map of the study area showing locations of sampling stations. S1: Aliğa, S2: Yeni Foça, S3: Güzelbahçe, S4: Özbek, S5: Özdere, S6: Kuşadası, S7: Taşburun, and S8: Akbük.

Coast and Akbük Bay (Didim, Aydın), respectively. These stations are used as sheltering places for commercial fishing vessels, recreational fishing boats, and yachts (Figure 1).

Station numbers sites, geographical coordinates, and sampling dates for each study site are given in Table 1.

Methods

Ascidians were sampled in July 2016. At each station, ascidians were collected by examining associated subtidal structures such as ropes, wheels, pontoons, buoys, tires, chains, walls, plastic structures and other floating materials. Specimens were taken by hand from artificial and natural substrates using hand tools to scrape the ascidians from surfaces.

Specimens were photographed in situ and were collected from a depth zone of 0-1 m. In addition to photographs, observation notes (i.e. color, shape, substrate type, and depth) were recorded, and then the samples were narcotized with menthol crystals for relaxation and preserved in fixatives (mixture of 850 ml of seawater, 50 ml of distilled water, 100 ml of full-strength formaldehyde, and 1 gram of sodium borate) for identification. Moreover, few ascidians were preserved in ethanol 95% for future molecular studies.

Ascidians were separated, identified and solitary species were counted. Specimens were examined and then identified under an Olympus-SZX16 stereomicroscope. Dissected ascidians were stained with haemalum. Images were taken by using a digital camera. Authoritative and valid taxonomic keys and publications (Van Name 1921, 1931, 1945; Kott 1985, 1990, 1992, 2001; Monniot, Monniot, & Laboute, 1991; Monniot & Monniot, 2001; Rocha, Zanata, & Moren, 2012) were used for species identification.

Three replicate surface water samples were collected in each station with sampling bottles (precleaned and rinsed twice with seawater) and then physicochemical variables such as temperature, salinity, pH, and dissolved oxygen (DO) were measured by using a portable instrument

(HACH-HQ40D, Hach Lange, Düsseldorf, Germany).

Statistical Analysis

Community variables such as number of species, number of individuals were calculated for each sampling station. Frequency Index (F) was calculated according to Soyer (1970) in order to determine the frequency value [(constant (F≥49%), common (49%>F≥25%) and rare (F<25%)] of each ascidian species. The frequency index formula is the following: $f=m/M*100$, where m= the number of stations where the species was detected and M= the numbers of all stations. Data analysis was performed using STATISTICA v.7.1 (STATSOFT) software package. Spearman's rank order correlation test was used to examine relationships among the physicochemical variables, number of individuals, and the number of species.

Results

Taxonomic Composition

A total of 396 solitary ascidians were examined corresponding to 7 species which belonged to 5 families namely Ascidiidae (2), Cionidae (1), Molgulidae (1), Pyuridae (1), and Styelidae (2) (Table 2). However, three taxa could be determined only to genus level (*Didemnum* sp., *Aplidium* sp., and *Microcosmus* sp.) (Table 2). The species *Didemnum* sp. and *M. exasperatus* were found in only one station, while *Styela plicata* (Lesueur, 1823) and *Styela canopus* (Savigny, 1816) were observed at all sampling stations (Figure 2).

Three taxa, namely *Microcosmus* sp., *Botrylloides leachii* (Savigny, 1816), and *S. canopus* were found on both natural and artificial substrata (Table 2). The orders Aplousobranchia and Phlebobranchia were represented with three colonial forms and three solitary species, respectively. Furthermore, the order of Stolidobranchia had the greatest number of species (8), three of which were colonial and others were solitary.

Table 1. Stations numbers, sampling sites, geographical coordinates, and sampling dates of each site surveyed in Aegean coasts of Turkey

Station Numbers	Sampling Sites	Geographical Coordinates		Sampling Dates
		Latitude (N)	Longitude (E)	
S1	Aliağa	38°48'23.63"	26°58'28.86"	02.07.2016
S2	Yeni Foça	38°44'30.24"	26°50'19.47"	02.07.2016
S3	Güzelbahçe	38°22'39.56"	26°53'02.06"	02.07.2016
S4	Özbek	38°22'28.66"	26°40'56.10"	02.07.2016
S5	Özdere	38° 0'37.55"	27° 6'18.02"	03.07.2016
S6	Kuşadası	37°51'41.15"	27°15'20.10"	04.07.2016
S7	Taşburun	37°27'11.44"	27°13'07.87"	06.07.2016
S8	Akbük	37°23'11.70"	27°25'54.48"	06.07.2016

Table 2. Number of ascidian species and abundance of solitary ascidians recorded in the sampling stations. Substrate and substrate type, A: Artificial, N: Natural, Colonial C

	STATIONS								Substrate, Substrate Type
	S1	S2	S3	S4	S5	S6	S7	S8	
Phylum: Tunicata									
Classis: Ascidiacea									
Order: Aplousobranchia Lahille, 1886									
Family: Didemnidae Giard, 1872									
<i>Didemnum</i> sp.			C						on Lamellibranch, N
Family: Polyclinidae Milne Edwards, 1841									
<i>Aplidium</i> sp.			C			C			Wheels, A
* <i>Polyclinum constellatum</i> Savigny, 1816						C		C	Ropes, A
Order: Phlebobranchia Lahille, 1886									
Family: Ascidiidae Herdman, 1882									
<i>Ascidia mentula</i> Müller, 1776				1	1		1		Ropes, wheels, buoys, A
<i>Asciella aspersa</i> (Müller, 1776)	4				2				Ropes, wheels, buoys, A
Family: Cionidae Lahille, 1887									
<i>Ciona robusta</i> Hoshino & Tokioka, 1967	4		15		2	1			Ropes, A
Order: Stolidobranchia Lahille, 1886									
Family: Molgulidae Lacaze-Duthiers, 1877									
<i>Molgula manhattensis</i> (De Kay, 1843)				4			3	1	Ropes, A
Family: Pyuridae Hartmeyer, 1908									
<i>Microcosmus exasperatus</i> Heller, 1878								3	Ropes, A
<i>Microcosmus</i> sp.	4	61			62	8	1	1	Ropes, wheels, buoys, on <i>S. plicata</i> , A, N
Family: Styelidae Sluiter, 1895									
<i>Botrylloides leachii</i> (Savigny, 1816)	C		C	C	C	C		C	Ropes, wheels, buoys, on bivalve shell and <i>S. plicata</i> , A, N
<i>Botryllus schlosseri</i> (Pallas, 1766)	C					C			Wheels, A
<i>Styela canopus</i> (Savigny, 1816)	5	6	3	12	21	5	7	25	on <i>S. plicata</i> , <i>M. squamiger</i> , N
<i>Styela plicata</i> (Lesueur, 1823)	13	8	10	9	30	7	29	25	Ropes, wheels, buoys, A
** <i>Symplegma brakenhielmi</i> (Michaelsen, 1904)								C	on Lamellibranch, N
Colonial	+	+	+	+	+	+		+	
Total species number	8	4	5	5	7	8	6	8	
Total number of soliter ascidians at stations	32	75	28	26	118	21	45	55	

*First record for the Turkish Sea fauna. **First record for the Aegean Sea fauna.

The ascidian species identified in the study area along with their systematic classification and ecological information (substrate type) are presented in Table 2. The highest ascidian species richness (8) occurred at stations S1, S6, and S8, while the lowest species richness (2) was found at Station S2. Of these recorded species, S3 and S4 stations were represented by five species followed by S7 station six species and S5 station seven species (Table 2).

A high abundance of solitary ascidians was recorded at Station S5 situated in Özdere (118 individuals) with *S. plicata*, *S. canopus*, *Microcosmus* sp., and *B. leachii* being the most frequent species. However, the lowest number of individuals was encountered at station S6 situated in Kuşadası (21) (Table 2).

The species *Microcosmus* sp. accounted for 34.6%, *S. plicata* 33.1%, *S. canopus* 21.2%, *C. robusta* 5.6%, *M. manhattensis* 2.0%, *A. aspersa* 1.5%, *A. mentula* 1.3%, and *M. exasperatus* 0.8% of the solitary ascidian community at all sampling stations.

According to frequency index values, *S. plicata* was constant at S1, S8, and S7 stations, *S. canopus* was constant at S4 and S8 stations, *Microcosmus* sp. was constant at S2 and S5 stations,

and also *C. robusta* was constant at S3 (Table 3).

Non-Indigenous Ascidians

Species were classified by origin status as native, cryptogenic or established non-indigenous (Zenetos *et al.*, 2017). The species, *Ascidia mentula* Müller, 1776 was classified as native (Figure 2), while *B. leachii*, *Botryllus schlosseri*, *Molgula manhattensis*, and *S. canopus* labeled as cryptogenic (Figure 3) in the Mediterranean Sea. Seven species, namely such as *P. constellatum*, *Asciella aspersa*, *C. robusta*, *M. exasperatus*, *S. plicata*, and *S. brakenhielmi* were assigned as established non-indigenous species in the Mediterranean Sea (Figure 2, 3). However, three taxa, which were identified only to the genus level, could not be assigned to any of the above categories.

Taxonomic Remarks

Order APLOUSOBRANCHIA Lahille, F. (1886)

Family POLYCLINIDAE Milne Edwards, 1841

Genus *Polyclinum* Savigny, 1816

Polyclinum constellatum Savigny, 1816

Sites (Figure 1): Station S6, S8




Figure 2. a) *Ascidia mentula* Müller, 1776, S5-Özdere, 03.07.2016; b) *A. mentula*, S5-Özdere, 03.07.2016, dissected; c) *A. mentula*, S5-Özdere, 03.07.2016, stigmata and papillae; d) *Ascidiella aspersa* (Müller, 1776), S5-Özdere, 03.07.2016, stigmata and papillae; e) *A. aspersa*, S5-Özdere, 03.07.2016; f) *Ciona robusta* Hoshino & Tokioka, 1967, S6-Kuşadası, 04.07.2016, without tunic; g) *C. robusta*, S6-Kuşadası, 04.07.2016, dorsal tubercule; h) *C. robusta*, S6-Kuşadası, 04.07.2016, dissected; i) *C. robusta*, S6-Kuşadası, 04.07.2016, stigmata and papillae; j) *Molgula manhattensis* (De Kay, 1843), S5-Özdere, 03.07.2016; k) *Microcosmus exasperatus* Heller, 1878, S8-Akbük, 06.07.2016; l) *M. exasperatus*, S8-Akbük, 06.07.2016. Scale bar a, f, h, j: 3 mm; b, e, k, l: 2 mm; g: 1 mm.

Table 3. Frequency index values of solitary taxa (F<25: Rare, 25≤F≤49: Common, F>49: Constant)

Species	S1	S2	S3	S4	S5	S6	S7	S8
<i>Ascidia mentula</i>	6.25			3.85	0.85		2.44	
<i>Ascidiella aspersa</i>	12.50				1.69			
<i>Ciona robusta</i>	12.50		53.57		1.69	4.76		
<i>Molgula manhattensis</i>				15.38			7.32	1.82
<i>Microcosmus exasperatus</i>								5.45
<i>Microcosmus</i> sp.	12.50	81.33			52.54	38.10	2.44	1.82
<i>Styela canopus</i>	15.63	8.00	10.71	46.15	17.80	23.81	17.07	45.45
<i>Styela plicata</i>	40.63	10.67	35.71	34.62	25.42	33.33	70.73	45.45

Examined material: 19 colonies, S6, Kuşadası, 37°51'41.15" (N) 27°15'20.10" (E), 1 m, 04 July 2016
Examined material: 4 colonies, S8, Akbük (Didim), 37°23'11.70" (N) 27°25'54.48" (E), 1 m, 06 July 2016

The non-indigenous *P. constellatum* Savigny, 1816, which is reported for the first time in Turkish waters, was especially abundant at stations S6 and S8. The biggest colony of *P. constellatum*, which collected from station S6, was measured 6.9 cm in

diameter by 5.3 cm in height. It formed ball-shaped colonies filled with a gelatinous tunic, with the zooids located only around the outer edge (Goodbody, 2003) (Figure 4a-d). It had similar characteristics as reported by Rocha and Costa (2005). The studied colonies of *P. constellatum* were encrusting and on the colony surface, they had dark green with white or beige stellate systems (Figure 4d). There were from 10 to 20 rows of stigmata and transverse muscle fibers




Figure 3. a) *Microcosmus* sp.S6-Kuşadası, 04.07.2016, without tunic; b) *Microcosmus* sp., S6-Kuşadası, 04.07.2016, dorsal tubercule; c) *Microcosmus* sp., S6-Kuşadası, 04.07.2016, intestine and gonads; d) *Botrylloides leachii* (Savigny, 1816), S8-Akbük, 06.07.2016; e) *Botryllus schlosseri* (Pallas, 1766), S6-Kuşadası, 04.07.2016; f) *B. schlosseri*, S6-Kuşadası, 04.07.2016, zooids; g) *Styela canopus* (Savigny, 1816) S7-Taşburun, 06.07.2016; without tunic; h) *S. canopus*, S7-Taşburun, 06.07.2016; dissected; i) *Styela plicata* (Lesueur, 1823), S2-Yeni Foça, 02.07.2016, dissected; j) *S. plicata*, S2-Yeni Foça, 02.07.2016, stigmata and papillae; k) *Symplegma brakenhielmi* (Michaelsen, 1904), S8-Akbük, 06.07.2016; l) *S. brakenhielmi*, S8-Akbük, 06.07.2016, zooid; Scale bar a, b, f, g, h, i: 3 mm; c: 2 mm; l: 1 mm.

between them (Figure 4e). The abdomen was short with the distal end of the esophagus bending at a right angle, leaving the stomach horizontal to the thorax. It brooded inside the atrial cavity or in a small pouch projecting from the thorax at the anus level (Figure 4f).

The shape of oral siphon was tubular, fringed by six long and triangular lobes (Figure 4g). And also it had a large circular atrial siphon and its dorsal languet had diminute lobes. Its margin of pharyngeal transverse vessels was papillated (Rocha, Zanata, & Moren, 2012) (Figure 4h). The stomach was globular with a smooth wall and the mid-intestine was short (Figure 4g-i). The intestine was twisted and the bilobed anus ended between the seventh and the ninth rows of stigmata. Anus was located at the 8th transverse vessel in zooids (Figure 4g). Its abdomen was more or less 90° position in relation to the thorax. The post-abdomen was sac-like and the ovary of this species surrounded by male follicles (Rocha & Costa, 2005) (Figure 4e-i).

The established NIS *S. brakenhielmi* formed flattened, encrusting colonies of gray-green color. It had gonads in both of sides of the body wall, and the slightly lobed testis. The maximum colony diameter was 3.3 cm. The colony was observed at station S8

and it is reported for the first time in the Aegean ecoregion (Antoniadou *et al.*, 2016). Due to having, tubercular prominences as small raised regions on the surface of the tunic of the siphons *C. intestinalis* type A individuals in this study were accepted as *C. robusta* (Brunetti *et al.*, 2015). The measured minimum and maximum lengths of the solitary species are presented at Table 4.

Distribution Patterns of the Species

A literature search was done to investigate the identified species distribution in various regions of the world and in Turkish Seas. Worldwide and Turkish Seas distribution of the recorded ascidian species were presented in Table 5.

Physicochemical Variables

The mean±standard errors (SE) of physicochemical variables of the seawater samples at all stations are given in Table 6. The mean water temperature levels ranged from 25.70±0.04 °C at Station S8 to 28.67±0.02 °C at Station S3. As for salinity, the present data showed that sampling

stations revealed narrow variations, the mean salinity values changed from 31.73 ± 0.21 at Station S6 to 40.43 ± 0.02 ppt at Station S7 (Table 6).

A great variability was not observed in recorded pH levels. Moreover, the water pH is slightly basic and the mean pH values which ranged from 7.77 ± 0.01 at Station S6 to 8.09 ± 0.01 at Station S7. In addition, the water conductivity was slightly basic and the mean conductivity levels varied from 48.77 ± 0.74 at Station S6 to 60.70 ± 0.01 mS/cm at Station S7. The average DO values changed between 5.18 ± 0.01 at Station S6 and 7.01 ± 0.05 mg/l at Station S3 (Table 6).

Correlation with Abiotic Variables

According to Spearman's rank correlation analysis there was a negative correlation between temperature ($^{\circ}\text{C}$) and the number of species ($r = -0.7120$, $P < 0.05$). Moreover, there was a positive relationship between *S. plicata* and pH ($r = 0.7143$, $P < 0.05$).

Discussion

The current study aimed to characterize the ascidian assemblages of the eastern Aegean coasts and investigate the presence of non-indigenous species. According to the findings of this study, *S. plicata* and *S. canopus* were frequently occurring species at all sampling stations. Skinner, Barboza and Rocha (2016) mentioned that *S. plicata*, *S. canopus*, *S. brackenhielmi*, *P. constellatum*, and *M. exasperatus* have a wide geographic distribution and are generally found in marinas and harbors. *Ascidiella aspersa* is also another widespread species that is introduced in many parts of the world. Moreover, it is assigned as an established alien species in the Mediterranean Sea (López-Legentil *et al.*, 2015; Zenetos *et al.*, 2010).

Stations with the highest species richness (S1, S6, and S8) hosted a high number of alien species. The increasing maritime traffic (e.g. fishing vessels and international cruise ships) and anthropogenic impacts (e.g. domestic sewage discharge) at these A




Figure 4. (a-d) *Polyclinum constellatum* Savigny 1816-colony, S6-Kuşadası, 04.07.2016, (e-i) zooid of *P. constellatum*, S6-Kuşadası, 04.07.2016. Scale bar d: 2 cm; e-i: 1 mm.

Table 4. The measured minimum and maximum lengths of solitary species at the sampling stations

Species	Minimum Length (cm)	Stations	Maximum Length (cm)	Stations
<i>Ascidia mentula</i>	2.8	S7	4.6	S7
<i>Ascidiella aspersa</i>	1.9	S1	4.4	S1
<i>Ciona robusta</i>	1.9	S1	11.3	S3
<i>Microcosmus exasperatus</i>	1.9	S7	2.8	S7
<i>Microcosmus sp.</i>	4.9	S2	4.9	S2
<i>Styela canopus</i>	0.2	S3	2.1	S4
<i>Styela plicata</i>	0.8	S5	7.5	S3

Table 5. Worldwide and Turkish Seas distribution of the recorded ascidian species in this study

	Worldwide Distribution	Turkish Seas Distribution
<i>Polyclinum constellatum</i> Savigny, 1816	Guam, Brazil, Pacific Mexico, Atlantic Canada, Brazil, Trinidad, France, North Pacific Ocean, Panama, Bermuda, USA (Florida, Gulf of Mexico), South Africa, West Indian Ocean, West Pacific, Hawaii, South California, Suez Canal, Mediterranean Sea.	Kuşadası Bay, Akbük Bay
<i>Ascidia mentula</i> Müller, 1776	Mediterranean, Adriatic to the Straits of Gibraltar, the Atlantic coast of France, Spain into the English Channel, Greece, Algericas Bay (Southern Spain), Baléares, Tunisia Lagoons.	Dardanelles, Kepez Harbor, Aegean Sea, Black Sea, Levantine Sea
<i>Ascidiella aspersa</i> (Müller, 1776)	Argentina, New England, New Zealand, South Africa, South Australia, Tasmania	Black Sea, İzmir Bay, Aegean Sea, Black Sea, Sea of Marmara, Levantine Sea
<i>Ciona robusta</i> Hoshino & Tokioka, 1967	Mediterranean Sea, the Pacific Ocean (Australia, Japan, New Zealand, South Korea, West coast of North America), the Atlantic coasts of South Africa.	Black Sea, Sea of Marmara, Levantine Sea
<i>Molgula manhattensis</i> (De Kay, 1843)	California Harbors, China/Korea, Europe, NE Pacific, Netherlands, South Australia, Tasmania, Western Atlantic, Sweden, Norway, UK, Portugal, Spain, Morocco, Canada, USA, Argentina.	Aegean Sea, Sea of Marmara, Mersin Bay
<i>Microcosmus exasperatus</i> Heller, 1878	Atlantic Panama, Guam, Hawaii, India, Mediterranean Sea, Trinidad, Bermuda, Caribbean, SE USA, Brazil, South Africa, West Africa, Australia, Tunisia, Lebanon, Israel, North-Western Cyprus.	Alsancak Harbor (İzmir Bay), Levantine Sea
<i>Botrylloides leachii</i> (Savigny, 1816)	South Australia, Tasmania, Northeast Atlantic, Indo West Pacific, Mediterranean, the Black Sea, Australia, Europe.	Mersin Harbor
<i>Botryllus schlosseri</i> (Pallas, 1766)	Atlantic Canada, California Harbors, India, New England, San Francisco Bay, South Africa, South Australia, Tasmania, US West coast.	Marmara Sea, Black Sea, Aegean Sea, Mersin Harbor, Marmara Sea, Mersin Harbor, İzmir Bay
<i>Styela canopus</i> (Savigny, 1816)	Atlantic Panama, Atlantic Canada, Adriatic Sea, Brazil, Mexico, California Harbors, Mediterranean, Guam, Indian Ocean, Australia, West Pacific, Hawaii, Easter Island.	Mersin Harbor, Aegean Sea
<i>Styela plicata</i> (Lesueur, 1823)	Atlantic Canada, Brazil, California Harbors, China/Korea, Gulf of Mexico, South Africa, South Australia, Tasmania, the Mediterranean Sea (Iberian Peninsula), the Northeastern Atlantic Ocean (Iberian Peninsula, Canary Islands), the Northwestern Atlantic Ocean (US east coast), the Southwestern Atlantic ocean (Brazil), the Northwestern Pacific Ocean (Japan and China), the Southwestern Pacific Ocean (Australia), the Northeastern Pacific Ocean (US West coast), the Southwestern Indian Ocean (South Africa), Hong Kong, China, East Asian Seas.	Mersin Harbor, Aegean Sea
<i>Symplegma brakenhielmi</i> (Michaelsen, 1904)	California Harbors, Guam, Hawaii, Mediterranean Sea, Trinidad, Caribbean, Brazil, Atlantic Oceans, Indian Ocean, West Pacific, Lebanon, Israel.	Levantine Sea, Fethiye Bays, Finike Bays

number of invasive ascidian species introductions have been linked to hull fouling stations could facilitate the introduction and spread of non-native species from recreational boating (Lambert & Lambert, 1998; Smiard *et al.*, 2017).

In this study, *S. plicata* and *S. canopus* were observed at all sampling stations (Table 2).

Styela plicata is a solitary ascidian native to NW Pacific Ocean (Carlton, 2009; Pineda, López-Legentil, & Turon, 2011) and *S. canopus* is originated from the Indo-West Pacific (Kott, 1985). They were commonly found in warm and temperate oceans, usually colonizing widely fluctuating environments, particularly marinas, harbours, and places used for

Table 6. Physicochemical variables (mean±SE) of the seawater at each sampling station

Stations	Temperature (°C)	Salinity (ppt)	pH	Conductivity (mS/cm)	DO (mg/l)
S1	26.03±0.27	33.53±0.13	7.80±0.02	51.23±0.38	6.03±0.01
S2	27.33±0.05	35.03±0.11	7.84±0.01	52.97±0.42	6.09±0.03
S3	28.67±0.02	39.50±0.00	8.04±0.00	58.87±0.06	7.01±0.05
S4	28.13±0.05	39.53±0.05	7.92±0.01	59.20±0.10	6.52±0.04
S5	28.30±0.04	40.03±0.02	8.05±0.01	59.63±0.12	6.82±0.02
S6	26.10±0.04	31.73±0.21	7.77±0.01	48.77±0.74	5.18±0.01
S7	26.63±0.11	40.43±0.02	8.09±0.01	60.70±0.01	6.67±0.01
S8	25.70±0.04	38.20±0.09	7.91±0.01	57.07±0.38	5.56±0.07

recreational and commercial fishing and boating. These species are able to survive in disturbed habitats and may dominate the sessile communities and reduce local diversity (Stachowicz, Whitlatch, & Osman, 1999). In addition, *S. plicata* (Lesueur, 1823) has been categorized as a harmful invasive species in several regions around the world (Pineda, López-Legentil, & Turon, 2011). The alien species *S. plicata*, *S. canopus*, *P. constellatum*, and *S. brackenhielmi* have been reported from both polluted and non-polluted environments (Marins, Novaes, Rocha, & Junqueira, 2010; Skinner, Barboza, & Rocha, 2016).

The results of this study showed that collected *S. plicata* individuals were fouled by *S. canopus* and *Microcosmus* sp. specimens. According to Yakovis and Artemieva (2017) most ascidians generally prefer to reside on barnacles, empty barnacle tests, and other ascidians.

In the present study, a negative relation was found between temperature (°C) and the number of species. Thiyagarajan and Qian (2003); Tracy and Reyns (2014) mentioned that temperature is an important factor which can affect the community composition, distribution and diversity of ascidians.

The circumtropical ascidian *S. brackenhielmi* is globally distributed in the Atlantic (Van Name, 1921; Monniot, 1983, Goodbody, 2003, Rocha *et al.* 2005), Indian (Millar, 1954; Monniot & Monniot, 1997) and Pacific Oceans (Monniot & Monniot, 2001; Lambert 2003). It occurs in the Red Sea and the Suez Canal (Michaelsen, 1918) and according to Abdel Messeih (1994) it can be a Lessepsian migrant. It has not previously reported from the Aegean Sea (Çınar, 2014; Antoniadou *et al.*, 2016). In the Mediterranean, it was reported from Lebanon (Bitar & Kouli-Bitar, 2001; Bitar, Ocana, & Ramos-Esplá, 2007), Israel (Shenkar, 2008) and the Levantine coast of Turkey (Çınar *et al.*, 2006). Moreover, it was also reported from Fethiye and Finike Bays (the Turkish Levantine Sea/Mediterranean coast) (Ulman *et al.*, 2017).

The species *M. exasperatus* has probably reached the Mediterranean via the Suez Canal; it has an extra-tropical Indo-Pacific distribution and its distribution in the eastern Mediterranean is restricted (Izquierdo-Muñoz, Diaz-Valdéz, & Ramos-Esplá, 2009; Shenkar & Loya, 2009). Alien species of tropical/subtropical Indo-Pacific origin, which are

present in Egypt, Israel, and Turkey were probably introduced via the Suez Canal, shipping traffic, ballast water, fouling of ship hulls, aquaculture, and aquarium trade (Zenetos *et al.*, 2010; Galil *et al.*, 2016). Transportation, shipping activities, ballast water, and hull fouling of recreational yachts, boats, and fishing vessels can be important vectors for the spread of ascidian species (Ashton, Boos, Shucksmith, & Cook, 2006).

The invasive tunicate *P. constellatum* is recorded for the first time from Turkish coastal waters (Table 5). Being a fouling organism, it often develops on artificial substrata in harbours (Monniot & Monniot, 1997). In the Mediterranean Sea, it was first recorded from Port Said in Egypt by Halim and Abdel Messeih (2016). This species has an invasive potential (Locke, 2009) so that it should be investigated.

Invasive alien species and NIS are a focus of good environmental status in the Marine Strategy Framework Directive (European Commission, 2008) and Aichi Biodiversity Target 9 (CBD, 2010), respectively due to their significant ecological and economic impacts on marine communities, aquaculture, and humans. As a result, invasive and alien ascidians should be monitored regularly to avoid possible economic and/or environmental impact. Additional taxonomic and molecular studies are needed to enhance knowledge about the ascidian diversity and NIS of the Turkish Seas.

Acknowledgements

I would like to express my very great appreciation to Dr. Noa Shenkar for kind assistance in teaching identification techniques of ascidians, for her constant encouragement and valuable helps. I would like to thank to Dr. Rosana Moreira da Rocha and who very kindly shared her taxonomic expertise and valuable publications on ascidians. I wish to express my gratitude to Dr. Gretchen Lambert for providing helpful information to identify ascidian specimens. Special thanks to Onur Önen for all his help during the field study. I am grateful to Dr. Şermin Açıık for providing me the camera for species photos and her valuable helps. Thanks are due to the several anonymous reviewers for their constructive suggestions and comments that helped to upgrade this

manuscript. Finally, I gratefully acknowledge all the people who help in some aspect to this work.

References

- Abdel Messeih, M. (1994). *Taxonomical Studies on Ascidiaceans in the Egyptian Mediterranean Waters and the Suez Canal* (Unpublished Ph.D. thesis) Faculty of Science, University of Alexandria, Alexandria, Egypt.
- Antoniadou, C., Gerovasileiou, V., & Bailly, N. (2016). Ascidiacea (Chordata: Tunicata) of Greece: an updated checklist. *Biodiversity Data Journal*, 4: e9273. <https://doi.org/10.3897/BDJ.4.e9273>
- Artüz, M.L., Sönmez, B., & Kubanç, C. (2014). First record of *Phyllochaetopterus socialis* (Annelida: Polychaeta) from a hydrothermal vent site in the Sea of Marmara, Turkey. *Marine Biodiversity Records*, 7: 1-8. <http://dx.doi.org/10.1017/S1755267214000347>
- Ashton, G., Boos, K., Shucksmith, R., & Cook, E. (2006). Rapid assessment of the distribution of marine non-native species in marinas in Scotland. *Aquatic Invasions*, 1(4): 209-213. <http://dx.doi.org/10.3391/ai.2006.1.4.3>
- Aslan, H. (2006). Ascidianlar ve Akdeniz Havzasındaki dağılımları. *Ege Üniversitesi Su Ürünleri Dergisi*, 23(1/1): 29-31.
- Aslan-Cihangir, H., Izquierdo-Muñoz, A., Pancucci-Papadopoulou, M.A., Ramos-Esplá, A., & Can-Yılmaz, E. (2011). *Microcosmus polymorphus* Heller, 1877 (Tunicata: Ascidiacea: Pyuridae)-a new addition to the fauna of the Turkish coasts. *Turkish Journal of Fisheries and Aquatic Sciences*, 11: 163-165. <http://dx.doi.org/10.4194/trjfas.2011.0121>
- Băcescu, M. (1961). Le rôle des îles dans la dispersion récentes des espèces indo-pacifiques en Méditerranée occidentale et quelques observations sur la faune marine de L'île des Serpents, en comparaison avec celle peuplée par les parages bosphoriens de la Mer Noire". In: V. Le peuplement des îles Méditerranéennes et la problématique de l'insularité, Banyuls-sur-Mer. Paris, France: Centre National de la Recherche Scientifique, 241-253 (in French).
- Bitar, G., & Kouli-Bitar, S. (2001). Nouvelles données sur la faune et la flore benthiques de la côte Libanaise. Migration Lessepsienne [New data on benthic fauna and flora on the coast Lebanese. Lessepsian migration]. *Thalassia Salentina* 25:71-74.
- Bitar, G., Ocana, O., & Ramos-Esplá, A.A. (2007). Contribution of the Red Sea alien species to structuring some benthic biocenosis in the Lebanon coast (Eastern Mediterranean). *Rapports de la Commission Internationale pour l'Exploration de la Mer Méditerranée*, 38:437.
- Brunetti, R., Gissi, C., Pennati, R., Caicci, F., Gasparini, F., & Manni L. (2015). Morphological evidence that the molecularly determined *Ciona intestinalis* type A and type B are different species: *Ciona robusta* and *Ciona intestinalis*. *Journal of Zoological Systematics and Evolutionary Research*, 53: 186-193. <http://dx.doi.org/10.1111/jzs.12101>
- Carlton, J.T. (2009). Deep Invasion Ecology and the Assembly of Communities in Historical Time. Rilov, G., Crooks, J.A., editors. Biological Invasions in Marine Ecosystems: Ecological, Management, and Geographic Perspectives: *Ecological Studies*, 204:13-56.
- Çinar, M.E., Bilecenoglu, M., Öztürk, B., & Can, A. (2006a). New records of alien species on the Levantine coast of Turkey. *Aquatic Invasions*, 1: 84-90. <http://dx.doi.org/10.3391/ai.2006.1.2.6>
- Çinar, M.E., Katağan, T., Öztürk, B., Egemen, Ö., Ergen, Z., Kocatas, A., Önen, M., Kirkim, F., Bakir, K., Kurt, G., Dağlı, E., Kaymakçı, A., Açık, Ş., Doğan, A., & Özcan, T. (2006b). Temporal changes of soft-bottom zoobenthic communities in and around Alsancak Harbour (Izmir Bay, Aegean Sea) with special attention to the autecology of exotic species. *Marine Ecology*, 27: 229-246. <http://dx.doi.org/10.1111/j.1439-0485.2006.00102.x>
- Çinar, M.E., Katağan, T., Koçak, F., Öztürk, B., Ergen, Z., Kocatas, A., Önen, M., Kirkim, F., Bakir, K., Kurt, G., Dağlı, E., Açık, Ş., Doğan, A., & Özcan, T. (2008). Faunal assemblages of the mussel *Mytilus galloprovincialis* in and around Alsancak Harbour (Izmir Bay, eastern Mediterranean) with special emphasis on alien species. *Journal of Marine Systems*, 71: 1-17. <http://dx.doi.org/10.1016/j.jmarsys.2007.05.004>
- Çinar, M.E., Bilecenoglu, M., Öztürk, B., Katağan, T., Yokeş, M.B., Aysel, V., Dağlı, E., Açık, Ş., Özcan, T., & Erdoğan, H. (2011). An updated review of alien species on the coasts of Turkey. *Mediterranean Marine Science*, 12: 257-315. <http://dx.doi.org/10.12681/mms.34>
- Çinar, M.E. (2014). Checklist of the phyla Platyhelminthes, Xenacoelomorpha, Nematoda, Acanthocephala, Myxozoa, Tardigrada, Cephalorhyncha, Nemertea, Echiura, Brachiopoda, Phoronida, Chaetognatha, and Chordata (Tunicata, Cephalochordata, and Hemichordata) from the coasts of Turkey. *Turkish Journal of Zoology*, 38: 698-722. <http://dx.doi.org/10.3906/zoo-1405-70>
- Çinar, M.E. (2016). The alien ascidian *Styela clava* now invading the Sea of Marmara (Tunicata: Ascidiacea). *ZooKeys*. 563: 1-10. <http://dx.doi.org/10.3897/zookeys.563.6836>
- Colombo, A. (1885). Raccolte zoologiche eseguite dal R. Pirascafa Washington nella campagna abissale Talassografica dell'anno. *Rivista Marittima*, 18: 2253 (in Italian).
- Convention for Biodiversity (CBD), (2010) Decision X/2. Strategic plan for biodiversity 2011-2020 and the Aichi biodiversity targets, Nagoya, Japan, 18-29 October 2010.
- Demir, M. (1952). Boğaz ve Adalar Sahillerinin omurgasız dip hayvanları. İstanbul, Turkey: İstanbul University (in Turkish).
- Demir, M. (1954a). Report on the plankton of the southeastern coast of the Black Sea. *Hidrobiologia*, 1: 284-286.
- Demir, M. (1954b). Türkiye'ye sahil veren iki büyük içdenizin tipik olan bazı fizikoşimik hususiyetleri ve bunların faunaya tesirleri. *Hidrobioloji Mecbuaları Serisi*, A 2: 144-163 (in Turkish).
- Diñçaslan, Y.E., & Öber, A. (2004). İzmir Körfezi'nde yaşayan bazı Ascidiacea (Tunicata) üyelerinin dorsal tüberkül yapıları. *Ege Üniversitesi Fen Fakültesi Dergileri Serisi*, 21: 119-121.
- EC. (2008). Marine strategy framework directive (2008/56/EC). Retrieved from <http://eurlex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2008:164:0019:0040:EN:PDF>

- Ergen, Z. (1967). İzmir Körfezi'nin tespit edilen başlıca planktonik organizmaları. *Ege Üniversitesi Fen Fakültesi İlimi Raporlar Serisi*, 47: 1-27 (in Turkish).
- Ergen, Z., Kocataş, A., Katarğan, T., & Çınar, M.E. (1994). Gencelli Limanı (Aliağa-İzmir) bentik faunası. *Ege Üniversitesi Fen Fakültesi Dergileri Serisi B*, 16: 1047-1059.
- Gab-Alla, A.F.A. (2008). Distribution of the sea squirt *Ecteinascidia thurstoni* Herdman, 1890 (Ascidiacea: Perophoridae) along Suez Canal and Egyptian Red Sea coasts. Institute of Oceanology PAS, Sopot, Poland. *Oceanologia*, 50(2): 239-253.
- Galil, B.S., Marchini, A., & Occhipinti-Ambrogi, A. (2016). East is east and West is west? Management of marine bioinvasions in the Mediterranean Sea. *Estuarine Coastal and Shelf Science*, <http://dx.doi.org/10.1016/j.ecss.2015.12.021>
- Geldiay, R., & Kocataş, A. (1972). Note preliminaire sur les peuplements benthiques du golfe d'Izmir. *Scientific Monographs of the Faculty of Science, Ege University*, 12: 3-33 (in French).
- Goodbody, I. (2003). The ascidian fauna of Port Royal, Jamaica. I. Harbor and mangrove dwelling species. *Bulletin of Marine Science*, 73(2): 457-476.
- Gopalakrishnan, S., Meenakshi, V.K., & Shanmuga, P.D. (2011). Antipyretic and Analgesic activity of *Phallusia nigra* Savigny, 1816. *Annals of Biological Research*, 2 (4): 192-6.
- Gökalg, N. (1972). Edremit, Bodrum ve İskenderun Körfezleri'nin plankton durumunun karşılaştırmalı incelenmesi, *İstanbul Üniversitesi Hidrobioloji Araştırma Enstitüsü Yayınları*, 3: 1-71 (in Turkish).
- Gözcüoğlu, B. (2011). Denizlerimizin Sakinleri İstanbul, Turkey: Gökce Ofset Basım Yayın Sanayi.
- Izquierdo-Muñoz, A., Diaz-Valdéz, M., & Ramos-Esplá, A.A. (2009). Recent non-indigenous ascidians in the Mediterranean Sea. *Aquatic Invasions*, 4: 59-64. <http://dx.doi.org/10.3391/ai.2009.4.1.5>
- Katsanevakis, S., Wallentinus, I., Zenetos, A., Leppäkoski, E., Çınar, M.E., Öztürk, B., Grabowski, M., Golani, D., & Cardoso, A.C. (2014). Impacts of marine invasive alien species on ecosystem services and biodiversity: a pan-European review. *Aquatic Invasions*, 9(4): 391-423.
- Kiseleva, M.I. (1961). Qualitative and quantitative aspects of benthos in the Aegean coast of the Dardanel. *USSR Science Academy Studies in Sevastopol Biological Station*, 14: 135-146.
- Kiseleva, M.I. (1969). Composition and distribution of benthos in the near-Bosporus Black Sea region. In: Vodyanitsky, V.A., Greze, V.N., Dolgopolskaya, M.A., Pavlova, E.V., editors. Water cycle through Bosporus and his influence on hydrology and biology of the Black Sea. Nauk Dum Kiev, 233-253.
- Kocataş, A. (1978). İzmir Körfezi kayalık sahillerinin bentik formları üzerinde kalitatif ve kantitatif araştırmalar. *Scientific Monographs of the Faculty of Science, Ege University*, 12: 1-93.
- Koçak, F., Ergen, Z., & Çınar, M.E. (1999). Fouling organisms and their developments in a polluted and an unpolluted marina in the Aegean Sea (Turkey). *Ophelia*, 50(1): 1-20.
- Koçak, F., & Küçüksezgin, F. (2000). Sessile fouling organisms and environmental parameters in the marinas of the Turkish Aegean coast. *Indian Journal Marine Science*, 29: 149-157.
- Kott, P. (1985). The Australian Ascidiacea, Part 1: Phlebobranchia and Stolidobranchia. *Memories of Queensland Museum*, 23: 1-440.
- Kott, P. (1990). The Australian Ascidiacea, Part 2: Aplousobranchia (1). *Memories of Queensland Museum*, 29(1): 1-226.
- Kott, P. (1992). The Australian Ascidiacea, Part 3: Aplousobranchia (2). *Memories of Queensland Museum*, 32(2): 375-620.
- Kott, P. (2001). The Australian Ascidiacea, Part 4: Aplousobranchia (3), Didemnidae. *Memories of Queensland Museum*, 47(1): 1-407.
- Lakkis, S., & Toklu, B. (2007). Contribution to the study of zooplankton community in Iskenderun Bay (North-eastern Mediterranean), *Rapport Commission internationale Mer Méditerranée*, 38: 527.
- Lambert, C.C., & Lambert, G. (1998) Non-indigenous ascidians in southern California harbors and marinas. *Marine Biology*, 130: 675-688
- Lambert, G. (2003). Marine biodiversity of Guam: the Ascidiacea. *Micronesica*, 35/36: 588-597.
- Lambert, G., Karney, R.C., Rhee, W.Y., & Carman, M.R. (2016) Wild and cultured edible tunicates: a review. *Management of Biological Invasions*, 7(1): 59-66. <http://dx.doi.org/10.3391/mbi.2016.7.1.08>
- Lemaire, P. (2011). Evolutionary crossroads in developmental biology: the tunicates. *Development*, 138: 2143-2152. <http://dx.doi.org/10.1242/dev.048975>
- Locke, A. (2009). A screening procedure for potential tunicate invaders of Atlantic Canada. *Aquatic Invasions*, 4(1): 71-79. <http://dx.doi.org/10.3391/ai.2009.4.1.7>
- López-Legentil, S., Legentil, M.L., Erwin, P.M., & Turon, X. (2015). Harbor networks as introduction patchworks: contrasting distribution patterns of native and introduced ascidians. *Biological Invasions*, 17: 1623-1638. <http://dx.doi.org/10.1007/s10530-014-0821-z>
- Marins, F.O., Novaes, R.L.M., Rocha, R.M., & Junqueira, A.O.R. (2010). Non-indigenous ascidians in port and natural environments in a tropical Brazilian bay. *Zoologia*, 27: 213-221. <http://dx.doi.org/10.1590/S1984-46702010000200009>
- Marques, A.C., Klöh, A.S., Migotto, A.E., Cabral, A.C., Rigo, A.P.R., Bettim, AL, Razzolini, E.L., Bettim, A.L., Razzolini, E.L., Cascon, H.M., Bardi, J., Kremer, L.P., Vieira, L.M., Bezerra, L.E.A., Haddad, M.A., Oliveira, Filho, R.R., Gutierrez, S.M.M., Miranda, T.P., Franklin, Jr.W., & Rocha, R.M. (2013). Rapid assessment survey for exotic benthic species in the São Sebastião Channel, Brazil. *Latin America Journal of Aquatic Research*, 41(2):265-285. In: Marques, A.C., Lotufo, L.V.C., Paiva, P.C., Chaves, P.T.C., & Leitão, S.N. (Guest Editors) Exotic benthic species in São Sebastião, Brazil. Proceedings of the 3rd Brazilian Congress of Marine Biology. 10.3856/vol41-issue2-fulltext-6
- Mavili, S. (1987). Türkiye'nin Ege Denizi kıyıları yuzey sularında bulunan Appendicularia (Tunicata) türleri. *Doğa Türk Zooloji Dergisi*, 11: 128-135.
- Michaelsen, W. (1918). Ascidia ptychobranchia und dictyobranchia des Roten Meeres. *Denkschr. Akad. Wiss. Wien. (Math.-Nat. Kl.)* 95(10): 1-120.
- Millar, R.H. (1954). On a collection of ascidians from South Africa. *Proceedings of the Zoological Society of London* 125 (1): 169-22.
- Monniot, C. (1983). Ascidies littorales de Guadeloupe. 4. Styelidae. *Bulletin du Muséum National D'Histoire*

- Naturelle*, 4e sér., A, 5 (2): 423-456.
- Monniot, C., Monniot, F., & Laboute, P. (1991). Coral reef ascidians of New Caledonia, Orstom, Paris, 247 pp.
- Monniot, C., & Monniot, F. (1997). Records of ascidians from Bahrain, Arabian Gulf, with three new species. *Journal of Natural History*, 31: 1623-1643. <http://dx.doi.org/10.1080/00222939700770871>
- Monniot, C., & Monniot, F. (2001). Ascidians from the tropical western Pacific. *Zoosystema*, 23(2): 201-383. <http://dx.doi.org/10.1080/00222939700770871>
- Mutlu, E., & Ergeç, M.B. (2008). Spatio-temporal distribution of soft bottom epibenthic fauna on the Cilician shelf (Turkey) Mediterranean Sea. *Revista de Biología Tropical*, 56: 1919-1946.
- Nalbantoğlu, U. (1955). Uskumru balıklarının mide muhteviyatı. *Hidrobioloji Mecbuaları Serisi*, A, 3, 1-9.
- Okuş, E., Altıok, H., Yüksek, A., Yılmaz, N., Aslan Yılmaz, A., Ünsal Karhan, S., Demirel, N., Müftüoğlu, E., Demir, V., Zeki, S., Kalkan, E., & Taş, S. (2007). Biodiversity in western part of the Fethiye Bay. *Journal of Black Sea/Mediterranean Environment*, 13: 19-34.
- Ostroumoff, A. (1894). Dal'neishie materialyi k estestvennoi istoriyii Bosfora. Le Bulletin de l'Académie impériale des sciences de Saint-Petersbourg, 74: 1-46 (in Russian).
- Ostroumoff, A. (1896). Otchet o dragirovkax i planktonnyix ulovaxekspeditsii "Selyanika", Bulletin de l'Académie impériale des sciences de Saint Petersburg, 5: 33-92 (in Russian).
- Öztürk, B., Aktan, Y., Topaloğlu, B., Keskin, C., Karakulak, S., Ozturk, A.A., Dede, A., & Turkozan, O. (2004). Marine Life of Turkey in the Aegean and Mediterranean Seas, İstanbul, Turkey: Turkish Marine Research Foundation.
- Pennati, R., & Rothbacher, U. (2015). Bioadhesion in ascidians: a developmental and functional genomics perspective. *Interface Focus*, 5: 20140061. <http://dx.doi.org/10.1098/rsfs.2014.0061>
- Pınar, E. (1974). Türkiye'nin bazı limanlarında fouling-boring organizmalar ve antifouling-antiboring boyaların bunlar üzerine etkisi. *Ege Üniversitesi Fen Fakültesi İlmî Raporlar Serisi*, 170: 1-67.
- Pineda, M.C., López-Legentil, S., & Turon, X. (2011). The Whereaboutsof an ancient wanderer: Global phylogeography of the solitary ascidian *Styela plicata*. *PLoS ONE*, 6(9): e25495. <https://doi.org/10.1371/journal.pone.0025495>
- Ramos-Espla, A.A., Izquierdo, A., & Cinar, M.E. (2013). *Microcosmus exasperatus* (Asciacea: Pyuridae), current distribution in the Mediterranean Sea. *Marine Biodiversity Records*, 6: e89. <http://dx.doi.org/10.1017/S1755267213000663>
- Rocha, R.M., & Costa, L.V.G. (2005). Ascidians (Urochordata: Ascidiacea) from Arraial do Cabo, RJ, Brazil. *Iheringia, Série Zoologia*, 95 (1): 57-64. <http://dx.doi.org/10.1590/S1984-4670201200100005>
- Rocha, R.M., Zanata, T.B., & Moren, T.R. (2012). Keys for the identification of families and genera of Atlantic shallow water ascidians. *Biota Neotropica*, 12(1): 269-303. <http://dx.doi.org/10.1590/S1984-4670201200100005>
- Shenkar, N. (2008). *Ecological aspects of the ascidian community along the Israeli coasts* (PhD thesis): Tel Aviv University, Tel Aviv, Israel.
- Shenkar, N., & Loya, Y. (2009). Non-indigenous ascidians along the Mediterranean coast of Israel. *Marine Biodiversity Records*, 2: 1-7. <http://dx.doi.org/10.1017/S1755267209990753>
- Shenkar, N., & Swalla, B.J. (2011). Global diversity of Ascidiacea. *PLoS One*, 6: e20657. <http://dx.doi.org/10.1371/journal.pone.0020657>
- Shenkar, N., Gittenberger, A., Lambert, G., Rius, M., Rocha, R.M., Swalla, B.J., & Turon, X. (2017). Ascidiacea World Database. Accessed through: World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=103732> on 2017-07-10
- Simard, N., Pelletier-Rousseau, M., Clarke Murray, C., McKindsey, C.W., Therriault, T.W., Lacoursière-Roussel, A., Bernier, R., Sephton, D., Drolet, D., Locke, A., Martin, J.L., Drake D.A.R., & McKenzie, C.H. (2017). National Risk Assessment of Recreational Boating as a Vector for Marine Non-indigenous Species. DFO Can. Sci. Advis. Sec. Res. Doc. 2017/006. vi + 95 p.
- Skinner, L.F., Barboza, D.F., & Rocha, R.M. (2016). Rapid Assessment Survey of introduced ascidians in a region with many marinas in the southwest Atlantic Ocean, Brazil. *Management of Biological Invasion*, 7(1): 13-20. <http://dx.doi.org/10.3391/mbi.2016.7.1.03>
- Soyer, J. (1970). Bionomie benthique du plateau continental de la côte catalane française. III. Les peuplements de Copepodes harpacticoides (Crustacea). *Vie et Milieu*, 21: 337-511.
- Stachowicz, J.J., Whitlatch, R.B., & Osman, R.W. (1999). Species diversity and invasion resistance in a marine ecosystem. *Science*, 286: 1577-1579. <http://dx.doi.org/10.1126/science.286.5444.1577>
- Tarkan, A.N. (2000). Abundance and distribution of zooplankton in coastal area of Gökçeada Island (northern Aegean Sea). *Turkish Journal of Marine Science*, 6: 201-214.
- Thiyagarajan, V., & Qian, P.Y. (2003). Effect of temperature, salinity and delayed attachment on development of the solitary ascidian *Styela plicata* (Lesueur). *Journal of Experimental Marine Biology and Ecology*, 290: 133-146. [http://dx.doi.org/10.1016/S0022-0981\(03\)00071-6](http://dx.doi.org/10.1016/S0022-0981(03)00071-6)
- Tracy, B., & Reynolds, N. (2014). Spatial and temporal patterns of native and invasive ascidian assemblages in a southern California embayment. *Aquatic Invasions*, 9: 441-455. <http://dx.doi.org/10.3391/ai.2014.9.4.03>
- Ulman, A., Ferrario, J., Occhipti-Ambrogio, A., Arvanitidis, C., Bandi, A., Bertolino, M., Bogi, C., Chatzigeorgiou, G., Çiçek, B.A., Deidun, A., Ramos-Esplá, A., Koçak, C., Lorenti, M., Martinez-Laiz, G., Merlo, G., Princisgh, E., Scribano, G., & Marchini A. (2017). A massive update of non-indigenous species records in Mediterranean marinas. *PeerJ*, 5: e3954. <http://dx.doi.org/10.7717/peerj.3954>
- Uysal, A. (1976). Türkiye suları Ascidiaları. *İstanbul Üniversitesi Fen Fakültesi Hidrobiyoloji Araştırma Enstitüsü Yayınları*, 15: 1-29.
- Van Name, W.G. (1921). Ascidians of the West Indian region and southeastern United States. *Bulletin of the American Museum of Natural History*, 44: 283-494.
- Van Name, W.G. (1931). New North and South American ascidians. *Bulletin of the American Museum of Natural History*, 61: 207-225.
- Van Name, W.G. (1945). North and South American Ascidians. *Bulletin of the American Museum of*

- Natural History*, 84: 1-476.
- Yakovis, E., & Artemieva, A. (2017). Cockles, barnacles and ascidians compose a subtidal facilitation cascade with multiple hierarchical levels of foundation species. *Scientific Reports*, 7: 1-11. <http://dx.doi.org/10.1038/s41598-017-00260-2>
- Zenetos, A., Çinar, M.E., Crocetta, F., Golani, D., Rosso, A., Servello, G., Shenkar, N., Turon, X., & Verlaque, M. (2017). Uncertainties and validation of alien species catalogues: The Mediterranean as an example. *Estuarine Coastal and Shelf Science*, 191: 171-187. <http://dx.doi.org/10.1016/j.ecss.2017.03.031>
- Zenetos, A., Gofas, S., Verlaque, M., Çinar, M.E., García Raso, J.E., Bianchi, C.N., Morri, C., Azzurro, E., Bilecenoglu, M., Frogliã, C., Siokou, I., Violanti, D., Sfriso, A., San Martín, G., Giangrande, A., Katagan, T., Ballesteros, E., Ramos-Espla, A., Mastrototaro, F., Ocaña, O., Zingone, A., Gambi, M.C., & Streftaris, N. (2010). Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union's Marine strategy framework directive (MSFD). Part I. Spatial distribution. *Mediterranean Marine Science*, 11(2): 381-493. <http://dx.doi.org/10.12681/>
- Zhan, A., Briski, E., Bock, D.G., Ghabooli, S., & MacIsaac, H.J. (2015) Ascidians as models for studying invasion success. *Marine Biology*, 162: 2449-2470. <http://dx.doi.org/10.1007/s00227-015-2734-5>