



## Biochemical Content (Fatty Acids, Sterols, Lipophilic Vitamins, Total Protein, MDA, GSH, GSSG) of *Noctiluca scintillans* in the Southeastern Black Sea

Nurgül Şen Özdemir<sup>1,\*</sup>, Fatma Caf<sup>2</sup>, Ali Muzaffer Feyzioğlu<sup>3</sup>, İlknur Yıldız<sup>4</sup>

<sup>1</sup> Bingöl University, Faculty of Agriculture, Department of Fisheries, Bingöl, Turkey.

<sup>2</sup> Bingöl University, Vocational School of Technical Sciences, Department of Veterinary Medicine, Bingöl, Turkey.

<sup>3</sup> Karadeniz Technical University, Sürmene Faculty of Marine Sciences, Department of Marine Sciences and Technology Engineering, Trabzon, Turkey.

<sup>4</sup> Karadeniz Technical University, Marine Sciences and Technology Institute, Karadeniz Technical University, Trabzon, Turkey.

\* Corresponding Author: Tel.: +90.505 6334072;  
E-mail: nsozdemir@bingol.edu.tr

Received 07 September 2016  
Accepted 11 September 2016

### Abstract

*Noctiluca scintillans* reached the highest abundance (5105 ind/m<sup>3</sup>) in April; and April was an important period in the Black Sea ecosystem because of the seasonal thermocline. While the relationship between temperature and abundance was important ( $P < 0.05$ ,  $r^2 = -0.8$ ), the relationship was not important between salinity and abundance ( $P < 0.05$ ,  $r^2 = -0.06$ ); and Chlorophyll-a (Chl-a) and abundance ( $P < 0.05$ ,  $r^2 = -0.2$ ). It was determined that total lipid was 0.5% in wet weight (WW). It was shown that the main Saturated Fatty Acids (SFA) were 16:0 (24%) and 18:0 (6%), main Monounsaturated Fatty Acids (MUFA) were 18:1 $\omega$ -9c (22%) and 16:1 $\omega$ -7 (4%), main Polyunsaturated Fatty Acids (PUFA) were 18:2 $\omega$ -6c (19%) and 20:4 $\omega$ -6c (8%). Phytosterols were the most important sterols in *N. scintillans*; and stigmaterol was 60  $\mu$ g/gr and  $\beta$ -sterol was 8  $\mu$ g/gr as the most important ones in phytosterols.  $\alpha$ -tocopherol was the highest amount (10.8  $\mu$ g/g) in lipophilic vitamins. It was found that Malondialdehyde (MDA) was 65  $\mu$ g/g, total protein was 10 mg/g, Glutathione (GSH) was 10565  $\mu$ g/g and GSSG was 49  $\mu$ g/g. Consequently, PUFA and  $\alpha$ -tocopherol were the most important biochemical parameters and  $\alpha$ -tocopherol may reduce toxicity in blooms of *N. scintillans* because of inhibitory effect of  $\alpha$ -tocopherol on lipid oxidation. Additionally, we can say that *N. scintillans* is a resistant dinoflagellate against oxidative stress.

**Keywords:** PUFA, phytosterols, toxicity, lipid oxidation.

### Introduction

*Noctiluca scintillans* (Macartney) Kofoid & Swezy is a large-size heterotrophic dinoflagellate. It is reported widely as a "red tide" organism in temperate, subtropical and tropical seas and is well known for its luminescence. *Noctiluca* does not produce toxins but it is reported as a harmful algal with oxygen depletion, gill clogging, and high ammonia levels on the basis of mass mortalities of finfish in farms associated (Okaichi and Nishio, 1976). It has been reported that *N. scintillans* may act as a vector of phycotoxins to higher trophic levels, by feeding on toxigenic microalgae (Escalera *et al.*, 2007). The blooms generally occur from spring to summer in many parts of the world, and can cause extreme events as increasing of the cells concentrate near the surface (Elbrächter and Qi, 1998; Miyaguchi *et al.*, 2006; Baek *et al.*, 2009; Padmakumar *et al.* 2010). The blooms are resulted often in a strong pinkish red or orange in some places as Australia (Hallegraeff, 1991), Thailand (Sriwoon *et al.* 2008), Gulf of Mannar, India (Gopakumar *et al.*, 2009) Japan, Hong

Hong, China (Huang and Qi, 1997), Turkey including the Bosphorus, Marmara Sea, Dardanelles (Unsal *et al.* 2003, Turkoglu *et al.*, 2004, Turkoglu and Buyukates, 2005, Turkoglu, 2010; Turkoglu and Erdogan, 2010; Turkoglu, 2013), the Black Sea (Porumb, 1992; Turkoglu and Koray, 2002; 2004) and in the Southeastern coast of the Black Sea (Kopuz *et al.*, 2014). *N. scintillans* blooms are known to occur after diatom blooms ((Kjørboe, 1998; Tiselius and Kiorobe, 1998; Dela-Cruz *et al.*, 2002) because availability of phytoplankton as prey is one of the important factors for the variation in abundance of *N. scintillans* (Elbrächter and Qi, 1998). Therefore, we used Chl-a concentration as the basic criteria of the phytoplankton biomass. However, the abundance of *N. scintillans* shows temporal variations from seasons to years, and its blooms have been reported to correlate with environmental factors. However, the initial trigger of the largescale bloom formations have not specifically been attributed to the any particular condition. The cause of the large-scale bloom formation in coastal embayments is still controversial (Miyagucci *et al.*, 2006).

*N. scintillans* is an opportunistic omnivorous dinoflagellate and has a rapid reproduction capability, together with polyphagous feeding behavior enables outbursts in populations under favorable conditions, frequently dominating the >200 µm fraction of plankton (Uhlig and Sahling, 1990; Shanks and Walters 1996; Elbrachter and Qi, 1998). The diet of the species consists of a broad spectrum of prey, including phytoplankton, nauplii and eggs of zooplankton (acartia), anchovy, organic detritus, and bacteria (Hattori, 1962; Sekiguchi and Kato, 1976; Schaumann et al., 1988; Kirchner et al., 1996; Elbrachter and Qi, 1998; Quevedo et al., 1999), thus enabling the species to prey on items theoretically higher up in the trophic web. It is one of the common and numerous components of the heterotrophic plankton in the Black Sea (Erkan et al., 2000; Kovalev et al., 2001; Feyzioglu and Sivri, 2003; Özdemir and Ak, 2012; Mikaelyan et al., 2014) and has an important role on the pelagic ecosystem of the Black Sea (Oguz et al., 2001 a, b). Thus, we thought that to obtain information about some biochemical parameters of *N. scintillans* would be useful to create source. - We especially researched these parameters in the period when there was the most abundance. Biochemical parameters as lipids are important biochemical compounds in trophic transfer in marine food webs - because they are rich in carbon with a very high energy value (Parrish, 1988). They can be used as biomarkers in ecology (Budge et al., 2006; Litzow et al., 2006). Especially, PUFA, including docosahexaenoic acid (20:6ω3, DHA), eicosapentaenoic acid (20:5ω3, EPA) and α-linolenic acid (18:3ω3, ALA) etc., are among the crucial classes of fatty acids. They are critical structural components and precursors of signaling molecules, which are involved in many diverse biological and biochemical processes, and important to maintain physiological functions in consumers (Ackman et al. 1980; Sargent et al., 1987; Caramujo et al., 2008; Lund et al., 2008). Additionally, proteins play extremely important roles in most biological processes of living beings, such as enzymatic

catylasis, transport and storage, coordinated motion, mechanical support, immune protection, generation and transmission of nerve impulses, and control of growth and differentiation (Zaia et al., 1998). MDA level constitutes a good marker of peroxidative damage to cell membranes (Gil et al., 2002). Thus, it is used as an indicator of lipid peroxidation. In addition, it is a highly reactive three carbon dialdehyde produced as a byproduct of PUFA peroxidation (Janero, 1990) and also during arachidonic acid (ARA) metabolism for the synthesis of prostaglandins (Marnette, 1999). GSH - plays an important role in protecting these from oxidative damage (López-Barea and Gómez-Ariza, 2006). GSH is the most abundant antioxidant in aerobic cells and is synthesized from l-glutamate, l-cysteine, and l-glycine in two ATP requiring steps catalyzed by the enzymes g-glutamylcysteine ligase and GSH synthetase. The cysteine thiol moiety gives antioxidant properties of GSH. The thiol is oxidized by cellular pro-oxidants, such as free radicals and reactive aldehydes, to form GSSG (Owen and Butterfield, 2010).

To our knowledge, this paper is the first report about biochemical content of *N. scintillans*. The main objective of this study is to investigate some biochemical parameters as lipophilic vitamins, fatty acids, sterols, MDA, GSH and GSSH in *N. scintillans*, which has a wide range of food, almost non-predator and causes blooms; also, to determine the relation between abundance and environmental conditions.

## Materials and Methods

### Study Area and Sampling

The study was performed in the Southern part of the Black Sea at a coastal station with coordinates 40° 57' 12" N - 40° 9' 30" E (Figure 1). *N. scintillans* samples were collected with a 200 µm mesh Hydro-Bios net with a mouth diameter of 110 cm. The vertical haul was used from the entire oxic layer



Figure 1. Sampling station in the Southeastern Black Sea.

starting from a depth of 130 m up to the surface layer. Samplings for abundance were made on a single day for every corresponding month between March 2012 and February 2013 aboard KTU's research vessel Yakamoz. The water depth was 250 m. Water samples were taken using Nansen bottles on the surface layer. Temperature and salinity values were measured with a conductivity-temperature-depth-oxygen CTD profiler (CTD, General Oceanic Idronaut, 316).

#### Abundance of *N. scintillans*

*N. scintillans* cells were counted under BH<sub>2</sub> stereo microscope. The abundance was calculated as the number of individuals in m<sup>3</sup>. In the basis, the volume of filtered water and the total number of individuals were used (Harris et al., 2000).

#### Chlorophyll-a Analysis

Water samples were collected from the surface layer for Chlorophyll-a (Chl-a) assay. The collected samples were filtered through 0.45 µm cellulose acetate membrane. The filtered samples were immediately frozen at -20°C and kept until analysis. Before the analysis, Chl-a was extracted from the filtered samples using 90% acetone, and was assayed with spectrometric method (Parsons et al., 1984).

#### Lipid Extraction

The samples were homogenized with tissue fragmentation buffer. The homogenate was centrifuged at 5000 rpm for 5 min at 4°C. After centrifuging, the supernatant part was used for GSH, MDA, and GSSG. The remaining homogenate pellets were extracted in hexane: isopropanol (3:2 v/v) (Hara and Radin, 1978) and used by centrifuging for the ADEK vitamin and fatty acid analysis.

#### Derivatization and Analysis of Fatty Acids

An aliquot was taken from the supernatant part of the sample pellet and 5 mL 2% H<sub>2</sub>SO<sub>4</sub> in methanol was added. The mixture was vortexed and then kept in the oven for 12h (50°C). Then, after being cooled to room temperature, 5 mL of 5% NaCl was added and then it was vortexed. Fatty Acid Methyl Esters (FAME) were extracted with 2×5 ml hexane. FAME were treated with 5 ml 2% KHCO<sub>3</sub> solution and then the hexane phase was evaporated by the nitrogen flow; and then, samples were taken to auto sampler vials by dissolving in 1 ml fresh hexane (Christie, 1990).

The FAME were analyzed on a SHIMADZU GC 17 Ver. 3 Gas Chromatography (Kyoto, Japan). The GC column was 25 m of long Machery-Nagel (Germany) capillary column with an inner diameter of 0.25 µm and a 25 micron-thick film was used. The

column temperature was kept at 120-220°C, the injection temperature was kept at 240°C, and the detector temperature stayed constant at 280°C. The column temperature program was adjusted as 120-220°C, and the temperature increase was determined to be 5°C/min until 200 and 4°C/min from 200-220°C. It was kept at 220°C for 8 min and the total duration was set as 35 min. Nitrogen gas was used as the carrier gas. Before the analysis of the FAME, mixtures of standard fatty acid methyl esters were injected and the residence time of each fatty acid was determined. After this process, the necessary programming was made and the fatty acid methyl ester mixtures of the samples were analyzed (Christie, 1990).

#### Derivatization and Analysis of Lipophilic Vitamins and Sterols

The 5 mL supernatant was taken to 25 mL tubes with caps and 5mL KOH: methanol (1: 10 v/v) was added. The tubes were vortexed and then they were kept in the oven at 85°C for 15 min. They were taken and cooled to room temperature, and 5 mL of pure water was added and mixed. Lipophilic molecules were extracted with 2×5 mL hexane. The hexane phase was evaporated with nitrogen flow. It was dissolved in 1 mL acetonitrile: methanol (50 + 50% v/v) and then was taken to auto sampler vials and was analyzed.

Sterols were analyzed on a Shimadzu HPLC, by using LC-10 ADVP UV as the visible pump; SPD-10A VP as the detector; CTO- 10ASVP, as column oven; SIL-10ADVP as auto sampler, unit DGU-14A as degasser and Class VP software (Shimadzu, Kyoto Japan). The acetonitrile: methanol (60+40% v/v) was used in the mobile phase. The mobile phase flow rate was 1 mL. UV detector and the Supelcosil LC 18 (15×4.6 cm, 5 µm; Sigma, USA) column were used. Wave lengths of detection were 326 nm for vitamin A, 202 nm for vitamin E and 265 nm for vitamins D, K (Katsanidis and Addis, 1999).

#### Analysis of MDA, GSH, GSSG and Total Protein

The supernatant part was used for MDA, GSH and GSSG. 1mL perchloric acid (10% v/v) was added to 1 mL supernatant. After centrifuged, MDA, GSH and GSSG were analyzed on a Shimadzu HPLC. MDA was analyzed according to Karatepe et al., 2004. GSH was analyzed according to Yilmaz et al., 2009. Peaks were identified by comparison of retention times and spectra (multiwave length scan) of the standard and the samples. Total protein amount in *N. scintillans* was measured by method of Lowry et al., 1951.

#### Statistics

The data obtained were analyzed using the

Analysis of Variance (ANOVA) Method. Used software was STATISTICA 8.0 in case there were any significant differences.

## Results and Discussion

### Hydrography, Chl-a, and the Abundance of *N. scintillans*


Samplings of temperature, salinity, Chl-a, and the abundance of *N. scintillans* were carried out during March 2012-February 2013 (Figure 2). It was determined that *N. scintillans* reached the highest abundance in April during the sampling period (March 2012-February 2013). Therefore, biochemical content of *N. scintillans* was investigated in April.

During the sampling period (March 2012-February 2013) for hydrography, temperatures of sea surface fluctuated from 9°C in March to 28 °C in August; and a seasonal thermocline was observed to start at a depth of 20 m in April when *N. scintillans* was sampled for biochemical content. It was observed that the temperature of the sea surface increased (12°C) with the increase of the temperature in April. Water temperature at a depth of 40 m decreased to 7°C and the cold intermediate layer (CIL) observed during certain periods in the Black Sea dominates the region. CIL is characterized by temperatures less than 8°C (Alkan et al., 2013). In the same sampling period, salinity of the sea surface was 18 ppt and it reached 19 ppt in 100 m depending on the depth. Agirbas et al. (2014) reported that the thickness of the CIL was observed to be larger (65-130 m) in the coastal station of the Southeastern Black Sea (Trabzon coastline) and the salinity varied greatly within the range of 18 to 21‰ in the CIL during March 2010-December 2010. Measurements at Southeastern Coastal Site of the Black Sea (off Trabzon) took a decade (2001-2011) and were documented by Alkan et al. (2013). They indicated that the temperature had the lowest values in the range of 7-10°C during March and the highest values as 25-29°C during July-August for all the years. And also, the surface mixed layer is

characterized by salinity values between 17.5-18.0 psu (Alkan et al., 2013).

The mean integrated concentration of Chl-a during the sampling period (March 2012-February 2013) was 1.08 µg/L in the surface layer. Chl-a was the lowest value in April (0.54 µg/L) and while Chl-a was 1.41 µg L<sup>-1</sup> in March, it decreased sharply in April and increased again to 1.65 µg L<sup>-1</sup> in May (Figure 1). Chl-a concentration was used as the basic criteria of the phytoplankton biomass and researched intensively seasonal, annual and monthly variations (Vedernikov and Demidov, 1993; Oguz et al., 2002; Kopelevich et al., 2002; Moncheva, 2003; Agirbas et al., 2014). The Black Sea is characterized with two seasonal peaks of Chl-a in Autumn and Spring (Lalli and Parsons, 2004). Intense concentrations of Chl-a occurs during autumn and two subsequent secondary peaks in late winter-early spring (February-March) and early summer (June) (Alkan et al., 2013).

During the sampling period (March 2012-February 2013), *N. scintillans* was the most abundant (5105 ind/m<sup>3</sup>) in April, following March (1358 ind/m<sup>3</sup>) and February (888 ind/m<sup>3</sup>) (Figure 2). Although the relation between the temperature and the abundance was important ( $P < 0.05$ ,  $r^2 = -0.8$ ), the relation between the salinity and the abundance was not important ( $P < 0.05$ ,  $r^2 = -0.06$ ). Also, the relation between Chl-a and the abundance was not important ( $P < 0.05$ ,  $r^2 = -0.2$ ). Özdemir and Ak (2012) reported that the rate of *N. scintillans* was 57%, 98% and 85%, respectively, in April, May and June, in all of the zooplankton species in the Southeastern Black Sea (Trabzon Coast). But they did not give individual amounts. In their study, abundance of *N. scintillans* had the highest rate in May. But, only two species represented zooplankton abundance. In the study, the abundance of *N. scintillans* was 57% in April, but the abundance rate was given in five zooplankton species. Also, the influence of the environmental factors should not be forgotten. Therefore the abundance of *N. scintillans* can change in the same season. An unusual blooming of *N. scintillans* was observed in April, 2011 in the same place by Kopuz et al. (2014).



**Figure 2.** Abundance and Chl-a of *N. scintillans* (ind/m<sup>3</sup>) during the sampling periods

## Biochemical Content

In marine ecosystems, lipids provide the densest form of energy which is transferred from algae to vertebrates via zooplankton. Also, they contain essential fatty acids and sterols which are considered to be important drivers of ecosystem health and stability. Fatty acids and sterols are also susceptible to oxidative damage leading to cytotoxicity and a decrease in membrane fluidity. The physical characteristics of biological membranes can be defended from the influence of changing temperature, pressure, or lipid peroxidation by altering the fatty acid and sterol composition of the lipid bilayer. The influence of essential lipids, lipid oxidation, and membrane composition on food web structure and function will become increasingly important in the context of global warming and ozone depletion (Parrish, 2013). In the present study, the biochemical parameters as fatty acids, sterols and lipophilic vitamins have been investigated.

Table 1 represents the FA composition (percent of total FA) of *N. scintillans* from dinoflagellate in April, 2012. The total lipid amount was determined as 0.5% in Wet Weight (WW). Some fatty acids, or their derivatives, are thought to work synergistically with other known toxins such as domoic acid (Wang and Shimizu, 1990; Wang et al., 1993), diarrhetic shellfish poison (DSP) (Lawrence et al., 1994), or the neurotoxin brevetoxin (Boer et al., 2012). Interestingly, de Boer et al. (2012) found that two essential fatty acids, i.e. the C<sub>20</sub> PUFA, ARA and EPA, were more toxic than the C<sub>18</sub> PUFA octadecatetraenoic acid (OTA, 18:4 $\omega$ -3). Since ARA and OTA have the same number of double bonds, and ARA is more toxic than EPA, the degree of unsaturation does not seem to be a key determinant of

toxicity level (Parrish 2013). EPA with 2% and DHA with 3% were the two PUFA mainly detected in April, 2012. Fatty acid composition of algal lipids varies widely with species, habitat, light, salinity, pollution and environmental conditions (Kim et al., 1996; Ratana-Arporn and Chirapart, 2006) and in most studies 16:0 is predominant (Khotimchenko et al., 2002; Li et al., 2002; Gressler et al., 2010; Leblond et al., 2003). Dinoflagellates are the major contributors to the marine food web and are second only to diatoms as primer producers of organic matter in the Ocean (Tappan, 1980). Dinoflagellates can form extensive blooms known as red-tide in coastal areas (Hallegraeff, 1993). Therefore, they are a major source of essential long-chain highly unsaturated fatty acids (HUFA), sterols, and other nutrients to other marine species. They are also an important source of organic matter in marine sediment (De Leeuw et al., 1983). HUFA are high in dinoflagellates (Holz, 1981; Nichols et al., 1984). Also, feeding preference of *N. scintillans* would modify the quantity and distribution of  $\omega$ -3 polyunsaturated fatty acids in the ambient environment (Zhang et al. 2015; Zhang et al., 2016). Moreover, altering the amount and ratio of some HUFA (DHA and EPA) available to higher trophic levels, such as copepods, fish larvae or hydromedusae (Zhang et al., 2016).

Mansour et al. (1999) reported that saturated fatty acids were dominated by 16:0 and its concentration ranged from 9.0% to 24.8% in terms of the total fatty acids in different marine dinoflagellates species. They indicated that MUFA and PUFA's level changed according to the species. While they existed at high levels in some species, they existed at low levels in some species. In this study, the major MUFA were oleic acid (18:1 $\omega$ -9) in *N. scintillans* (22%) and it was the second most abundant compound. The

**Table 1.** Fatty acid composition of *N. scintillans*

Fatty Acids	%
16:0	24.34±0.91
C18:0	6.11 ±0.56
C23:0	1.03 ±0.08
∑SFA	31.48±0.43
16:1 $\omega$ -7	3.96±0.18
18:1 $\omega$ -9t	1.03±0.43
18:1 $\omega$ -9c	21.99±1.34
20:1 $\omega$ -9	0.83±0.06
∑MUFA	27.81±1.49
18:2 $\omega$ -6t	1.68±0.08
18:2 $\omega$ -6c	18.49±2.26
18:3 $\omega$ -3c	7.12±0.69
20:4 $\omega$ -6	8.13±0.76
20:5 $\omega$ -3 (EPA)	2.21±0.06
22:6 $\omega$ -3 (DHA)	3.08±0.11
∑PUFA	40.71±2.22
∑HUFA	13.42±0.81

same result was given by Leblond *et al.* (2003) and they determined that 16:0 was the first dominant and, 18:1 $\omega$ -9 was the second dominant fatty acid in *Karenia brevis* (dinoflagellate). But, Holz (1981) and Parrish *et al.* (1994) reported that MUFA was not usually abundant in dinoflagellates. Mansour *et al.* (1999) observed that the elevated content (11.8%) of in *Gymnodinium sp.* (dinoflagellate) was unexpected. The fatty acid profile can distinguish among algal classes but it may be limited use for the identification of species within the same algal class (Holz, 1981; Volkman, 1989). Viso and Marty (1993) determined the profiles of fatty acids of nine different marine algal groups, and they were even able to define the species-specific lipid compositions.

Free sterols are dominated in sediments of the Black Sea and the sterols are clearly of dinoflagellate origin (Boon *et al.*, 1979). The sterols are essential for all eukaryotes. They are components of the membranes and have a function in regulation of the membrane fluidity and permeability. Sterols also play an important role as precursors of many steroid hormones including vitamin D and brassinosteroids as well as for a wide range of secondary metabolites such as saponins and glycolalkaloids (Piironen *et al.*, 2000). Also, sterols are a source of oxygenated lipids: the oxysterols which are the sterols bearing a second oxygen function. These compounds have a variety of biological properties including cytotoxicity and effects on specific enzymes (Parrish, 1991). Phytosterols are present in small amounts, and two common examples are the stigmasterol and the sitosterol (Abidi, 2001). Phytosterols are bioactive compounds, which can be found in a great variety of plant-based foods (Brufau *et al.*, 2008). Phytosterols were the most important sterols in *N. scintillans*. Stigmasterol was 60  $\mu$ g/g and  $\beta$ -sterol was 8  $\mu$ g/g from phytosterols.

In our study, it was shown that  $\alpha$ -tocopherol (vitamin E) had the highest amount (10.8  $\mu$ g/g) in lipophilic vitamins, following D<sub>2</sub> (3.5  $\mu$ g/g) and K<sub>2</sub> (0.2  $\mu$ g/g).  $\alpha$ -tocopherol is an important antioxidant, and antioxidants play a major role in reducing or preventing the lipid peroxidation (Aitken *et al.*, 1989). One of the most effective antioxidants is  $\alpha$ -tocopherol as it breaks the free radical chain reaction by forming a relatively stable tocopheroxyl radical (Selley *et al.*, 1991).  $\alpha$ -tocopherol is identified as the major naturally occurring tocopherol in the lipids of marine organisms (Ackman and Cormier, 1978; Parazo *et al.*, 1998). It is essential for integrity and optimal function of reproductive, muscular, circulatory, nervous and immune systems (Hoekstra, 1975; Sheffy and Schultz, 1979; McDowell, 2000). The most important determinant of vitamin E requirements is the dietary concentration of unsaturated fatty acids as PUFA are highly susceptible to auto-oxidation (Nacka *et al.*, 2001). Animals ingesting high levels of PUFA require high concentrations of vitamin E to protect tissue lipids from free radical attack (Debieer *et al.*, 2002;

Lammi-Keefe and Jensen, 1984; Machlin, 1991). PUFA and  $\alpha$ -tocopherol prevent the occurrence of oxidative stress in the marine organisms, and a correlation between increased dietary PUFA and  $\alpha$ -tocopherol requirement has been found in blue tilapia (Roen *et al.* 1990) turbot (Stephan *et al.* 1995), carp (Runge *et al.*, 1992) and Atlantic salmon (Waagboe 1991). High level  $\alpha$ -tocopherol probably reflects higher degree of antioxidant protection required  $\omega$ -3 PUFA-rich organisms (Hamre and Lie 1995). It has also an accelerating effect on reproduction (Kahn-Thomas and Enesco, 1982; Shalaby *et al.*, 2004). This effect could be possibly attributed to the well-known antioxidant effect of  $\alpha$ -tocopherol (Shalaby *et al.*, 2004). Therefore, higher  $\alpha$ -tocopherol amount than the other lipophilic vitamins may cause blooms in suitable temperature condition, because of reproduction accelerating effect of the  $\alpha$ -tocopherol in *N. scintillans*. Additionally, the most important reason of indirect toxicity of *N. scintillans* may be the inhibitory effect of  $\alpha$ -tocopherol on lipid oxidation.

In this study, the MDA level was found as 65  $\mu$ g/g, the total protein amount was found as 10 mg/g, the GSH level was found as 10565  $\mu$ g/g and the GSSG level was found as 49  $\mu$ g/g in *N. scintillans*. MDA was used as an indicator of lipid peroxidation, and protein carbonyl content, which provides an indication of protein oxidative damage. It is a highly reactive three carbon dialdehyde produced as a byproduct of polyunsaturated fatty acid peroxidation (Janero, 1990) and also during arachidonic acid metabolism for the synthesis of prostaglandins (Marmette, 1999). MDA can combine with several functional groups on molecules including proteins, lipoproteins, RNA and DNA (Sevilla *et al.*, 1997). GSH is a key contributor to the cellular antioxidant defense system and to the maintenance of the intracellular redox milieu for the preservation of thiol disulfide redox states of proteins. GSH is also involved in cellular signaling, regulation and redox activation of transcription factors, and thiol-disulfide exchange reactions. GSH oxidation to GSSG are resulted in intracellular redox imbalance as reflected in a decreased GSH-to-GSSG ratio (GSH/GSSG), a condition often associated with oxidative stress (Dalle-Donne *et al.*, 2007). In our study, GSH/GSSG was measured as 216  $\mu$ g/g in *N. scintillans*. It was shown that MDA level was lower than GSH, and GSH/GSSG was quite high. We can say that *N. scintillans* is a resistant dinoflagellate against oxidative stress. Fidan *et al.* (2008) showed that GSH, as natural antioxidant, levels were found to be the same as MDA due to increase and decrease of oxidative stress in *Carassius carassius* in Eber Lake. MDA level increases when the natural antioxidant system is insufficient (Şahan *et al.*, 2003). The increase of its level in winter could be because of the cold conditions of the fish environment, and the lipid catabolism occurs in cold seasons to get much more energy for their metabolisms (Fidan *et al.*, 2008).

Because, Fidan *et al.* (2008) indicated that MDA levels varied during the seasons and the highest level was in winter and the lowest level was in spring and summer seasons. They observed that the highest GSH level was in winter season.

## Conclusion

*N. scintillans* is one of the most important heterotrophic dinoflagellates in the Black Sea ecosystem. It creates blooms. But it is not toxic. In this study, some biochemical parameters of *N. scintillans* were investigated in the most abundant periods (April) of *N. scintillans* together with biological and physiological factors. We determined that April was an important period in the Black Sea because seasonal thermocline occurred and the abundance of *N. scintillans* reached the highest amount in April. Consequently, we found a negative relationship between the temperature and the abundance of *N. scintillans*. However, the between the salinity, Chl-a and abundance of *N. scintillans* was not a relationship. Additionally, we can say that PUFA,  $\alpha$ -tocopherol and GSH were the most important parameters in bloom of *N. scintillans*.  $\alpha$ -tocopherol and GSH may reduce toxicity in *N. scintillans* because of inhibitory effect of  $\alpha$ -tocopherol on lipid oxidation, and key role in the cellular antioxidant defense system of GSH and *N. scintillans* is a resistant dinoflagellate against oxidative stress. Future studies are required to establish new methods for a few cell. In addition, effect of biochemical content on the bloom formation of *N. scintillans* should research with annually series. However, it is difficult to provide enough sample amounts for biochemical analyses on *N. scintillans* and enough sample amount can provide in most abundant or bloom periods. Therefore we thought that the study would contribute to future studies about biochemical content of *N. scintillans*. Moreover, the study is the first known record about biochemical content of *N. scintillans*.

## References

- Abidi, S.L. 2001. Chromatographic analysis of plant sterols in foods and vegetable oils. *Journal of Chromatography, A* 935(1-2): 173-201. doi: 10.1016/S0021-9673(01)00946-3.
- Ackman, R.G. and Cormier, M.G. 1978.  $\alpha$ -Tocopherol in some Atlantic fish and shellfish with particular reference to live holding without food. *Journal of the Fisheries Research Board of Canada*, 24: 357-373. doi:10.1139/f67-031.
- Ackman, R., Sebedio, J. and Kovacs, M. 1980. Role of eicosenoic and docosenoic fatty acids in fresh water and marine lipids. *Marine Chemistry*, 9: 157-164. doi: 10.1016/0304-4203(80)90034-1.
- Agirbas, E., Feyzioglu, A.M. and Kopuz, U. 2014. Seasonal Changes of Phytoplankton Chlorophyll a, Primary Production and their Relation in the Continental Shelf Area of the South Eastern Black Sea. *Turkish Journal of Fisheries and Aquatic Sciences*, 14: 713-726. doi: 10.4194/1303-2712-v14\_3\_14.
- Aitken, R.J., Clarkson, J.S. and Fische, L.S. 1989. Generation of reactive oxygen species lipid peroxidation and human sperm function, *Biology of Reproduction*, 41: 83-197. doi: 10.1095/biolreprod41.1.183.
- Alkan, A., Zengin, B., Serdar, S. and Oğuz, T. 2013. Long-Term (2001-2011) Temperature, salinity and Chlorophyll-a variations at a Southeastern Coastal site of the Black Sea. *Turkish Journal of Fisheries and Aquatic Sciences*, 13: 57-68. doi: 10.4194/1303-2712-V143\_1\_08.
- Baek, S.H., Shimode, S., Kim, H.C., Han, M.Y. and Kikuchi, T. 2009. Strong bottom-up effects on phytoplankton community caused by a rainfall during spring and summer in Sagami Bay, Japan. *Journal of Marine Systems*, 75: 253-264. doi: 10.1016/j.jmarsys.2008.10.005.
- Boon, Rijkstra, W.I.C., de Lange, F., De Leeuw, J.W., Yoshioka and M., Shimizu, Y. 1979. Black Sea sterol-a molecular fossil for dinoflagellate blooms. *Nature*, 277: 125-127. doi: 10.1038/277125a0.
- Brufau, G., Canela, M.A. and Rafecas, M. 2008. Phytosterols: physiologic and metabolic aspects related to cholesterol-lowering properties. *Nutrition Research* 28: 217-25. doi: 10.1016/j.nutres.2008.02.003.
- Budge, S.M., Iverson, S.J. and Koopman, H.N. 2006. Studying trophic ecology in marine ecosystems using fatty acids: a primer on analysis and interpretation. *Marine Mammal Science*, 22: 759-801. doi: 10.1111/j.1748-7692.2006.00079.x.
- Christie, W.W. 1990. Gas chromatography and lipids: A practical guide. Bridgewater, Somerset: The Oily Press.
- Caramujo, M.J., Boschker, H.T. and Admiraal, W. 2008. Fatty acid profiles of algae mark the development and composition of harpacticoid copepods. *Freshwater Biology*, 53: 77-90.
- Dalle-Donne, I., Rossi, R., Giustarini, D., Colombo, R. and Milzani, A. 2007. S-glutathionylation in protein redox regulation. *Free Radical Biology and Medicine*, 43: 883-898. doi: 10.1016/j.freeradbiomed.2007.06.014.
- De Boer, M.K., Boeree, C., Sjollem, S.B., de Vries, T., Rijnsdorp, A.D., Buma, A.G.J. 2012. The toxic effect of the marine raphidophyte *Fibrocapsa japonica* on larvae of the common flatfish sole (*Solea solea*), *Harmful Algae*, 17: 92-101. doi: 10.1016/j.hal.2012.03.005.
- De Leeuw, J.W., Rijkstra, W.I.C., Schenck, P. A. and Volkman, J.K. 1983. Free, esterified, and residual bound sterols in Black Sea Unit I sediments. *Geochimica Cosmochimica Acta*, 47:455-465. doi: 10.1016/0016-7037(83)90268-5.
- Debier, C., Pomeroy, Van, P.P., Wouwe, N, Mingolet, E., Baret, P.V. and Larondelle, Y. 2002. Dynamics of vitamin A in grey seal (*Halichoerus grypus*) mothers and pups throughout lactation. *Canadian Journal of Zoology*, 80: 1262-1273. doi: 10.1139/z02-107.
- Dela-Cruz, J., Ajani, P., Middleton, H. J. and Suthers, M. I. (2003) Population growth and transport of the red tide dinoflagellate *Noctiluca scintillans*, in the coastal waters off Sydney, Australia, using cell diameter as a tracer. *Limnology and Oceanography*, 48: 656-674.

- doi:10.4319/lo.2003.48.2.0656.
- Elbrachter, M. and Qi, Y.Z. 1998. Aspects of *Noctiluca* (Dinophyceae) population Dynamics. In: Physiological ecology of harmful algal blooms, (eds., DM. Anderson, AD. Cambella, GM. Hallegraeff), Springer, London, pp. 662.
- Erkan, F., Gucu, A. and Zagorodnyaya, J. 2000. The diel Vertical distribution of zooplankton in the Southeast Black Sea. Turkish Journal of Zoology, 24: 417-427.
- Escalera, L., Pazos, Y., Morono and A., Reguera, B. 2007. *Noctiluca scintillans* may act as a vector of toxigenic microalgae. Harmful Algae, 6:317-320. doi: 10.1016/j.hal.2006.04.006.
- Feyzioglu, A.M. and Sivri, N. 2003. Seasonal changes of *Noctiluca scintillans* Kofoids and Swezy in Trabzon Coast, Eastern Black Sea. E.U. Journal of Fisheries and Aquatic Sciences 20: (1-2): 75- 79. doi: 10.12714/egefas.2003.20.1.5000157043.
- Fidan, A.F., Cigerci, I.H., Konuk, M., Küçükuyurt, I., Aslan, R. and Dündar, Y. 2008. Determination of some heavy metal levels and oxidative status in *Carassius carassius* L., 1758 from Eber Lake, Environmental Monitoring and Assessment, 147(1): 35-41. doi: 10.1007/s10661-007-0095-3.
- Gil, P., Farinas, F., Casado, A. and Lo'pez-Ferna'ndez, E. 2002. Malondialdehyde: a possible marker of ageing. Gerontology, 48: 209-214. doi: 10.1159/000058352.
- Quevedo, M., Gonzalez-Quiros, R. and Anadon, R. 1999. Evidence of heavy predation by *Noctiluca scintillans* on *Acartia clausi* (copepoda) eggs off the central Cantabrian coast (NW Spain). Oceanologica, 22: 127-131. doi: 10.1016/S0399-1784(99)80039-5.
- Gopakumar, G., Sulochanan, B. and Venkatesan, V. 2009. Bloom of *Noctiluca scintillans* (Macartney) in Gulf of Mannar, southeast coast of India. Journal Marine Biological Association of India, 51(1): 75- 80.
- Gressler, V., Yokoya, N., Fujii, M., Colepicolo, P., Filho, J., Torres, R. and Pinto, E. 2010. Lipid, fatty acid, protein, amino acid and ash contents in four Brazilian red algae species. Food Chemistry, 120:585-590. doi: 10.1016/j.foodchem.2009.10.028.
- Hallegraeff, G.M., 1993. A review of harmful algal blooms and their apparent global increase. Phycologia, 32: 79-99. doi: 10.2216/i0031-8884-32-2-79.1.
- Hallegraeff, G.M. 1991. Aquaculturists guide to harmful Australian microalgae. 1st edn., Fish. Ind. Train. Board, Tasmania/CSIRO Div. Fisher., Hobart, pp 111.
- Hamre, K. and Lie, O. 1995. Alpha-tocopherol levels in different organs of Atlantic salmon (*Salmo salar* L.) - effect of smoltification, dietary levels of n-3 polyunsaturated fatty acids and vitamin E. Comp. Biochemistry & Physiology, 111(A): 547-554.
- Harris, R., Wiebe, P., Lenz, J., Skjoldal, H.R. and Huntley, M. 2000. ICES zooplankton methodology manual. Academic Press, UK, pp 684.
- Hattori, S. 1962. Predatory activity of *Noctiluca* on Anchovy eggs. Bull. Tokai Reg. Fish. Res. Lad. 9: 211-220.
- Hoekstra, W.G. 1975. Biochemical function of selenium and its relation to vitamin E. Federation Proceedings, 34: 2083-2089.
- Holz Jr., G.G. 1981. Non-isoprenoid lipids and lipid metabolism of marine flagellates. In: Bio- chemistry and Physiology of Protozoa, (eds., M Levandowsky, SH Hunter), Academic Press, New York, pp. 301-332.
- Huang, C. and Qi, Y. 1997. The abundance cycle and influence factors on red tide phenomena of *Noctiluca scintillans* (Dinophyceae) Dapeng Bay, the South China Sea. Journal of Plankton Research, 19(3): 303-318. doi: 10.1093/plankt/19.3.303.
- Janero, D.R. 1990. Malondialdehyde and thiobarbituric acid-reactivity as diagnostic indices of lipid peroxidation and peroxidative tissue injury. Free Radical Biology and Medicine, 9: 515-540. doi: 10.1016/0891-5849(90)90131-2.
- Kahn-Thomas, M. and Enesco, H.E. 1982. Effect of  $\alpha$ -tocopherol and culture method on reproduction of *Turbatrix aceti*. Journal of Nematology, 14(4):496-500.
- Karatepe, M. 2004. Simultaneous determination of ascorbic acid and free malondialdehyde in human serum by HPLC/UV. LC-GC North America, 22(4): 362-365.
- Katsanidi, E. and Addis, P.B. 1999. Novel HPLC analysis of tocopherols and cholesterol in tissue. Free Radical Biology and Medicine, 27: 1137-1140. doi:10.1016/S0891-5849(99)00205-1.
- Kim, M., Dubacq, J., Thomas, J. and Giraud, G. 1996. Seasonal variations of triacylglycerols and fatty acids in *Fucus serratus*. Phytochemistry, 43: 49-55. doi: 10.1016/0031-9422(96)00243-9.
- Kjørboe, T. 1998. Intensive aggregate formation with low flux during an upwelling-induced diatom bloom. Limnology and Oceanography, 43: 104-116. doi:10.4319/lo.1998.43.1.0104.
- Kirchner, M., Sahling, G., Uhlig, G., Gunkel, W. and Klings, K.W. 1996. Does the red tide-forming dinoflagellate *Noctiluca scintillans* feed on bacteria? Sarsia, 81: 45-55. doi: 10.1080/00364827.1996.10413610.
- Khotimchenko, S., Vaskovsky, V. and Titlyanova, T. 2002. Fatty acids of marine algae from the Pacific coast of North California. Botanica Marina, 45: 17-22. doi: 10.1515/BOT.2002.003.
- Kopelevich, O.V., Sherberstov, S.V., Yunev, O., Baştürk, O., Finenko, Z.Z., Nikonov, S. and Vedernikov, V.I. 2002. Surface chlorophyll in the Black Sea over 1978-1986 derived from satellite and in situ data. Journal of Marine Systems, 36(3-4):145-160. doi: 10.1016/S0924-7963(02)00184-7.
- Kopuz, U., Feyzioglu, A.M. and Valente, A. 2014. An unusual red-tide event of *Noctiluca scintillans* (Macartney) in the Southeastern Black Sea. Turkish Journal of Fisheries and Aquatic Sciences, 14: 261-268. doi: 10.4194/1303-2712-V14\_1\_28.
- Kovalev, A.V., Mazzocchi, M.G., Siokou-Frangou, I. and Kideys, A.E. 2001. Zooplankton of the Black Sea and the Eastern Mediterranean: similarities and dissimilarities. Mediterranean Marine Science, 2(1): 69-77. doi: 10.12681/mms.277.
- Lalli, C.M. and Parsons, T.R. 2004. Biological Oceanography an Introduction. University of British Columbia, Vancouver, Canada, 314.
- Lammi-Keefe, C.J. and Jensen, R.G. 1984. Lipids in human milk: a review. 2: Composition and fat-soluble vitamins. Journal of Pediatric Gastroenterology Nutrition, 3: 172-198. doi: 10.1097/00005176-198403000-00004.
- Lawrence, J.F., Chadha, R.K., Ratnayake, W.M.N. and Truelove, J.F. 1994. An incident of elevated levels of unsaturated free fatty acids in mussels from Nova Scotia and their toxic effect in mice after intraperitoneal injection. Natural Toxins, 2(5): 318-32. doi: 10.1002/nt.2620020511.
- Leblond, J.D., Evans, T.J. and Chapman, P.J. 2003. The



- biochemistry of dinoflagellate lipids, with particular reference to the fatty acid and sterol composition of a *Karenia brevis* bloom. *Phycologia*, 42(20): 324-331. doi: 10.2216/i0031-8884-42-4-324.1.
- Li, X., Fan, X., Han, L. and Lou, Q. 2002. Fatty acids of some algae from the Bohai Sea. *Phytochemistry*, 59: 157-161. doi: 10.1016/S0031-9422(01)00437-X.
- Litzow, M.A., Bailey, K., Prahl, F.G. and Heintz, R. 2006. Climate regime shifts and reorganization of fish communities: the essential fatty acid limitation hypothesis. *Marine Ecology Progress Series*, 315: 1-11. doi: 10.3354/meps315001.
- López-Barea, J. and Gómez-Ariza, J.L. 2006. Environmental proteomics and metallomics. *Proteomics*, 6: S51-S62. doi: 10.1002/pmic.200500374.
- Lowry, O.H., Rosebrough, N.J. and Farr, A.L. 1951. Randall RJ: Protein measurement with folin phenol reagent. *The Journal of Biological Chemistry*, 193 (1): 265-275.
- Lund, E.D., Chu, F.L., Harvey, E. and Adlof, R. 2008. Mechanism(s) of long chain n-3 essential fatty acid production in two species of heterotrophic protists: *Oxyrrhis marina* and *Gyrodinium dominans*. *Marine Biology*, 155: 23-36. doi: 10.1007/s00227-008-1003-2.
- Machlin, L.J. 1991. *Handbook of Vitamins*. Marcel Dekker, Inc., New York and Basel.
- McDowell, L.R. 2000. *Vitamins in animal and human nutrition*. (ed., I.A. Ames), Iowa State University Press, pp. 793. doi: 10.1002/9780470376911.
- Mansour, M.P., Volkman, J.K., Jackson, A.E. and Blackburn, S.I. 1999. The fatty acid and sterol composition of five marine dinoflagellates. *Journal of Phycology*, 35: 710-720. doi: 10.1046/j.1529-8817.1999.3540710.x.
- Marnette, L.J. 1999. Generation of mutagens during arachidonic acid metabolism. *Cancer and Metastasis Reviews*, 13: 303-308. doi: 10.1007/BF00666100.
- Mikaelyan, A.S., Malej, A., Shiganova, T.A., Turk, V., Sivkovitch, A.E., Musaeva, E.I., Kogovsek, T. and Lukasheva, T.A. 2014. Populations of the red tide forming dinoflagellate *Noctiluca scintillans* (Macartney): A comparison between the Black Sea and the Northern Adriatic Sea. *Harmful Algae*, 33: 29-40. doi: 10.1016/j.hal.2014.01.004.
- Miyaguchi, H., Fujiki, T., Kikuchi, T., Kuwaha, V.S. and Toda, T. 2006. Relationships between the bloom of *Noctiluca scintillans* and environmental factors in the coastal waters of Sagami Bay, Japan. *Journal of Plankton Research*, 28(3):313-324. doi: 10.1093/plankt/fbi127.
- Monncheva, S. 2003. On the recent state of the Black Sea Ecosystem biological response to eutrophication-are there signs of recovery uncertainties. In: Preliminary Assessment of the previous biogeochemical, hydrophysical, and hydrobiological cruise data within the Black Sea, pp. 53.
- Nacka, F., Cansell, M., Meleard, P. and Combe, N. 2001. Incorporation of alphatocopherol in marine lipid-based liposomes: in vitro and in vivo studies. *Lipids*, 36:1313-1320. doi:10.1007/s11745-001-0846-x.
- Nichols, P., Jones, G.J., de Leeuw, J.W. and Johns, R.B. 1984. The fatty acid and sterol composition of two marine dinoflagellates. *Phytochemistry*, 23:1043-1047. doi: 10.1016/S0031-9422(00)82605-9.
- Oğuz, T., Ducklow, H.W. and Purcell, J.E. 2001a. Modeling the response of top-down control exerted by gelatinous carnivores on the Black Sea pelagic food web. *Journal of Geophysical Research*, 106: 4543-4564. doi: 10.1029/1999JC000078.
- Oğuz, T., Malanotte-Rizzoli, P. and Ducklow, H.W. 2001b. Simulations of phytoplankton seasonal cycle with multi-level and multi-layer physical-ecosystem models: the Black Sea example. *Ecological Modeling*, 144: 295-314. doi: 10.1016/S0304-3800(01)00378-7.
- Oğuz, T., Deshpande, A.G. and Malanotte-Rizzoli, P. 2002. The role of mesoscale processes controlling biological variability in the Black sea coastal waters: interferences from sea WIFS-derived surface chlorophyll field. *Continental Shelf Research*, 22: 1477-1492. doi: 10.1016/S0278-4343(02)00018-3.
- Okaichi, T. and Nishio, S. 1976. Identification of ammonia as the toxic principle of red tide of *Noctiluca miliaris*. *Bulletin of the Plankton Society of Japan*, 23: 75-80.
- Owen, J.B. and Butterfield, D.A. 2010. Measurement of oxidized/reduced glutathione ratio. *Methods in Molecular Biology*, 648: 269-277. doi: 10.1007/978-1-60761-756-3\_18.
- Özdemir, G.P. and Ak, O. 2012. Quality and quantitative Changes of phytoplankton in the South East Black Sea (Trabzon Coasts). *Yunus Research Bulletin*, 4: 13-25.
- Padmakumar, K.B., SreeRanjima, G., Fanimol, C.L., Menon, N.R. and Sanjeevan, V.N. 2010. Preponderance of heterotrophic *Noctiluca scintillans* during a multi-species diatom bloom along the southwest coast of India. *International Journal of Oceans and Oceanography*, 4(1): 55-63.
- Parish, E. 1991. The biosynthesis of oxysterols in plants and microorganisms. In: *Physiology and Biochemistry of Sterols*, (eds., GW Patterson, WD Nes), pp. 324-336.
- Parrish, C.C. 2013. Lipids in marine ecosystems. *ISRN Oceanography*, 2013: 1-16. doi: 10.5402/2013/604045.
- Parsons, T.R., Maita, Y. and Lailli, C.M. 1984. *A manual of chemical and biological methods for Seawater Analysis*. Pergamon Press, Greet Britain, pp. 173.
- Parazo, M.P.M., Lall, S.P., Castell, J.D. and Ackman, R.G. 1998. Distribution of  $\alpha$ - and  $\gamma$ -tocopherols in Atlantic salmon (*Salmo salar*) tissues. *Lipids*, 33: 697-704. doi: 10.1007/s11745-998-0259-x.
- Piironen, V., Lindsay, D.G., Miettinen, T.A., Toivo, J. and Lampi, A.M. 2000. Plant sterols: biosynthesis, biological function and their importance to human nutrition. *Journal of the Science of Food and Agriculture*. 80: 939-966. doi: 10.1002/(SICI)1097-0010(20000515)80:7<939::AID-JSFA644>3.0.CO;2-C.
- Porumb, F. 1992. On the development of *Noctiluca scintillans* under eutrophication of Romanian Black Sea waters. *Science of the total environment*, Supplement 1992, Elsevier Publishers BV, Amsterdam, pp. 907-920.
- Ratana-Arporn, P. and Chirapar, A. 2006. Nutritional evaluation of tropical green seaweeds *Caulerpa lentillifera* and *Ulva reticulata*. *Kasetsart Journal: Natural Sciences*, 40: 75-83.
- Roem, A.J., Kohler, C.C. and Stickney, R.R. 1990. Vitamin E requirement of the blue tilapia, *Oreochromis aureus* (Steindachner), in relation to

- dietary lipid level. *Aquaculture*, 87: 155-164. doi: 10.1016/0044-8486(90)90272-O.
- Runge, G., Steinhart, H. and Schwarz, F.J. M. 1992. Influence of type of fats and  $\alpha$ -tocopherol acetate additions to the feed rations on the tocopherol and tocotrienol composition of carp (*Cyprinus carpio* L.). *Journal of Animal Physiology and Animal Nutrition*, 67: 16-24. doi: 10.1111/j.1439-0396.1992.tb00577.x.
- Shalaby, M.A., El Zorba, H.Y. and Kamel, G.M. 2004. Effect of  $\alpha$ -tocopherol and simvastatin on male fertility in hypercholesterolemic rat. *Pharmacological Research*, 50(2): 137-142. doi: 10.1016/j.phrs.2003.10.013.
- Sargent, J.R., Parkes, R., Mueller-Harvey, I. and Henderson, R. 1987. Lipid biomarkers in marine ecology. In: *Microbes in the sea*, (ed., M.A. Sliegh), Ellis Horwood Limited, Chichester, UK, pp. 119-138.
- Sekiguchi, H. and Kato, T. 1976. Influence of *Noctiluca*'s predation on the Acartia population in Ise Bay, central Japan. *Journal of Oceanographical Society of Japan*, 32: 195-162. doi: 10.1007/BF02107121.
- Shanks, A. and Walters, K. 1996. Feeding by a heterotrophic dinoflagellate in marine snow. *Limnology and Oceanography*, 41: 177-181. doi: 10.4319/lo.1996.41.1.0177.
- Schaumann, K., Gerdes, D. and Hesse K.J. 1988. Hydrographic and biological characteristics of a *Noctiluca scintillans* red-tide in the German Bight, 1984. *Meeresforschung*, 32: 77-91.
- Selley, M.L., Lacey, M.J., Bartlett, M.R. and Copeland, C.M. and Ardlie, N.G. 1991. Content of significant amounts of a cytotoxic endproduct of lipid peroxidation in human semen, *Journal of Reproduction and Fertility*, 92: 291-298. doi: 10.1530/jrf.0.0920291.
- Sevilla, C.L., Mahle, N.H., Eliezer, N., Uzieblo, A., O'Hara, S.M., Nokubo, M., Miller, R., Rouzer, C.A. and Marnett, L.J. 1997. Development of monoclonal antibodies to the malondialdehyde-deoxyguanosine adduct, pyrimidopurine. *Chemical Research in Toxicology*, 10: 172-180. doi: 10.1021/tx960120d.
- Sheffy, B.E. and Schultz, R.D. 1979. Influence of vitamin E and selenium on immune response mechanisms. *Federation Proceedings*, 38: 2139-2143.
- Sriwoon, R., Pholpunthin, P., Lirdwitayaprasit, T. 2008. Population dynamics of green *Noctiluca scintillans* (dinophyceae) associated with the monsoon cycle in the upper gulf of Thailand. *Journal of Phycology*, 44: 605-615. doi: 10.1111/j.1529-8817.2008.00516.x.
- Stéphan, G., Guillaume, J., Lamour, F. 1995. Lipid peroxidation in turbot (*Scophthalmus maximus*) tissue: effect of dietary vitamin E and dietary n-6 or n-3 polyunsaturated fatty acids. *Aquaculture*, 130: 251-268. doi: 10.1016/0044-8486(94)00322-F.
- Şahan, A., Kurutaş, E. and Dikel, S. 2003. Liver Antioxidant Systems and Lipid Peroxidation in Sea Bass (*Dicentrarchus labrax*) Adapted to Fresh Water. *Turkish Journal of Veterinary and Animal Sciences*, 27: 1261-1267.
- Tappan, H. 1980. *The Paleobiology of Plant Protists*. Freeman, San Francisco, pp. 1027.
- Tiselius, P. and Kjørboe, T. 1998. Colonization of diatom aggregates by the dinoflagellate *Noctiluca scintillans*. *Limnology and Oceanography*, 43: 154-159. doi: 10.4319/lo.1998.43.1.0154.
- Turkoglu, M., Unsal, M., Ismen, A., Mavili, S., Sever, T.M., Yenici, E., Kaya, S. and Coker, T. 2004. Dynamics of lower and high food chain of the Dardanelles and Saros Bay (North Aegean Sea). *TUBITAKYDABAG Tech. Fin. Rep.*, No:101Y081, Canakkale, (in Turkish), pp. 314.
- Turkoglu, M. and Buyukates, Y. 2005. Short time variations in density and bio-volume of *Noctiluca scintillans* (Dinophyceae) in Dardanelles. XIII. Natl. Fish. Symp., 01-04 Semptember 2005, Canakkale, Turkey, Abstr. Book (Abstracts), (in Turkish), pp.59.
- Turkoglu, M. 2010. Temporal variations of surface phytoplankton, nutrients and chlorophyll-a in the Dardanelles (Turkish Straits System): A coastal station sample in weekly time intervals. *Turkish Journal of Biology*, 34(3): 319-333.
- Turkoglu, M. and Erdogan, Y. 2010. Diurnal variations of summer phytoplankton and interactions with some physicochemical characteristics under eutrophication of surface water in the Dardanelles (Çanakkale Strait, Turkey). *Turkish Journal of Biology*, 34(2): 211-225.
- Turkoglu, M. and Koray, T. 2002. Phytoplankton species succession and nutrients in Southern Black Sea (Bay of Sinop), *Turkish Journal of Botany*, 26: 235-252.
- Turkoglu, M. and Koray, T. 2004. Algal blooms waters of the Sinop Bay in the Black Sea, Turkey. *Pakistan Journal of Biology Sciences*, 7(9): 1577-1585. doi: 10.3923/pjbs.2004.1577.1585.
- Turkoglu, M. 2013. Red tides of the dinoflagellate *Noctiluca scintillans* associated with eutrophication in the Sea of Marmara (the Dardanelles, Turkey), *Oceanologia*, 55(3): 709-732. doi: 10.5697/oc.55-3.709.
- Uhlig, G. and Sahling G. 1990. Long-term studies on *N. scintillans* in the German Bight population dynamics and red tide phenomena. *Netherlands Journal of Sea Research*, 25: 101-112. doi: 10.1016/0077-7579(90)90012-6.
- Unsal, M., Turkoglu, M. and Yenici, E. 2003. Biological and physicochemical researches in the Dardanelles (Canakkale strait). *TUBITAK-YDABAG Tech. Fin. Rep.*, No: 101Y075, Canakkale, (in Turkish), pp. 92.
- Vedernikov, V.I. and Demidov, A.B. 1993. Primary production and chrophyll in deep regions of the Black Sea, *Oceanology*, 33: 193-199.
- Viso, A.C. and Marty, J.C. 1993. Fatty acids from 28 marine microalgae. *Phytochemistry*, 34: 1521-1533. doi: 10.1016/S0031-9422(00)90839-2.
- Waagbø, R., Sandnes, K., Sandev, A. and Lie, O. 1991. Feeding three levels of n-3 polyunsaturated fatty acids at two levels of vitamin E to Atlantic salmon (*Salmo salar*): Growth and chemical composition. *Fzrk.Dzr. Skr., Ser. Ernreznng*, 4: 51-63.
- Volkman, J.K. 1989. Fatty acids of microalgae used as feedstocks in aquaculture. In: *Fats for the Future*, (ed., RC Cambie), Ellis Horwood Limited: Chichester, UK, pp. 263-283.
- Yilmaz, O., Keser, S., Tuzcu, M., Guvenc, M., Cetintas, B., Irtegun, S., Tastan, H. and Sahin, K. 2009. A Practical HPLC Method to Measure Reduced (GSH) and Oxidized (GSSG) Glutathione Concentrations in Animal Tissues. *Journal of Animal and Veterinary Advances*, 8(2): 343-347.
- Wang, R., Maranda, L., Hargraves, P.E. and Shimizu, Y. 1993. Chemical variation of *Nitzschia pungens* as demonstrated by the co-occurrence of domoic acid and bacillariolides. In: *Toxic Phytoplankton Blooms in the*

- Sea, (eds., T.J. Smayda, Y. Shimizu), Elsevier, pp. 637-641.
- Wang, R. and Shimizu, Y. 1990. Bacillariolides I and II, a new type of cyclopentane eicosanoids from the diatom *Nitzschia pungens*. Journal of the Chemical Society, 5: 413-414. doi: 10.1039/c39900000413.
- Zaia, D.A.M., Zaia, C.T.B.V. and Lichting, J. 1998. Determination of total protein by spectrophotometry: advantages and disadvantages of proposed methods. Quimica Nova, 21(6): 787-793. doi: 10.1590/S0100-40421998000600020.
- Zhang, S., Liu, H., Chen, B. and Wu, C.J. (2015) Effects of diet nutritional quality on the growth and grazing of *Noctiluca scintillans*. Marine Ecology Progress Series, 527: 73-85. doi: 10.3354/meps11219.
- Zhang, S., Liu, H., Guo, C. and Harrison, P.J. (2016). Differential feeding and growth of *Noctiluca scintillans* on monospecific and mixed diets. Marine Ecology Progress Series, 549: 27-40. doi: 10.3354/meps11702.