



Microsatellite Diversity and Population Structure of *Hypophthalmichthys molitrix* in Hatchery Populations of Punjab

Nadia Nazish^{1,*}, Khalid Abbas¹, Sajid Abdullah¹, Muhammad Anjum Zia²

¹ University of Agriculture, Faisalabad, Pakistan, Zoology, Wildlife and Fisheries, Faisalabad-Punjab, Pakistan.

² University of Agriculture, Department of Biochemistry, Faisalabad, Pakistan.

* Corresponding Author: Tel.: +920 3401770610;
E-mail: nadianazish@gmail.com

Received 07 September 2017
Accepted 29 November 2017

Abstract

The genetic integrity of the commercially important fish species is being compromised mainly due to artificial propagation. The present study evaluated the extent of genetic variability and assessed the population structure of Silver Carp (*Hypophthalmichthys molitrix*) in hatchery stocks of Punjab province, Pakistan using 16 species-specific microsatellite loci. The results revealed low-to-moderate scale of genetic variability in the hatchery stocks. The level of average observed heterozygosity varied from 0.3770 to maximum of 0.5874 in all the stocks. Most of the hatcheries showed heterozygote deficit. The average values of inbreeding coefficient ranged from 0.174 to 0.441. Most of the locus-population combinations showed significant departures from Hardy-Weinberg equilibrium. The AMOVA revealed significant genetic structuring in hatchery populations of *H. molitrix*. The UPGMA dendrogram showed the populations clustering irrespective of their geographical proximities. The current research work provides a comprehensive knowledge regarding the genetic structure and genetic diversity of *H. molitrix* populations which may be applicable for the management of the concerned populations to maintain their genetic quality in Pakistan.

Keywords: Silver Carp, DNA markers, population genetics.

Introduction

Like indigenous carps, artificial propagation of exotic carps is also posing a big threat to the genetic integrity of fish species including Silver Carp (*Hypophthalmichthys molitrix*) in Pakistan. For instance, hatchery stock deterioration is much debated issue in fisheries sector due to genetic erosion which can be explained by insufficient number of effective breeders, negative selection, poor broodstock management, hybridization and inbreeding among exotic fish species (Hauser & Seeb, 2008). The alterations in the hydrography and human interventions in the natural water bodies have degraded the nursing and breeding sites of fish (Diana, 2009). Deterioration of water quality and disruption in migration channels resulted in the ecological problems (Van Overzee & Rijnsdorp, 2015). Major cause behind the diminishing exotic fish stocks is overfishing and degraded fish biodiversity, making them more susceptible to climatic variations (Paukert *et al.*, 2016).

With the expansion of aquaculture in Pakistan, the demand for fish seed is also increased. About 99%

of fish seed is produced through artificial propagation by public and private hatcheries in Pakistan to meet requirement of expanding aquaculture practices (Khan, Shakir, Khan, Abid, & Mirza, 2008). Generally, stakeholders maintain few small sized brooders to keep production cost minimum; therefore, there is always a risk of producing poor quality seed (Munilkumar & Nandeesh, 2007). Utilization of small number of fish brooders, generation after generation, may cause reduction in genetic variability and result in indiscriminate hybridization, inbreeding and loss of genetic purity of Chinese fish species (Ellergern & Galtier, 2016). Therefore, regular monitoring of genetic variability in fish hatchery stock is essential to detect whether breeding programs are affecting genetic integrity by reducing genetic variation (De-Cara, Villanueva, Toro, & Fernandez, 2013) which is lacking in Pakistan. By upgrading and managing standards of breeding activities and brood stock in hatchery operations, aquaculture outputs can be enhanced and their influence on capture fisheries biodiversity can be reduced (Xu *et al.*, 2015).

DNA markers techniques are extensively utilized to investigate different issues pertaining to genetics of aquaculture species (Khoo, Lim, & Phang,

2011). Selectively neutral markers can provide an insight into dissemination of genetic diversity within and among the fish populations (Avisé, 2010) and could be helpful to assess the level of genetic diversity in cultured stock (Askari, Shabani, & Miandare, 2013). Microsatellite markers have been considered for analysis of hatchery fish stocks (Defaveri, Viitaniemi, Leder, & Merila, 2013). Moreover, the microsatellite markers have been developed for Silver carp for monitoring the influence of conventional hatchery practices on genetic integrity of species (Liao *et al.*, 2007) to provide valuable information for their effective management in hatcheries. The main objective of the current study was to evaluate the extent of genetic diversity of *H. molitrix* in hatchery populations of Punjab by using microsatellite markers.

Materials and Methods

Specimens of *H. molitrix* were collected from eight selected hatcheries from different districts of the Punjab province of Pakistan (Table 1) designated in map (Figure 1). Thirty individuals from each population were collected in ice boxes and transported to the laboratory for preservation at -20 °C till DNA extraction. DNA extraction from dorsal muscle tissues was done with slight modifications in the methods of Yue & Orben (2005) and that of Green & Sambrook (2012).

The quantitative and the qualitative assessment of DNA was done on 0.8% agarose gel. To get

optimum PCR amplification, the concentration of each DNA sample was quantified for dilution. The quantification of DNA samples was done using UV spectrophotometer (NanoDrop 2000c, Thermo Scientific, USA). The final concentration of the DNA samples was adjusted to almost 50 ng/mL using TE to have uniform concentration of each fish sample.

Sixteen species-specific microsatellite markers were used for genotyping the *H. molitrix* individuals at specific temperature (Liao *et al.*, 2007) (Table 2). The primers were purchased from the company Gene-Link, USA. The PCR reaction was carried out in a 20 µl reaction mixture in labeled PCR tubes using thermal cycler (Multigene, LabNet, USA). The mixture included 0.8 µl of each primer set of 10 µM concentration, 0.4 µl dNTPs of 10 mM concentration, 2.0 µl of 10 x PCR buffer (20 mM), 1.5 µl MgCl₂ (20 mM), 0.4 µl (2 U/ µl) *Taq* DNA polymerase (Fermentas, USA) and about 50 ng of the template DNA. The PCR cycles were adjusted for 5 minutes denaturation at 94°C, 32 cycles of 1 minute at 94°C, 30 seconds at a primer-specific temperature for annealing, 1 minute at 72°C and final elongation for 4 minutes at 72°C. The amplicons were tested using 2% agarose (Pei, Costumbrado, Hsu & Kim, 2012) and finally resolved on polyacrylamide gel followed by silver staining (Qu., Li, Wu, & Yang, 2005). The gel imaging was done in gel imaging system (UVCI, Major Science, USA)

The allelic data obtained through individual genotyping were analyzed by using different analytical programs/software. FSTAT (ver. 2.9.3.2)

Table 1. Sampling details of *H. molitrix* from selected hatcheries of Punjab, Pakistan

Sr. No	Population Name	Code	District	Collection date
1	Central Fish Seed Hatchery, (Manawan)	LHR	Lahore	14.11.2014
2	Fish seed hatchery, Mureedkey	GWH	Gujranwala	19.03.2015
3	Fish seed hatchery, Satiana Road,	FSD	Faisalabad	03.06.2014
4	Fish seed hatchery, Mian Chanu	KWH	Khanewal	14.01.2014
5	Fish seed nursing unit, Farooqabad	SHK	Sheikhupura	27.01.2015
6	Fish seed nursing farm, Sargodha	SGD	Sargodha	30.10.2014
7	Fish Seed Nursing Farm, Sialkot	SKT	Sialkot	02.09.2014
8	Fish seed nursing farm, Toba Tek Singh	TTS	Toba Tek Singh	19.06.2014




Figure 1. UPGMA dendrogram based genetic relatedness between the hatchery populations of *H. Molitrix*.

Table 2. Characterization of microsatellites for *H. molitrix* (Liao et al., 2007)

Locus	Motif	Primer sequence (5'-3')	Annealing temperature (°C)
BL42	(TG)14	F: TGCCGATGTTATGTTTGCT R: TGCTTGTGGGTGAGTTTCT	52
BL56	(AC)16	F: TTAGGTGAACCCAGCAGC R: AAGAAGCATTAGTGCAGATGAGTAC	54
BL64	(TG)23	F: GCCAGGCTAGAAGAACCACC R: TTGCAGCACAGTTACCAAGACA	54
BL67	(AC)17	F: GGCAGGCTTCAAAGGACA R: CTCGCCCTCATTGTGCGTA	54
BL73	(AC)6	F: TGACTTTACACGGCTCCA R: TTA CTCTGTTATGGTGGGTCA	54
BL83	(AC)6	F: CTATCCGCCCTGTTCTGA R: ACCAAACATCCCTCAAGC	54
BL90	(CA)19	F: ATGCGAGGGTGGATGATGGG R: GGAAAGCAAAGCCTGGACTA	52
BL106-2	(AC)14	F: TTTAATTTCTTAGCTGGACACG R: CACTCCTCTCCCTCGTAAAT	54
BL108	(GT)9	F: GATGAATCGCAGGGCGTGAGG R: GCAGAACACGCACAATGGAGA	54
BL109	(TG)21	F: GTGTCCTGGATTCTAGCCG R: CATGAGAGAAACACCTGAACA	54
BL116	(CT)15	F: GCGGGATGAGTTTGAAGAA R: TATGGACTGGACTGCTGGAT	54
BL123	(TG)9	F: GCGACAGGAACAGTAAAAC R: CAAAGAAGGCACAAAGGATT	54
BL132	(AC)10	F: CTTTGACTGCTGGTTGGTTGT R: TTTCTTGCTTCCCTGGCTTCT	54
BL138	(TG)8	F: ACTGAAAACATCACTGCCACG R: ACTGAAAACATCACTGCCACG	54
BL145	(TG)12	F: GTGATTGGACGGGATGAACTA R: TCTTTCTTTTCTGTCCGAGGG	52
BL167	(AC)6	F: AGTGCGCCTAAAAGTAAAAG R: AACCAGGACGAATCAAGTA	52

(Goudet, 2002) and POPGENE (ver. 1.31) (Yeh, Yang, Boyle, Ye, & Mao, 1999) softwares were implemented to calculate the allelic richness (A_r), allelic frequency, expected and observed heterozygosity (H_e & H_o) and inbreeding coefficient F_{IS} . The population differentiation F_{ST} for sampling locations was calculated following Weir and Cockerham (1984). To adjust the statistical significance of deviations from Hardy-Weinberg equilibrium (HWE) sequential Bonferroni correction (Rice, 1989) was used and for each locus sample type-1 error of $\alpha = 0.05$ was maintained. The presence of null alleles was calculated by MicroChecker 2.2.3 (Van Oosterhout, Hutchinson, Wills, & Shipley, 2004). Analysis of molecular variance (AMOVA) was conducted using ARLEQUIN ver. 3.1 (Excoffier, Smouse, & Quattro, 1992) to analyze molecular variance as well as to estimate the hierarchical partitioning in genetic diversity. The software TFGPA ver. 1.3 (Miller, 1997) was employed to construct UPGMA dendrogram based on Nei's (1972) unbiased distance.

Results

The present study assessed the level of genetic

diversity and the population structure of Silver Carp (*Hypophthalmichthys molitrix*) in eight hatchery populations in different districts of Punjab, Pakistan.

In the hatchery stocks of *H. molitrix*, genetic diversity was measured in terms of allelic richness (A_r), observed (H_o) and expected (H_e) heterozygosity, inbreeding coefficient (F_{IS}) and Hardy-Weinberg equilibrium (HWE) over the sixteen microsatellite loci. The data set was analysed for presence and frequency of null alleles at all the loci. There were certain loci which yielded null alleles in an inconsistent pattern. No one of the loci showed null allele frequency more than 10%. All the screened loci were found to be polymorphic by following allele frequency criteria at P 0.95 criterion. The highest average values of allelic diversity were observed in the LHR ($A_r = 6.522$) to a minimum for TTS population ($A_r = 4.732$). The highest value for the allelic diversity ($A_r = 11.93$) was observed in the LHR hatchery population of *H. molitrix* over the locus BL109, while the lowest value at the locus BL67 ($A_r = 2.00$) in SHK population of *H. molitrix* (Table 3).

Low-to-moderate level of average observed heterozygosity was found in the present study that varied from 0.377 (TTS) to a maximum of 0.5875 LHR population of *H. molitrix*. The values of average

Table 3. Genetic structure analysis of *H. molitrix* in the hatchery stock selected from various districts of Punjab, Pakistan

Populatio ns	Parameters																				Average SE
	Loci																				
	BL42	BL56	BL64	BL67	BL73	BL83	BL90	BL106	BL108	BL109	BL116	BL123	BL132	BL138	BL145	BL167	BL167	BL167	BL167	BL167	
LHR	A_v	6	5.998	7.993	8.926	3.933	4.933	7.867	3.997	11.93	3.933	6	6	6	7.997	4	6.522				
	H_v		0.4000	0.4333	0.6667	0.4	0.7	0.6333	0.6333	0.6333	0.7	0.4667	0	0.5667	0.7333	0.6667	0.3333	0.5875			
	H_s		0.5272	0.6428	0.7222	0.7594	0.6217	0.6178	0.73	0.6694	0.8661	0.5606	0.7628	0.7383	0.7728	0.7983	0.7161	0.7055			
	F_{IS}		0.244	0.325	0.094	-0.486	-0.109	-0.008	0.138	0.149	0.209	0.184	0.23	0.248	0.068	0.181	0.271	0.174			
	HWE		↓	0.0490	0.0590	0.0002	1.0000	0.4814	↓	0.0205	0.1572	0.3829	0.2162	0.4540	1.0000	0.1016	0.1259				
SHK	A_v		3.966	4.867	2.997	2	4	5	5.933	6	3.933	6	6	6	4	4	4.968				
	H_v		0.2333	0.3	0.5333	0.4	0.7	0.7	0.4	0.6	0.6333	0.6	0.4667	0.6	0.7	0.4333	0.527				
	H_s		0.5394	0.5533	0.5267	0.42	0.6367	0.7306	0.7172	0.62	0.825	0.5322	0.7211	0.7472	0.7078	0.6367	0.6683	0.6442			
	F_{IS}		0.583	0.471	0.004	0.065	-0.083	0.059	0.456	0.049	0.248	0.202	0.184	0.39	0.169	-0.083	0.366	0.194			
	HWE		0.0022	0.0225	0.7241	1.0000	1.0000	0.7060	0.0010	0.2870	0.2162	0.0546	0.1572	0.2162	0.1549	1.0000	0.0068				
GWH	A_v	5	5.93	2.933	2.933	4	5.997	8.86	6.997	3.997	10.926	3.933	7	6	5.997	5	5.781				
	H_v		0.4667	0.4333	0.6333	0.4333	0.5	0.6	0.4667	0.5667	0.7	0.5333	0.4	0.6333	0.7667	0.5667	0.5333	0.552			
	H_s		0.605	0.59	0.5128	0.455	0.545	0.7528	0.7817	0.76	0.5539	0.8661	0.7556	0.7694	0.8333	0.6928	0.7489	0.6747			
	F_{IS}		0.208	0.262	-0.219	0.065	0.099	0.219	0.248	0.4	-0.006	0.088	0.484	0.193	0.097	0.198	0.303	0.178			
	HWE		0.4655	0.0597	0.4613	0.6817	0.0219	0.0017	1.0000	0.0068	1.0000	0.5886	0.0017	1.0000	1.0000	0.0116	0.1284				
FSD	A_v	3.999	2	2	3	4	4.933	4.933	5	5	10.993	3.933	5.997	5	4.996	5	4.732				
	H_v		0.4333	0.0667	0.1	0.1333	0.4	0.6667	0.3	0.5	0.5333	0.3333	0.3667	0.2333	0.5333	0.4333	0.4	0.377			
	H_s		0.5517	0.5	0.4861	0.545	0.6406	0.7156	0.6433	0.7428	0.8633	0.6083	0.7372	0.6678	0.6717	0.7006	0.5711	0.6475			
	F_{IS}		0.219	0.866	0.8	0.763	0.39	0.085	0.546	0.342	0.179	0.397	0.515	0.66	0.222	0.396	0.205	0.441			
	HWE		0.1332	0.0000	0.0000	0.0000	0.0037	0.0546	0.0010	0.1016	0.0219	0.0117	0.0000	0.0002	0.1259	0.0030	1.0000				
SGD	A_v	5	2.997	2	4	4	5	4.997	5	3	10.99	3.997	7	6	5.997	6	4.9987				
	H_v		0.3333	0.5	0.4667	0.4	0.5333	0.5333	0.5667	0.5	0.5667	0.5667	0.5667	0.4667	0.7333	0.4667	0.5	0.5062			
	H_s		0.5367	0.5228	0.4911	0.5861	0.6344	0.755	0.665	0.71	0.8667	0.6072	0.7506	0.7661	0.7828	0.7472	0.7117	0.6693			
	F_{IS}		0.318	0.06	0.067	0.333	0.176	0.309	0.164	0.311	0.321	0.084	0.261	0.405	0.08	0.39	0.313	0.247			
	HWE		0.0490	0.4814	0.7270	0.0749	0.0315	0.4179	0.0324	0.1320	0.0598	0.1549	0.0225	0.0219	0.2162	0.0004	0.4485				
SKT	A_v	4	3	3	3.933	4	5	4.997	5	3.997	8.996	3.933	6	5	4.997	5.933	4	4.7366			
	H_v		0.4333	0.4667	0.5333	0.4	0.5667	0.5667	0.6	0.4333	0.6333	0.6	0.4	0.6333	0.5333	0.4	0.5208				
	H_s		0.61	0.5228	0.56	0.5061	0.6261	0.7117	0.6428	0.7428	0.8611	0.5428	0.7994	0.7178	0.6978	0.6789	0.6939	0.6579			
	F_{IS}		0.305	0.077	0.065	0.226	0.112	0.22	0.083	0.253	0.092	0.218	0.265	0.456	0.109	0.231	0.437	0.214			
	HWE		0.1549	0.7149	1.0000	0.4485	0.2881	0.0205	0.4789	0.4179	0.7234	0.5886	0.1259	0.0002	0.4540	0.0315	0.0563				
KWH	A_v	4.998	2	2	2.933	4	5	5	4	10	3.933	7	6	6	6	3	4.804				
	H_v		0.4333	0.3	0.3667	0.4	0.6333	0.5333	0	0.5333	0.5333	0.4333	0.5333	0.7333	0.5667	0.4667	0.4937				

Table 3. Continued

H_e	0.5778	0.4861	0.4861	0.4994	0.6633	0.6967	0.6661	0.700	0.6994	0.8511	0.5606	0.7911	0.6933	0.7789	0.7567	0.5844	0.6557
F_{IS}	0.25	0.367	0.262	0.215	0.062	0.25	0.414	0.254	0.253	0.349	0.184	0.466	0.247	0.075	0.267	0.218	0.2583
HWE	0.1341	0.0617	0.2559	0.2559	0.1549	0.0068	0.0205	0.0749	0.0017	0.0730	0.1320	0.0038	0.0749	1.0000	0.0563	0.2881	
A_e	5	2.999	3	3.933	4	4	5	6	4	8.996	4	5.997	6	5	6	4	4.87
H_o	0.4333	0.3667	0.4667	0.4333	0.4667	0.7333	0.6333	0.6667	0.4667	0.7	0.5667	0.6333	0.6333	0.6667	0.5333	0.3	0.5437
H_e	0.5939	0.5294	0.5228	0.5139	0.6322	0.69	0.7344	0.7333	0.7039	0.825	0.6417	0.7506	0.7828	0.7711	0.7378	0.6917	0.6784
F_{IS}	0.223	0.297	0.124	0.173	0.278	-0.046	0.154	0.108	0.352	0.168	0.134	0.161	0.207	0.152	0.293	0.578	0.209
HWE	0.2747	0.0597	0.7060	0.4485	0.0315	0.7109	0.4467	0.1572	0.1320	1.0000	0.4789	0.2559	0.1284	0.3784	0.0563	0.0038	

Statistically significant values are marked with asterisks. * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$; ns = non-significant

Here (A_e) designated for allelic richness, (H_o) observed heterozygosity, (H_e) expected heterozygosity, (F_{IS}) inbreeding index and (HWE) deviation from Hardy-Weinberg equilibrium

expected heterozygosity ranged from 0.6475 (TTS population) to maximum 0.7055 (LHR population).

Inbreeding coefficient (F_{IS}) for each hatchery population of *H. molitrix* was calculated across all the loci. The mean values of F_{IS} ranged from 0.174 in the LHR population to a highest of 0.441 for the TTS population indicating heterozygote deficiency in all populations. The χ^2 values (for loci) clearly indicated the significant deviation from Hardy-Weinberg equilibrium (*HWE*) in all the hatchery populations of *H. molitrix*. Out of 128 tests, 38 tests showed significant deviation from *HWE* in the hatchery populations of *H. molitrix*. Non-significant deviations from *HWE* were observed at Locus BL138 in all the hatchery populations.

Across all the microsatellite loci, the pairwise genetic differentiation (F_{ST}) among hatchery stocks of *H. molitrix* was analysed by FSTAT. The study revealed statistically significant ($P < 0.01$) differentiation among most of the population pairs which reflects that most of hatchery populations are genetically non-homogenous (Table 4). The values of F_{ST} among the population pairs ranged from 0.0014 (FSD-GWH) to 0.0297 (TTS-KWH). The results of AMOVA indicated that most of the variation percentage lie within the individuals (39.54%). The variation between populations and between samples within population was measured as 27.31% and 33.16%, respectively (Table 5).

The UPGMA dendrogram was constructed based on Nei's unbiased genetic distance among all the examined hatchery populations of *H. molitrix* by using the software TFGA. The dendrogram showed hatchery populations of *H. molitrix* in two clusters/groups (Figure 2).

Discussion

In Pakistan, many studies have been conducted on nutrition, toxicology, physiology, growth performance and culture of Chinese carps but genetic issues pertaining to the subject species remained unattended. The available data on population genetics of *H. molitrix* is overwhelmingly reported from China and few studies from other regions. The study on genetic structure of *H. molitrix* was not reported from Pakistan so far. Hence, this is the very first information describing the population structure of *H. molitrix* in hatchery conditions of Punjab, Pakistan. Lack of expertise and infrastructure for undertaking genetic studies are the obvious reasons for the scarcity of work on genetic status of Chinese carp and even native fish species (King, Eackles, & Chapman, 2011).

The present study was conducted to underpin the genetic status of *H. molitrix* in selected hatchery populations of the Punjab province. The methodology used was fairly efficient for the extraction of genomic DNA from muscle tissues, its qualitative and quantitative analysis and subsequent PCR-based genotyping of the fish individuals at target microsatellite loci. The species-specific microsatellite markers employed for genotyping of the sample individuals proved to be robust tools for genetic monitoring of hatchery populations.

The present study revealed that the genetic diversity in hatchery stocks was found at low-to-moderate scale. The probable reason for the limited allelic diversity in hatchery populations of *H. molitrix* is the invasiveness because this species is not indigenous to Pakistan and was introduced in the

Table 4. Pairwise F_{ST} (above diagonal) and geographical distance (Km) (below diagonal) in the hatchery populations of *H. molitrix*

Populations	LHR	SHK	GWH	TTS	FSD	SGD	SKT	KWH
LHR	-	0.0114	0.0076*	0.0109	0.0249	0.004*	0.0132	0.0187
SHK	56	-	0.0062*	0.0120	0.0078*	0.002*	0.0138	0.008*
GWH	72	54	-	0.0128	0.0014*	0.002*	0.009*	0.002*
TTS	265	213	250	-	0.0254	0.006*	0.0224	0.0297
FSD	184	133	183	82	-	0.003*	0.002*	0.006*
SGD	189	139	175	160	92	-	0.009*	0.003*
SKT	143	138	52	304	223	229	-	0.008*
KWH	298	310	369	415	203	245	415	-

* = Significant statistically at $P < 0.01$

Table 5. Hierarchical AMOVA in the hatchery populations of *H. molitrix*

Source of variance	df	MSS	Variance	% variation
Between populations	7	18.66	0.25	27.31
Between samples within Populations	232	6.43	1.23	33.16
Within individuals	240	4.21	3.32	39.54
Total	479	29.3	4.8	100




Figure 1. UPGMA dendrogram based genetic relatedness between the hatchery populations of *H. Molitrix*.

country during the recent past. During 1960s, the Chinese carps were introduced to Pakistan primarily in order to control the menace of vegetation in aquatic systems. Due to close ecological conditions, *H. molitrix* adapted well in Pakistan (DoF, 2005). Moreover, the technique of hypophysation for successful artificial spawning carps led to rapid expansion of Chinese carps in natural water bodies (Bondad-Reantaso, 2007). Although, there is no data available over the number of individuals of Chinese carps imported from China, the evidences suggest that the number was very small (in thousands). Secondly, the loss of allelic diversity due to mass production of fish seed with limited number of brooders in hatcheries is a common phenomenon in the Asian region (Aung, Nguyen, Supawadee, & Kamornrat, 2010). Microsatellite markers based studies on genetic structure of *H. molitrix* revealed that conventional breeding practices in hatcheries has resulted in allelic loss mainly due to low effective population size and other poor brood stock management practices (Ji, Gu, Mao, Zhu, & Sun, 2010; Ovenden *et al.*, 2016).

Low to moderate level of allelic diversity (2-11) was observed in hatchery populations of *H. molitrix* in the present study. The reason of limited allelic diversity is directly influenced by bottleneck and founder effect because the hatchery managers kept limited number of brooders in artificial propagation for commercial reasons (Wang *et al.*, 2008). Related findings about low to moderate level of the allelic diversity were reported by Liao *et al.* (2007) and allelic diversity ranged from 2 to 16 while developing primer for cross amplification of Silver Carp and Bighead Carp. This value can also be in the range of 2-7 in wild Silver Carp as described by Cheng, Liu, Yu, and Tong (2008).

The mean values of observed heterozygosity (H_o) varied from 0.3770 to 0.5874 in the hatchery populations of *H. molitrix*, in this study. The lower level of H_o in the hatchery stocks might correspond to the inbreeding and negative selection during hatchery propagation programs (Zhou, Zhao, Bi, & Hu, 2011). Secondly, allelic diversity plays a key role in determining the level of heterozygosity that was found limited in this study. The findings of this study

about H_o are in accordance to the inferences reported by Pang, Yu, and Tong (2015).

who described higher level of H_o in natural populations of *H. molitrix* from China. In this study, the mean values of expected heterozygosity (H_e) were recorded as relatively higher in comparison to H_o that is in accordance with the findings of Zhang, XU, Liu, Duan, and Shi (2012). Coscia, Chopelet, Waples, Mann, and Mariani (2016) described that the value of H_e directly correlate with the effective size (N_e) of population especially in case of wild fish. While, in hatchery fish stock the relatively H_e implied to large number of breeding individuals. Therefore, it is necessary to maintain sufficient census number in relation to their N_e for the sustainable fish hatchery stock (Hare *et al.*, 2011). Generally, real number of brooders that contributed in succeeding generations remain unknown due to large number of effective individuals utilized in hatcheries and short domestication history.

The average values of inbreeding coefficient (F_{IS}) ranged from 0.174 to 0.441 in the hatchery populations of *H. molitrix*. The impact of demographic processes in isolated small populations results in loss of genetic variability which increases the incidence of inbreeding depression which may lead to diminish population's existence probability Hayer *et al.* (2014).

The findings of the present study about F_{IS} are different from those reported by Guo, Yu, and Tong (2013) for *H. molitrix* due to inadvertent hybridization (that is an unnoticed phenomenon) in the hatcheries of Punjab that is adversely affecting the genetic integrity. The same has been reported by Sabbir, Khan, Sultana, Rahi, and Shah (2017) over the incidence of hybridization and introgression between the indigenous carps in the hatcheries of Bangladesh. The occurrence of hybridization in hatchery stocks is widespread than in the wild environments mainly caused by poor broodstock management practices in Asia.

The microsatellite loci screened in this study revealed low level of population differentiation among the hatchery populations *H. molitrix*. Generally, genetic differentiation is considered as

very weak with F_{ST} ranging from 0.00 to 0.05, medium with 0.05-0.15 and large genetic differentiation with 0.15-0.25 and higher for intra-specific groups (Wright, 1951). The range of F_{ST} values among all the hatchery population pairs of *H. molitrix* were observed from 0.0014 (FSD-GWH) to 0.0297 (TTS-KWH) and 0.0021 (FSD-KWH) to 0.0175 (TTS-GWH), respectively. Low gene flow and a large geographical distance (415 Km) between SKT-KWH population pair may have contributed to higher population differentiation. Conventional mixing of gene pools by human interventions may be the reason of lowest value of F_{ST} between the FSD-GWH population pair even with a larger geographical distance of approximately 183 Km. Low gene flow and a widespread geographical distance between population pair may be the possible reasons to higher population's differentiation (Li et al., 2011).

Over most of the loci in *H. molitrix* in hatchery populations, the χ^2 values clearly indicated the significant deviation from *HWE*. The deficiency in heterozygotes can be caused by inbreeding, presence of null alleles and admixture of distinct populations (Guo et al., 2013). Comparable findings have been reported for a wide range of freshwater fish species by Keller et al. (2013).

Analysis of molecular variance (AMOVA) is used to delineate partitioning of genetic divergence within and among groups. In the present study, the results of AMOVA revealed that most of the variation existed at the inter-population level, showing significant genetic structuring in hatchery populations of *H. molitrix*. These results are in agreement with Feng, Yu, Fu, He, and Tong (2014).

Based on Nei's unbiased genetic distance among all the examined hatchery populations of Silver Carp, the UPGMA dendrogram was constructed to explore the genetic related among them. Mostly, the dendrogram clustered the populations onto two major branches showing their genetic relatedness partially following their geographic distribution. The populations lying in the same cluster with limited genetic distance despite larger geographic distance reveal the brood stock management structure of fisheries department in the province. This owes to lack of plan for genetic management of fish genetic resources in Pakistan (Haque & Hoq, 2016). For perspective studies, it is suggested to develop novel markers for the local populations. A holistic study with comprehensive sampling and employment of a combined panel of different marker systems can better explore the underlying genetic issues. Future research on hybridization and genetic introgression between Chinese carps and related taxa can add to the existing understanding over the genetic structure of subject species.

References

Askari, G., Shabani, A., & Miandare, H. K. (2013).

- Application of molecular markers in fisheries and aquaculture. *Journal of Animal Science*, 2(4), 82-88.
- Aung, O., Nguyen, T. T. T., Poompuang, S., & Kamornrat, W. (2010). Microsatellite DNA markers revealed genetic population structure among captive stocks and wild populations of mrigal, *Cirrhinus cirrhosus* in Myanmar. *Aquaculture*, 299(1), 37-43. <http://doi.org/10.1016/j.aquaculture.2009.12.010>
- Avise, J.C. (2010). Perspective: Conservation genetics enters the genomics era. *Conservation Genetics*, 11, 665-669. <http://dx.doi.org/10.1007/s10592-009-0006-y>
- Bondad-Reantaso, M. (2007). Assessment of freshwater fish seed resources for sustainable aquaculture. FAO Fisheries Technical Paper. No. 501. Rome, FAO. 2007. 628p.
- Cheng, L., Liu, L., Yu, X., & Tong, J. (2008). Sixteen polymorphic microsatellites in Bighead Carp (*Aristichthys nobilis*) and cross-amplification in Silver Carp (*Hypophthalmichthys molitrix*). *Molecular Ecology Resource*, 8(3), 656-658. <http://dx.doi.org/10.1111/j1471-8286-02037-x>
- Coscia, I., Choquet, J., Waples, R.S., Mann, B.Q., & Mariani, S. (2016). Sex change and effective population size: Implications for population genetic studies in marine fish. *Heredity*, 117(4), 251-258. <http://dx.doi.org/10.1038/hdy.2016.50>
- De-Cara, M.A.R., Villanueva, B., Toro, M.A., & Fernandez, J. (2013). Using genomic tools to maintain diversity and fitness in conservation programmes. *Molecular Ecology*, 22(24), 6091-6099. <http://dx.doi.org/10.1111/mec.12560>
- Defaveri, J., Viitaniemi, H., Leder, E., & Merila, J. (2013). Characterizing genic and non-genic molecular markers: comparison of microsatellites and SNPs. *Molecular Ecology Resource*, 13(3), 377-392. <http://dx.doi.org/10.1111/1755-0998.12071>
- Diana, J.S. (2009). Aquaculture production and biodiversity conservation. *BioScience*, 59(1), 27-38. <http://dx.doi.org/10.1525/bio.2009.59.1.7>
- DoF. (2005). Fish and Fisheries in the Punjab - A Manual. Department of Fisheries, Punjab, Lahore.
- Ellergern, H., & Galtier, N. (2016). Determinants of genetic diversity. *Nature Reviews Genetics*, 17(7), 422-433. <http://dx.doi.org/10.1038/nrg.2016.58>
- Excoffier, L., Smouse, P.E., & Quattro, J.M. (1992). Analysis of molecular variance inferred from metric distances among DNA haplotypes: Application to human mitochondrial DNA restriction data. *Genetics*, 131(2), 479-491. PMID: PMC1205020.
- Feng, X., Yu, X., Fu, B., He, S., & Tong, J. (2014). Development of 159 transcript-associated microsatellite markers in Silver Carp (*Hypophthalmichthys molitrix*). *Conservation Genetics, Resources*, 6(1), 111-113. <http://dx.doi.org/10.1007/s12686-013-0017-3>
- Goudet, J. (2002). FSTAT Version 2.9.3.2, a program to estimate and test gene diversities and fixation indices. Institute of Ecology and Evolution, University of Lausanne, Switzerland.
- Green, M.R., & Sambrook, J. (2012). Molecular cloning: A laboratory manual. 4th Ed. Cold spring harbor laboratory press, New York, USA.
- Guo, W., Yu, X., & Tong, J. (2013). Development of 134 novel polynucleotide-repeat microsatellite markers in Silver Carp (*Hypophthalmichthys molitrix*). *Conservation Genetics Resource*, 5(2), 525-528.

- <http://dx.doi.org/10.1007/s12686-012-9843-y>
- Haque, N., & Hoq, T. (2016). Deterioration of genetic diversity: concern in hatchery populations of Catla (*Catla catla*, Hamilton 1822: Cypriniformes, Cyprinidae) of Sylhet district in Bangladesh. *International Journal of Current Microbiology and Applied Science*, 5(12), 171-181. <http://dx.doi.org/10.20546/ijcmas.2016.512.019>
- Hare, M.P., Nunney, L., Schwartz, M.K., Ruzzante, D.E., Burford, M., Waples, R.S., Ruegg, K., & Palstra, F. (2011). Understanding and estimating effective population size for practical application in marine species management. *Conservation Biology*, 25(3), 438-49. <http://dx.doi.org/10.1111/j.1523-1739.2010.01637.x>
- Hauser, L., & Seeb, J.E. (2008). Advances in molecular technology and their impact on fisheries genetics. *Fish and Fisheries*, 9(4), 473-486. <http://dx.doi.org/10.1111/j.1467-2979.2008.00306.x>
- Hayer, C.A., Breeggemann, J.J., Klumb, R.A., Graeb, B.D.S., & Bertrand K.N. (2014). Population characteristics of Bighead and Silver Carp on the northwestern front of their North American invasion. *Aquatic Invasions*, 9(3), 289-303. <http://dx.doi.org/10.3391/ai.2014.9.3.05>
- Ji, C.H., Gu, J.J., Mao, R.X., Zhu, X.P., and Sun, X.W. (2009). Analysis of genetic diversity among wild Silver Carp (*Hypophthalmichthys molitrix*) populations in the Yangtze, Heilongjiang and Pearl Rivers using microsatellite markers. *Journal of Fisheries of China*, 33(3), 364-371. [http://dx.doi.org/1000-0615\(2009\)03-0364-08](http://dx.doi.org/1000-0615(2009)03-0364-08)
- Keller, I., Wagner, C.E., Greuter, L., Mwaiko, S., Selz, O.M., Sivasundar, A., Wittwer, S., & Seehausen, O. (2013). Population genomic signatures of divergent adaptation, gene flow and hybrid speciation in the rapid radiation of Lake Victoria cichlid fishes. *Molecular Ecology*, 22(11), 2848-2863. <http://dx.doi.org/10.1111/mec.12083>
- Khan, A.M., Shakir, H.A., Khan, M.N., Abid, M., & Mirza, M.R. (2008). Ichthyofaunal survey of some freshwater reservoirs in Punjab. *Journal of Animal and Plant Science*, 18(4), 151-154.
- Khoo, G., Lim, T.M., & Phang, V.P.E. (2011). A review of PCR-based DNA fingerprinting using arbitrary primers in tropical ornamental fishes of South-East Asia. *Journal of Advanced Medical Research*, 1(2), 71-93.
- King, T.L., Eackles, M.S., & Chapman, D.C. (2011). Tools for assessing kinship, population structure, phylogeography and interspecific hybridization in Asian carps invasive to the Mississippi River, USA: Isolation and characterization of novel tetranucleotide microsatellite DNA loci in Silver Carp *Hypophthalmichthys molitrix*. *Conservation Genetics Resource*, 3(3), 397-401. <http://dx.doi.org/10.1007/s12686-010-9285-3>
- Li, S.F., Jia-Wei, X., Qi-Ling, Y., Cheng-Hui, W., Chapman, D.C., & Lu, G. (2011). Significant genetic differentiation between native and introduced Silver Carp (*Hypophthalmichthys molitrix*) inferred from mtDNA analysis. *Environmental Biology of Fishes*, 92(4), 503-511. <http://dx.doi.org/10.1007/s10641-011-9870-7>
- Liao, M., Yang, G, Wang, X., Wang, D., Zou, G., & Wei, Q. (2007). Development of microsatellite DNA markers of Silver Carp (*Hypophthalmichthys molitrix*) and their cross-species application in Bighead Carp (*Aristichthys nobilis*). *Molecular Ecology Notes*, 7(1), 95-99. <http://dx.doi.org/10.1111/j.1471-8286.2006.01542.x>
- Miller, M.P. (1997). Tools for population genetic analyses (TFPGA) v 1.3: A windows program for the analysis of allozyme and molecular genetic data. Flagstaff: Northern Arizona, University.
- Munilkumar, S., & Nandeesha, M.C. (2007). Aquaculture practices in Northeast India: current status and future directions. *Fish Physiology and Biochemistry*, 33(4), 399-412. <http://dx.doi.org/10.1007/s10695-007-9163-4>
- Nei, M. (1972). Genetic distance between populations. *The American Naturalist*, 106(949), 283-292.
- Ovenden, R., Leigh, G.M., Blower, D.C., Jones, A.T., Moore, A., Bustamante, C., Buckworth, R.C., Bennett, M.B., & Dudgeon, C.L. (2016). Can estimates of genetic effective population size contribute to fisheries stock assessments? *Journal of Fish Biology*, 89(6), 2505-2518. <http://dx.doi.org/10.1111/jfb.13129>
- Pang, M.X., Yu, X.M., Tong, J.G. (2015). The microsatellite analysis of genetic diversity of five Silver Carp populations in the three Gorges reservoir of the Yangtze River. *Acta Hydrobiologica Sinica*, 39(5), 869-876. <http://dx.doi.org/10.7541/2015.115>
- Paukert, C.P., Glazer, B.A., Hansen, G.J.A., Irwin, B.J., Jacobson, P.C., Kershner, J.L., Shuter, B.J., Whitney, J.E., & Lynch, A.J. (2016). Adapting inland fisheries management to a changing climate. *Fisheries*, 41(7), 374-384. <http://dx.doi.org/10.1080/03632415.2016.1185009>
- Pei, Y.L., Costumbrado, J., Hsu, C., & kim, Y.H. (2012). Agarose gel electrophoresis for the separation of DNA fragments. *Journal of Visualized Experiments*, 62: 3923. <http://dx.doi.org/10.3791/3923>
- Qu, L., Li, X., Wu, G., & Yang, N. (2005). Efficient and sensitive method of DNA silver staining in polyacrylamide gels. *Electrophoresis*, 26(1), 99-101. <http://dx.doi.org/10.1002/elps.200406177>
- Rice, W. R. (1989). Analysing tables of statistical tests. *Evolution*, 43(1), 223-225. <http://dx.doi.org/10.1111/j.1558-5646.1989.tb04220.x>
- Sabbir, W., Khan, Md. N., Sultana, S., Rahi, Md. L., & Shah, Md. S. (2017). Production of heterotic hybrid in Rohu (*Labeo rohita*) by crossing the riverine and hatchery strains. *International Journal of Innovation Sciences & Research*, 6 (2), 982-986.
- Van Oosterhout, C., Hutchinson, W.F., Wills D.P.M., & Shipley, P. (2004). Micro-checker: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes*, 4: 535-538. <http://dx.doi.org/10.1111/j.1471-8286.2004.00684.x>
- Van Overzee, H.M.J., & Rijnsdorp, A.D. (2015). Effects of fishing during the spawning period: Implications for sustainable management. *Reviews in Fish Biology and Fisheries*, 25(1), 65-83. <http://dx.doi.org/10.1007/s11160-014-9370-x>
- Wang, C.Z., Liang, H.W., Zou, G.W., Luo, X.Z., Li, Z., Tian, H., & Hu, G.F. (2008). Genetic variation analysis of two Silver Carp populations in the middle and upper Yangtze River by microsatellite. *Hereditas (Beijing)*, 10(30), 1341-1348.

- <http://dx.doi.org/10.3724/SP.J.1005.2008.01341>.
- Weir, B.S., & Cockerham, C.C. 1984. Estimating F-statistics for the analysis of population structure. *Evolution*, 38(6), 1358-1370. <http://dx.doi.org/10.2307/2408641>
- Wright, S. (1951). The genetical structure of populations. *Annals of Eugenics*, 15(4), 323-354. <http://dx.doi.org/PMID,24540312>
- Xu, K., Duan, W., Xiao, J., Tao, M., Zhang, C., Liu, Y., & Liu, S.J. (2015). Development and application of biological technologies in fish genetic breeding. *Science China Life Science*, 58(3), 187-201. <http://dx.doi.org/10.1007/s11427-015-4798-3>
- Yeh, F.C., Yang, R.C., Boyle, T.B.J., Ye, Z.H., & Mao, J.X. (1999). POPGENE, The user-friendly shareware for population genetic analysis, Molecular Biology Biotechnology Centre; University of Alberta Canada
- Yue, G.H., & Orban, L. (2005). A simple and affordable method for high-throughput DNA extraction from animal tissues for polymerase chain reaction. *Electrophoresis*, 26(16), 3081-3083. <http://dx.doi.org/10.1002/elps.200410411>
- Zhang, M.Y., XU, D.P., Liu, K., Duan, J.R., & Shi, W.G. (2012). Analysis of genetic diversity among Silver the artificially released populations of carps in the lower reach of Yangtze River with microsatellite markers. *Acta Agriculturae Universitatis Jiangxiensis*, 34(1), 41-146. <http://dx.doi.org/10.3969/j.issn.1000-2286.2012.01.026>
- Zhou, G., Zhao, X., Bi, Y., & Hu, Z. (2011). Effects of Silver Carp (*Hypophthalmichthys molitrix*) on spring phytoplankton community structure of three Gorges reservoir (China): results from an enclosure experiment. *Journal of Limnology*, 70(1), 26-32. <http://dx.doi.org/10.3274/JL11-70-1-05>