

Microalgae and Cyanobacteria, a Promising Source of Antimicrobial Molecules Against Aquatic Pathogen

Carlos Torres-Bayona^{1,*} , Jorge Leyva Rojas¹ , Ronield Fernandez¹ , Martha Prieto-Guevara² , Alexander Pulido¹ , Ignacio Moreno-Garrido^{3,*} 

¹Simón Bolívar University, Faculty of Basic and Biomedical Science, Microbiology Department, Barranquilla/Atlantico, 080002, Colombia.

²Córdoba University, Faculty of Veterinary Medicine and Zootechnics, Montería /Cordoba, 230002, Colombia.

³Institute of Marine Sciences of Andalucía -CSIC-, Department of Ecology and Coastal Management. (CSIC). Campus Universitario Río San Pedro, 11510 Puerto Real, Cádiz, Spain

How to cite

Torres-Bayona, C., Rojas, J.L., Fernandes, R., Prieto-Guevara, M., Pulido, A., Moreno-Garrido, I. (2023). Microalgae and Cyanobacteria, a Promising Source of Antimicrobial Molecules Against Aquatic Pathogen *Turkish Journal of Fisheries and Aquatic Sciences*, 23(2), TRJFAS21184. <https://doi.org/10.4194/TRJFAS21184>

Article History

Received 02 March 2022

Accepted 07 September 2022

First Online 23 September 2022

Corresponding Author

Tel.: +34686661373

E-mail: ignacio.moreno@icman.csic

Keywords

Antibacterial activity

Microalgae and cyanobacteria

Methanol extracts

Vibriosis

Antibiotic resistance

Abstract

Genus *Vibrio* is involved in common pathologies of aquaculture fish species, being responsible for significant economic losses for that industrial activity. Microalgae and cyanobacteria have demonstrated to count on bioactive compounds able to diminish mortality and morbidity by their use as functional food supply or by addition of those bioactive compounds previously purified. The main goal of this work has been to evaluate the anti-microbial effect (growth of bacterial population measured as changes in optical density) of 25 microalgae extracts on the bacterial strain *Vibrio anguillarum* CECT 522T. Stock microalgae extracts were obtained from lyophilized biomass treated with methanol 99.9%. A total of thirteen extracts exhibited antibacterial activity. The highest activity corresponded to *Monochrysis lutheri*, followed by *Hemiselmis cyclopea*, *Porphyridium cruentum*, *Tetraselmis rubens*, *Cryptomonas* sp, *Navicula* sp. The anti-bacterial activity was not related to a taxonomic group, indicating species-specific or even strain-specific activity.

Introduction

Aquaculture is an emerging industrial activity of great importance for the economic development of countries (Arunkumar *et al.*, 2020). Aquaculture production is currently estimated to supply almost half of the fish consumed by humans worldwide (FAO, 2020).

The high demand for aquaculture products, on many occasions, leads to an overpopulation of specimens per unit of production (Bouwmeester *et al.*, 2021), allowing the appearance of infectious outbreaks that cause large economic losses, thus becoming, one of the main limiting factors for the development of the aquaculture sector (Ina-Salwany *et al.*, 2019). Economic

losses due to morbidity, mortality, and investment costs in treatments for the control and prevention of infectious outbreaks using drugs are so high that they may even lead to the total collapse of some facilities (Bermúdez Almada *et al.*, 2017). In China economic losses are around 15-20% due to disease (Sudheesh & Cain, 2017). In Chile operating costs for pharmaceutical products are around the US \$700 million per year (Flores *et al.*, 2020).

Among the important pathologies of great economic and epidemiological implication in aquaculture is vibriosis. This bacterial disease is caused by some species of the genera *Vibrio* and *Photobacterium* (Istiqomah *et al.*, 2020). The

significance of these genera in world aquaculture lies in the wide range of species affected, including fish, crustaceans, mollusks, corals, and rotifers of any productive level, strong commercial system or small-scale rural aquaculture farms (Mohamad *et al.*, 2019).

One of the main etiological agents related to vibriosis is the species *Vibrio anguillarum*; This gram-negative pathogenic bacterium has been described as capable of causing hemorrhagic septicemia in more than 50 species of freshwater and marine fish, among which are species of great economic importance such salmon (*Salmo salar*), turbot (*Psetta maximum*), sea bass (*Dicentrarchus labrax*), sea bream (*Sparus aurata*), cod (*Gadus morhua*), eel (*Anguilla anguilla*) or ayu (*Plecoglossus altivelis*), as well as some bivalves and crustaceans (Behringer *et al.*, 2020; Derome *et al.*, 2016).

It is worth mentioning that, along with bacterial diseases and the great economic losses derived from the control or prophylactic treatment of the different pathologies associated with aquaculture, there is underlying collateral damage that derives from the excessive use of antimicrobial drugs: the appearance of multi-resistant bacterial strains with resistance genes which could be horizontally transferred to non-target organisms, including pathogens from terrestrial animals, including man (Santos & Ramos, 2018).

Currently, there is a pressing need to discover novel broad-spectrum antimicrobial active principles, whose molecules make it possible to evade current biochemical mechanisms of antimicrobial resistance, thus reducing the impact of bacterial diseases, and the increased antimicrobial resistance (Igarashi *et al.*, 2018). For decades, the main sources for obtaining antimicrobial molecules have been bacteria, fungi, and actinomycetes (Hutchings *et al.*, 2019); however, marine organisms have shown to be a promising natural bank of molecules with broad biological activity including antimicrobial activity (Nweze *et al.*, 2020).

Microalgae (*sensu lato*), represents a vast group of multi-diverse microorganisms, which includes cyanobacteria and dinoflagellates. It is estimated that there are at least 200,000 species, distributed in almost all aquatic and humid ecosystems globally (Norton *et al.*, 1996; Sydney *et al.*, 2019).

Although only a low percentage of microalgae species have been characterized (Ameen *et al.*, 2021), the literature describes multiple active compounds with diverse biological activity, of which many of these molecules are related to the taxonomic groups of this study. Among the main compounds with antimicrobial activity from microalgae, it could be mentioned: organic acids, fatty acids, unsaturated hydroxylated fatty acids, glycolipids, steroids, phenols, terpenoids, toxins (such as bacterial toxin lecithinase), phycobiliproteins, exopolysaccharides (EPS) and antimicrobial peptides (AMP) (Hayes *et al.*, 2017; Nájera-Arce *et al.*, 2018; Pina-Pérez *et al.*, 2017).

Microalgae have a wide adaptive metabolic plasticity ability to adapt their biochemical responses to their physiological state (Falaise *et al.*, 2016). This quality allows through the manipulation of physicochemical factors such as nutritional stress (Lin *et al.*, 2018; Lovio *et al.*, 2019) temperature, irradiance (Gonçalves *et al.*, 2019) pH, salinity (Jaiswal *et al.*, 2020; Musa *et al.*, 2019), the expression of different groups of metabolites of industrial interest can be induced (Elisabeth *et al.*, 2021).

Some studies demonstrated the bactericidal and fungicidal effect of microalgae extracts on pathogens of epidemiologic interests, such as *Staphylococcus aureus*, *E. coli* and *P. aeruginosa*, as well as on different species from the genus *Vibrio* (Dewi *et al.*, 2018; Jena & Subudhi, 2019; Jha *et al.*, 2017; Ruiz *et al.*, 2019; Schuelter *et al.*, 2019). On the other hand, there is a great interest in the food industry, especially in seafood products, due to the bioactives of microalgae, where the preservative and quality improvement effect of seafood has been demonstrated (Tavakoli *et al.*, 2021).

This work involves an important amount of microalgae strains (25) from different taxonomic classes, to be assayed as a possible source of new antimicrobial molecules. Antibacterial activity of different microalgal strains would be exploited to understanding the mechanism of antimicrobial action of some molecules and them optimal use in fish protection against bacterial pathogens. Additionally, this protection would not be related to synthetic antibiotics which can provoke, a) resistance of bacterial strains, and b) environmental problems due to antibiotic activity on non-target organisms. Antibacterial activity of any of those strains has not assayed before.

Materials and Methods

Microalgae and Cyanobacteria Cultures and Extracts

Microalgae strains were obtained from the Marine Microalgae Culture Collection (MMCC) at the Institute of Marine Sciences of Andalusia (CSIC), and the Microalgae Collection from the INTERREG Project ALGARED+ (Table 1). Algae were cultured in marine well water, enriched with Guillard's f/2 medium (Guillard & Ryther, 1962), and $500 \mu\text{g} \times \text{L}^{-1}$ silicate in the case of diatoms. Cells were grown in 15 L cylindrical methacrylate columns, aerated with $0,2 \mu\text{m}$ filtered air. Cells were cultured at $20 \pm 1 \text{ }^\circ\text{C}$, under continuous white light (around $30 \mu\text{mol quanta} \times \text{m}^2\text{s}^{-1}$) till enough biomass was reached. Then, cultures were centrifuged in a Westfalia suspension solid separator continuous flow centrifuge. Biomass obtained was washed with ammonium formiate (Sigma-Aldrich) (0.9%) to remove salt and then frozen to -80°C . After this, samples were lyophilized (Virtis Advantage Wizard 2.0, from Hucoa Erlöss).

The dry biomass of each microalgae strain was diluted in $10 \text{ ml} \cdot \text{g}^{-1}$ of 99.9% methanol, (Merck, HPLC

grade), then, was disrupted Sonic (Hielscher UP200S) at 240 W for 20 s on ice.

After this treatment, samples were stored for 24h at 4°C and then submitted to the same sonic disruption again and re-stored at 4°C for another 24 h. After this step, the samples were centrifuged and the supernatant was filtered with a syringe filter (nylon membrane pore size 0.22 µm, SinerLab). Then, the crude extracts were dried to 35°C in a rotary evaporator and then stored at -80°C till re-suspended in appropriate volumes of DMSO just before the assays assayed. Controls were treated with the same DMSO volumes, after checking the DMSO concentration used did not interfere with bacterial growth.

Bacterial Culture and Inhibitory Activity Bioassay.

The *Vibrio anguillarum* strain (CECT 522T) was supplied by the Biochemical and Molecular Biology Laboratory of the Chemical Department at the University of Huelva (Spain). The inoculum was stored at -80°C in marine agar and glycerol. Before experiments, the *V. anguillarum* strain was quickly thawed, re-seeded in agar Mueller Hinton (MH) (OXOID) prepared with filtered seawater and incubated at 26°C for 24 h, and then inoculated in MH broth and also incubated at 26°C for 24 h. Then, culture was adjusted to a MacFarland 0.5 optical patron, measuring optic density to adjust 0.08-0.1 DO at 595 nm.

Bioassays were carried out in 96 multi-well plates (NunClon) filled with 180 µL of optically adjusted *V. anguillarum* culture plus 20 µL of microalgae extracts diluted in 2% DMSO. To increase the possibility of observing any antimicrobial effect present in the extracts, the experiment was carried out by testing the maximum concentration obtained from each microalgae

respectively during the extraction process with 99.9% methanol.

Experiments were carried out with five replicate per microalgae extract, and the control was inoculated with 20 µL of 2% DMSO. Readings for OD were performed at time 0 and 24 h in a Multi-plate Spectrophotometer reader TECAN Genios 2000 (measuring wavelength 595 nm, band width 10 nm, reference wavelength 570 nm, number of flashes 25, no settling time. The incubation experimental time was 24h.

Data Analysis

Differences in *V. anguillarum* growth in response to the microalgae alcoholic extracts inoculation were analyzed with the Kruskal-Wallis test ($P < 0.05$) using the IBM SPSS Statistics for Windows, Version 22.0 (Armonk, NY. USA).

Results

Microalgae strains and their taxonomic group, as well as the final concentration of the crude extract used in the bioassay can be seen in Table 1.

The results of this study showed that thirteen methanolic extracts showed statistical differences related to the growth of *V. anguillarum* after 24 h, measuring such divergences in DO at 595 nm. Likewise, within these significances, two groups can be differentiated.

The first group showed high significant ($P < 0.001$) statistical differences with controls: *Hemiselmis Cyclopea*, *Porphyridium cruentum*, *Tetraselmis rubens*, *Cryptomonas sp.*, *Navicula sp.* and *Monochrysis lutheri*. This last strain showed the highest significant

Table 1. Microalgae strains and their taxonomic group. Final concentration of the crude extract used in the bioassay

Species (or strain)	Taxonomic group	Final extract concentration(mg mL ⁻¹)
<i>Ostreopsis cf. ovata</i>	Dinophyta	0.237
<i>Chaetoceros gracilis</i>	Bacillariophyta	0.149
<i>Gyrodinium dorsum</i>	Dinophyta	0.212
<i>Tetraselmis striata</i>	Prasinophyta	0.199
<i>Nannochloropsis sp</i>	Eustigmatophyta	0.240
<i>Chlorococcum sp</i>	Chlorophyta	0.153
<i>Amphora sp.</i>	Bacillariophyta	0.602
<i>Synechococcus sp</i>	Cyanophyta	0.213
<i>Chroomonas salina</i>	Cryptophyta	0.137
<i>Chlamydomonas reinhardtii</i>	Chlorophyta	0.177
<i>Navicula sp</i>	Bacillariophyta	0.155
<i>Diacronema vlkianum</i>	Haptophyta	0.203
<i>Ochromonas sp</i>	Chrysophyta	0.240
<i>Monochrysis lutheri</i>	Chrysophyta	0.140
<i>Dunaliella salina</i>	Chlorophyta	0.161
<i>Anabaena sp</i>	Cyanophyta	0.172
<i>Cryptomonas sp</i>	Cryptophyta	0.146
<i>Porphyridium cruentum</i>	Rhodophyta	0.257
<i>Tetraselmis rubens</i>	Prasinophyta	0.244
<i>Hemiselmis Cyclopea</i>	Cryptophyta	0.316
<i>Picochlorum maculatum</i>	Trebouxiophyceae	0.149
<i>Isochrysis galbana</i>	Haptophyta	0.159
<i>Nannochloropsis gaditana</i>	Eustigmatophyta	0.192
<i>Picochlorum oklahomensis</i>	Trebouxiophyceae	0.306
<i>Phaeodactylum sp</i>	Bacillariophyta	2.940

differences with controls (approximately 17% more bacterial growth inhibition than others). A second group, showed significant ($P < 0.05$) statistical differences with controls: *Phaeodactylum* sp., *Picochlorum oklahomensis*, *Nannochloropsis gaditana*, *Ochromonas* sp., *Chlamydomonas reinhardtii*, *Gyrodinium dorsum*, and *Chaetoceros gracilis*. The other 12 microalgae extracts showed no significant inhibitory effect on bacterial growth or stimulate bacterial proliferation, as was observed with the strain *Dunaliella salina* (Figure 1).

Discussion

In this study, antibacterial activity on *V. anguillarum* for methanol extracts of 13 from the 25 microalgae assayed strains has been stated. Although the most effective extracts were obtained from organisms including the classes Chrysophyta, Rhodophyta, and Prasinophyta, other organisms from those classes did not show the same activity, pointing out a species-specific (perhaps a strain-specific) antibacterial capacity. The possibility of activation of this capacity when microalgae and cyanobacteria are previously exposed to the pathogen should be checked in further studies.

Literature about antibacterial activity of microalgae extracts is scarce. Duff *et al.*, (1966) demonstrated the inhibitory effect of extracts from *Monochrysis lutheri*, *Hemiselmis* sp., *Cryptomonas* sp. and *Tetraselmis* sp., on different bacterial strains, mainly from marine environments. Some active antibacterial

metabolites from *M. lutheri* correspond to hydrocarbons, terpenoids, phenols (Metting & Pyne, 1986), phospholipases (Antia & Bilinski, 1967), and some lipidic compounds such as digalactosildiacil glycerol (GL) and phosfatidilcoline (PL) (Khomova *et al.*, 1994). Bashir *et al.* (2018) already demonstrated the bacteriostatic capacity of methanolic extracts of *C. reinhardtii*, *P. purpureum* *P. lutheri* and *Isochrysis* sp. on bacteria of epidemiologic interest such as *M. luteus*, *P. aeruginosa* and *S. aureus*.

Used as food supplements, microalgae have demonstrated a protective role for cultured fishes. Thus, when *M. lutheri* was added to composing feds, juvenile turbot (*Scophthalmus maximus*) survival increased (Spektorova *et al.*, 1986). Cultures salmons (*Salmo salar*) also demonstrated higher resistance to infections due to *Vibrio* sp. when microalgae *T. suecica* was included in the diet. (Falaise *et al.*, 2016).

Nowadays, alternatives to traditional antibiotic treatments are considered due to environmental and epidemiological consequences derived from the use of those substances. Some proposals include the use of bacteriophages, probiotics, enzymes or antimicrobial peptides (AMPs) (Ghosh *et al.*, 2019; Ture *et al.*, 2022).

Even some studies of antimicrobial activity through synergistic treatments with plant extracts have proven to be a good starting point to motivate future studies with microalgae extracts; which could contribute to a lower environmental effect than other substances (Bartual *et al.*, 2014; Cheesman *et al.*, 2017).




Figure 1. Effect of microalgae and cyanobacteria alcoholic extract on *Vibrio anguillarum* growth after 24 h culture in multi-well plates. Control corresponds to *Vibrio anguillarum* inoculate with 20 μ L of 2% DMSO. Statistical significance according to the Kruskal-Wallis test **($P < 0.001$); *($P < 0.05$) in the inhibition of bacterial growth in comparison with the control is signed with an asterisk.

Most of the used strains in this study were isolated from coastal environments. Those biocenosis are quite active in physic-chemical and biological terms (Rojas Jirón *et al.*, 2019), which could imply a better capacity of organisms inhabiting those biotopes to combat microbe pressure (Bartual *et al.*, 2020; Fuentes *et al.*, 2016; Salcedo & Coello, 2018). This is also an interesting point to be checked in order to test if strains of the same species belonging to ecosystems with different grades of physical-chemical (or even biological) pressure show different antibacterial activity.

Extraction with methanol permits the integration of a wide range of polar and non-polar compounds present in the microalgae and cyanobacterial biomass, which is an excellent first approach for a vast number of taxa. However, to achieve a greater range of extraction of compounds, it would be convenient to carry out successive fractions with different solvents of extreme and intermediate polarities, in this way a greater range of biological activity could be obtained (Cepas *et al.*, 2019).

In addition, it is considered that the application of techniques such as super-critical fluids (Sánchez-Camargo *et al.*, 2017) or microwave-assisted extraction (Kapoor *et al.*, 2018) could be a needed improvement for this type of study.

Our results stated that within the group of extracts that does not show an inhibitory effect against *V. anguillarum*, there is a treatment (*D. salina*) that imply an OD of 14.4% above the growth threshold of the control. Faced with this behavior, some hypotheses may arise, within which we could support two of them. The first could be related to the molecular profile of the methanolic extracts of the microalgae *D. salina*. Some studies, such as that of Ambrico *et al.*, (2020) have characterized the composition of some extracts of this microalgae, using solvents such as Methanol: chloroform, Hexane and ethanol. Within the main components of the extracts can be found, proteins (probably including free amino acids), carbohydrates, fatty acids, among others. This nutritional cocktail could favor the growth of *V. anguillarum*, in the event that there is no inhibitor agent, or if the latter is found in very low concentrations within the extracts.

On the other hand, it has been shown that some substances, with a certain degree of toxicity for some organisms, can exert a stimulating effect on growth when they are present at very low concentrations. This phenomenon it is known as Chemical Hormesis (in the case of chemical agents) and was reviewed by Calabrese and Baldwin, (2000); It is currently the subject of rigorous study to understand in more detail this effect on different organisms and toxic agents (Agathokleous *et al.*, 2021).

Due to the above, it could be presented that, some organic molecules that register antibacterial activity present in the extracts of *D. salina*, for example those described by Cakmak *et al.*, (2014) and Little *et al.*, (2021), could be found in very low concentrations, and

this can exert a stimulating bacterial proliferation, effect on the development of *V. anguillarum* as observed in our experiment.

Ethical Statement

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper

Funding Information

This research was funded by the Project INTERREG VA POCTEP-055_ALGARED_PLUS5_E (European union)., Asociación Universitaria Iberoamericana de Postgrado (AUIP) funded stays of Mr. Carlos Torres-Bayona in the ICMAN (CSIC).

Author Contribution

All authors have read and agree to the published version of the manuscript. Conceptualization, C.T.-B and I.M.-G.; methodology, C.T.-B, R.F and A.P.; software, C.T.-B and J.A.L.-R.; validation, C.T.-B, I.M.-G and J.A.L.-R.; formal analysis, C.T.-B, I.M.-G and J.A.L.-R.; investigation, C.T.-B, I.M.-G and.; data curation, C.T.-B, and J.A.L.; writing—original draft preparation, C.T.B, I.M-G and J.A.L.-R.; writing—review and editing, C.T.B, I.M-G, M.P-G and J.A.L.-R; supervision, I.M-G.; funding acquisition, I.M-G.

Conflict of Interest

The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Acknowledgements

Authors thanks Dr Rosa León Bañares, from the University of Huelva (Spain), for providing *V. anguillarum* strain. The authors also thank Dr Isidro Gonzales Collado for the support in experimentation at the Department of Organic Chemistry of the University of Cádiz (Spain).

References

- Ameen, F., AlNadhari, S., & Al-Homaidan, A.A. (2021). Marine microorganisms as an untapped source of bioactive compounds. *Saudi Journal of Biological Sciences*, 28(1), 224–231. <https://doi.org/10.1016/j.sjbs.2020.09.052>
- Antia, N.J., & Bilinski, E. (1967). A Bacterial-Toxin Type of Phospholipase (Lecithinase C) in a Marine Phytoplanktonic Chrysomonad. *Journal of the Fisheries Research Board of Canada*, 24(1), 201–204. <https://doi.org/10.1139/f67-014>

- Arunkumar, M., LewisOscar, F., Thajuddin, N., Pugazhendhi, A., & Nithya, C. (2020). In vitro and in vivo biofilm forming *Vibrio* spp: A significant threat in aquaculture. *Process Biochemistry*, 94, 213–223. <https://doi.org/10.1016/j.procbio.2020.04.029>
- Ambrico, A., Trupo, M., Magarelli, R., Balducchi, R., Ferraro, A., Hristoforou, E., ... & Molino, A. (2020). Effectiveness of *Dunaliella salina* extracts against *Bacillus subtilis* and bacterial plant pathogens. *Pathogens*, 9(8), 613. <https://doi.org/10.3390/pathogens9080613>
- Agathokleous, E., Wang, Q., Iavicoli, I., & Calabrese, E.J. (2021). The relevance of hormesis at higher levels of biological organization: hormesis in microorganisms. *Current Opinion in Toxicology*. <https://doi.org/10.1016/j.cotox.2021.11.001>
- Bartual, A., Arandia-Gorostidi, N., Cózar, A., Morillo-García, S., Ortega, M., Vidal, M., Cabello, A., González-Gordillo, J., & Echevarría, F. (2014). Polyunsaturated Aldehydes from Large Phytoplankton of the Atlantic Ocean Surface (42°N to 33°S). *Marine Drugs*, 12(2), 682–699. <https://doi.org/10.3390/md12020682>
- Bartual, A., Hernanz-Torrijos, M., Sala, I., Ortega, M.J., González-García, C., Bolado-Penagos, M., López-Urrutia, A., Romero-Martínez, L., Lubián, L.M., Bruno, M., Echevarría, F., & García, C.M. (2020). Types and Distribution of Bioactive Polyunsaturated Aldehydes in a Gradient from Mesotrophic to Oligotrophic Waters in the Alborán Sea (Western Mediterranean). *Marine Drugs*, 18(3), 159. <https://doi.org/10.3390/md18030159>
- Bashir, K.M.I., Lee, J.-H., Petermann, M.J., Shah, A.A., Jeong, S.-J., Kim, M.-S., Park, N.-G., & Cho, M.-G. (2018). Estimation of Antibacterial Properties of Chlorophyta, Rhodophyta and Haptophyta Microalgae Species. *Microbiology and Biotechnology Letters*, 46(3), 225–233. <https://doi.org/10.4014/mb.1802.02015>
- Behringer, D.C., Silliman, B.R., & Lafferty, K.D. (2020). *Marine Disease Ecology* (D.C. Behringer, B.R. Silliman, & K.D. Lafferty, Eds.; 1st ed.). Oxford University Press. <https://doi.org/10.1093/oso/9780198821632.001.0001>
- Bermúdez Almada, M.D.C., Espinosa Plascencia, A., Lara Espinoza, C.L., Rivera Domínguez, M., Astorga Cienfuegos, K.R., & Villalpando Canchola, E. (2017). Detección de vibrio mediante la amplificación de genes de patogenicidad en camarón *Litopenaeus vannamei* cultivado en un sistema tipo invernadero. *Investigación y Ciencia de La Universidad Autónoma de Aguascalientes*, 72, 20–29. <https://doi.org/10.33064/iycuaa201772218>
- Bouwmeester, M.M., Goedknecht, M.A., Poulin, R., & Thielges, D.W. (2021). Collateral diseases: Aquaculture impacts on wildlife infections. *Journal of Applied Ecology*, 58(3), 453–464. <https://doi.org/10.1111/1365-2664.13775>
- Cepas, V., López, Y., Gabasa, Y., Martins, C.B., Ferreira, J.D., Correia, M.J., Santos, L.M.A., Oliveira, F., Ramos, V., Reis, M., Castelo-Branco, R., Morais, J., Vasconcelos, V., Probert, I., Guilloud, E., Mehiri, M., & Soto, S.M. (2019). Inhibition of Bacterial and Fungal Biofilm Formation by 675 Extracts from Microalgae and Cyanobacteria. *Antibiotics*, 8(2), 77. <https://doi.org/10.3390/antibiotics8020077>
- Cheesman, M., Ilanko, A., Blonk, B., & Cock, I. (2017). Developing new antimicrobial therapies: Are synergistic combinations of plant extracts/compounds with conventional antibiotics the solution? *Pharmacognosy Reviews*, 11(22), 57. https://doi.org/10.4103/phrev.phrev_21_17
- Cakmak, Y.S., Kaya, M., & Asan-Ozusaglam, M. (2014). Biochemical composition and bioactivity screening of various extracts from *Dunaliella salina*, a green microalga. *EXCLI journal*, 13, 679.
- Calabrese, E.J., & Baldwin, L.A. (2000). Chemical hormesis: its historical foundations as a biological hypothesis. *Human & Experimental Toxicology*, 2–31. <https://doi.org/10.1191/096032700678815585>
- Derome, N., Gauthier, J., Boutin, S., & Llewellyn, M. (2016). The Rasputin Effect: When Commensals and Symbionts Become Parasitic (C.J. Hurst, Ed.; Vol. 3). Springer International Publishing. <https://doi.org/10.1007/978-3-319-28170-4>
- Dewi, I.C., Falaise, C., Hellio, C., Bourgougnon, N., & Mouget, J.-L. (2018). Anticancer, Antiviral, Antibacterial, and Antifungal Properties in Microalgae. In *Microalgae in Health and Disease Prevention* (pp. 235–261). Elsevier. <https://doi.org/10.1016/B978-0-12-811405-6.00012-8>
- Duff, D.C.B., Bruce, D.L., & Antia, N.J. (1966). THE ANTIBACTERIAL ACTIVITY OF MARINE PLANKTONIC ALGAE. *Canadian Journal of Microbiology*, 12(5), 877–884. <https://doi.org/10.1139/m66-120>
- Elisabeth, B., Rayen, F., & Behnam, T. (2021). Microalgae culture quality indicators: a review. *Critical Reviews in Biotechnology*, 41(4), 457–473. <https://doi.org/10.1080/07388551.2020.1854672>
- Falaise, C., François, C., Travers, M.-A., Morga, B., Haure, J., Tremblay, R., Turcotte, F., Pasetto, P., Gastineau, R., Hardivillier, Y., Leignel, V., & Mouget, J.-L. (2016). Antimicrobial Compounds from Eukaryotic Microalgae against Human Pathogens and Diseases in Aquaculture. *Marine Drugs*, 14(9), 159. <https://doi.org/10.3390/md14090159>
- FAO. (2020). *The State of World Fisheries and Aquaculture 2020. Sustainability in Action*. <https://doi.org/10.4060/ca9229en>
- Flores-Kossack, C., Montero, R., Köllner, B., & Maisey, K. (2020). Chilean aquaculture and the new challenges: Pathogens, immune response, vaccination and fish diversification. *Fish & Shellfish Immunology*, 98, 52–67. <https://doi.org/10.1016/j.fsi.2019.12.093>
- Fuentes, J., Garbayo, I., Cuaserna, M., Montero, Z., González-del-Valle, M., & Vílchez, C. (2016). Impact of Microalgae-Bacteria Interactions on the Production of Algal Biomass and Associated Compounds. *Marine Drugs*, 14(5), 100. <https://doi.org/10.3390/md14050100>
- Ghosh, C., Sarkar, P., Issa, R., & Haldar, J. (2019). Alternatives to Conventional Antibiotics in the Era of Antimicrobial Resistance. *Trends in Microbiology*, 27(4), 323–338. <https://doi.org/10.1016/j.tim.2018.12.010>
- Gonçalves, C.F., Menegol, T., & Rech, R. (2019). Biochemical composition of green microalgae *Pseudoneochloris marina* grown under different temperature and light conditions. *Biocatalysis and Agricultural Biotechnology*, 18, 101032. <https://doi.org/10.1016/j.bcab.2019.101032>
- Guillard, R.R.L., & Ryther, J.H. (1962). STUDIES OF MARINE PLANKTONIC DIATOMS: I. CYCLOTELLA NANA HUSTEDT, AND DETONULA CONFERVACEA (CLEVE) GRAN. *Canadian Journal of Microbiology*, 8(2), 229–239. <https://doi.org/10.1139/m62-029>
- Hayes, M., Skomedal, H., Skjånes, K., Mazur-Marzec, H., Toruńska-Sitarz, A., Catala, M., Isleten Hosoglu, M., & García-Vaquero, M. (2017). Microalgal proteins for feed,

- food and health. In *Microalgae-Based Biofuels and Bioproducts* (pp. 347–368). Elsevier.
<https://doi.org/10.1016/B978-0-08-101023-5.00015-7>
- Hutchings, M.I., Truman, A.W., & Wilkinson, B. (2019). Antibiotics: past, present and future. *Current Opinion in Microbiology*, 51, 72–80.
<https://doi.org/10.1016/j.mib.2019.10.008>
- Igarashi, M., Ishizaki, Y., & Takahashi, Y. (2018). New antituberculous drugs derived from natural products: current perspectives and issues in antituberculous drug development. *The Journal of Antibiotics*, 71(1), 15–25.
<https://doi.org/10.1038/ja.2017.126>
- Ina-Salwany, M.Y., Al-saari, N., Mohamad, A., Mursidi, F., Mohd-Aris, A., Amal, M.N.A., Kasai, H., Mino, S., Sawabe, T., & Zamri-Saad, M. (2019). Vibriosis in Fish: A Review on Disease Development and Prevention. *Journal of Aquatic Animal Health*, 31(1), 3–22.
<https://doi.org/10.1002/aah.10045>
- Istiqomah, I., Sukardi, Murwantoko, & Isnansetyo, A. (2020). Review Vibriosis Management in Indonesian Marine Fish Farming. *E3S Web of Conferences*, 147, 01001.
<https://doi.org/10.1051/e3sconf/202014701001>
- Jaiswal, K.K., Banerjee, I., Singh, D., Sajwan, P., & Chhetri, V. (2020). Ecological stress stimulus to improve microalgae biofuel generation: a review. *Octa Journal of Biosciences*, 8, 48–54.
- Jena, J., & Subudhi, E. (2019). Microalgae: An Untapped Resource for Natural Antimicrobials. In *The Role of Microalgae in Wastewater Treatment* (pp. 99–114). Springer Singapore. https://doi.org/10.1007/978-981-13-1586-2_8
- Jha, D., Jain, V., Sharma, B., Kant, A., & Garlapati, V.K. (2017). Microalgae-based Pharmaceuticals and Nutraceuticals: An Emerging Field with Immense Market Potential. *ChemBioEng Reviews*, 4(4), 257–272.
<https://doi.org/10.1002/cben.201600023>
- Kapooore, R., Butler, T., Pandhal, J., & Vaidyanathan, S. (2018). Microwave-Assisted Extraction for Microalgae: From Biofuels to Biorefinery. *Biology*, 7(1), 18.
<https://doi.org/10.3390/biology7010018>
- Khomova, T.v., Gusakova, S.D., & Glushenkova, A.I. (1994). Lipids of microalgae. III. Lipids of the biomass of *Monocystis lutheri* grown under the conditions of an accumulation culture. *Chemistry of Natural Compounds*, 30(3), 295–299. <https://doi.org/10.1007/BF00629960>
- Lin, Q., Zhuo, W.-H., Wang, X.-W., Chen, C.-P., Gao, Y.-H., & Liang, J.-R. (2018). Effects of fundamental nutrient stresses on the lipid accumulation profiles in two diatom species *Thalassiosira weissflogii* and *Chaetoceros muelleri*. *Bioprocess and Biosystems Engineering*, 41(8), 1213–1224.
<https://doi.org/10.1007/s00449-018-1950-z>
- Lovio Fragoso, J., Hayano Kanashiro, C., & Lopez Elias, J. (2019). Effect of different phosphorus concentrations on growth and biochemical composition of *Chaetoceros muelleri*. *Latin American Journal of Aquatic Research*, 47(2), 361–366. <https://doi.org/10.3856/vol47-issue2-fulltext-17>
- Little, S.M., Senhorinho, G.N., Saleh, M., Basiliko, N., & Scott, J.A. (2021). Antibacterial compounds in green microalgae from extreme environments: a review. *Algae*, 36(1), 61–72.
<https://doi.org/10.4490/algae.2021.36.3.6>
- Metting, B., & Pyne, J.W. (1986). Biologically active compounds from microalgae. *Enzyme and Microbial Technology*, 8(7), 386–394.
[https://doi.org/10.1016/0141-0229\(86\)90144-4](https://doi.org/10.1016/0141-0229(86)90144-4)
- Mohamad, N., Amal, M.N.A., Yasin, I.S.M., Zamri Saad, M., Nasruddin, N.S., Al-saari, N., Mino, S., & Sawabe, T. (2019). Vibriosis in cultured marine fishes: a review. *Aquaculture*, 512, 734289.
<https://doi.org/10.1016/j.aquaculture.2019.734289>
- Musa, M., Ayoko, G.A., Ward, Rösch, C., Brown, R.J., & Rainey, T.J. (2019). Factors Affecting Microalgae Production for Biofuels and the Potentials of Chemometric Methods in Assessing and Optimizing Productivity. *Cells*, 8(8), 851.
<https://doi.org/10.3390/cells8080851>
- Nájera-Arce, C., Álvarez-Fitz, P., Pérez-Castro, D., Toribio-Jiménez, J., & Castro-Alarcón, N. (2018). Actividad antibacteriana de diatomeas marinas aisladas de Acapulco, Guerrero, México. *Revista de Biología Marina y Oceanografía*, 53(2), 195.
<https://doi.org/10.22370/rbmo.2018.53.2.1293>
- Norton, T.A., Melkonian, M., & Andersen, R.A. (1996). Algal biodiversity. *Phycologia*, 35(4), 308–326.
<https://doi.org/10.2216/i0031-8884-35-4-308.1>
- Nweze, J.A., Mbaaji, F.N., Huang, G., Li, Y., Yang, L., Zhang, Y., Huang, S., Pan, L., & Yang, D. (2020). Antibiotics Development and the Potentials of Marine-Derived Compounds to Stem the Tide of Multidrug-Resistant Pathogenic Bacteria, Fungi, and Protozoa. *Marine Drugs*, 18(3), 145. <https://doi.org/10.3390/md18030145>
- Pina-Pérez, M.C., Rivas, A., Martínez, A., & Rodrigo, D. (2017). Antimicrobial potential of macro and microalgae against pathogenic and spoilage microorganisms in food. *Food Chemistry*, 235, 34–44.
<https://doi.org/10.1016/j.foodchem.2017.05.033>
- Rojas Jirón, S.F., Romero Peralta, J.N., Hernández Rivera, C.I., & Cea Navas, N.E. (2019). Dinámica de microalgas frente a la costa de las Peñitas-Poneloya, Nicaragua. *Revista Universitaria Del Caribe*, 22(1), 114–123.
<https://doi.org/10.5377/ruc.v22i1.8425>
- Ruiz-Güerrecá, D.A., Licea-Navarro, A.F., & Sánchez-Saavedra, M.D.P. (2019). Evaluation of antimycobacterial activity from marine and freshwater microalgae. *Revista de Biología Marina y Oceanografía*, 54(1), 82.
<https://doi.org/10.22370/rbmo.2019.54.1.1544>
- Salcedo, J., & Coello, D. (2018). Dinámica del plancton en la primera milla náutica frente a la provincia de El Oro, Ecuador. *Revista de Biología Tropical*, 66(2), 836.
<https://doi.org/10.15517/rbt.v66i2.33417>
- Sánchez-Camargo, A. del P., Ibáñez, E., Cifuentes, A., & Herrero, M. (2017). Bioactives Obtained from Plants, Seaweeds, Microalgae and Food By-Products Using Pressurized Liquid Extraction and Supercritical Fluid Extraction (pp. 27–51).
<https://doi.org/10.1016/bs.coac.2017.01.001>
- Santos, L., & Ramos, F. (2018). Antimicrobial resistance in aquaculture: Current knowledge and alternatives to tackle the problem. *International Journal of Antimicrobial Agents*, 52(2), 135–143.
<https://doi.org/10.1016/j.ijantimicag.2018.03.010>
- Schuelter, A.R., Kroumov, A.D., Hinterholz, C.L., Fiorini, A., Trigueros, D.E.G., Vendruscolo, E.G., Zahariva, M.M., & Módenes, A.N. (2019). Isolation and identification of new microalgae strains with antibacterial activity on food-borne pathogens. Engineering approach to optimize synthesis of desired metabolites. *Biochemical Engineering Journal*, 144, 28–39.
<https://doi.org/10.1016/j.bej.2019.01.007>

- Spektorova, L.V., Goronkova, O.I., Nosova, L.P., Albitskaya, O.N., & Danilova, G.Yu (1986). High-density culture of marine microalgae — Promising items for mariculture. *Aquaculture*, 55(3), 231–240. [https://doi.org/10.1016/0044-8486\(86\)90118-3](https://doi.org/10.1016/0044-8486(86)90118-3)
- Sudheesh, P.S., & Cain, K. (2017). Prospects and challenges of developing and commercializing immersion vaccines for aquaculture. *Integrative Biology*, 1, 1–20.
- Sydney, E.B., Schafranski, K., Barretti, B.R.V., Sydney, A.C.N., Zimmerman, J.F.D., Cerri, M.L., & Mottin Demiate, I. (2019). Biomolecules from extremophile microalgae: from genetics to bioprocessing of a new candidate for large-scale production. *Process Biochemistry*, 87, 37–44. <https://doi.org/10.1016/j.procbio.2019.09.012>
- Tavakoli, S., Regenstein, J.M., Daneshvar, E., Bhatnagar, A., Luo, Y., & Hong, H. (2021). Recent advances in the application of microalgae and its derivatives for preservation, quality improvement, and shelf-life extension of seafood. *Critical Reviews in Food Science and Nutrition*, 1–14. <https://doi.org/10.1080/10408398.2021.1895065>
- Ture, M., Altinok, I., Cebeci, A., & Caliskan, N. (2022). Characterization of novel bacteriophage HP-T19 that targets *Hafnia alvei*. *Aquaculture Research*, 53(2), 694–699. <https://doi.org/10.1111/are.15592>