



## Heavy Metal Levels in Four Commercial Fishes Caught in Sinop Coasts of the Black Sea, Turkey

Levent Bat<sup>1,\*</sup>, Hasan Can Öztekin<sup>1</sup>, Funda Üstün<sup>1</sup>

<sup>1</sup> Sinop University, Fisheries Faculty, 57000, Sinop, Turkey.

\* Corresponding Author: Tel.: +90.368 2876254/3312; Fax: +90.368 2876268;  
E-mail: leventbat@gmail.com

Received 23 December 2014  
Accepted 22 April 2015

### Abstract

The aim of this study is to evaluate the levels of eight heavy metals (aluminium, arsenic, copper, zinc, mercury, iron, cadmium and lead) in the muscle tissue of Red mullet (*Mullus barbatus*), Whiting (*Merlangius merlangus*), Mediterranean horse mackerel (*Trachurus mediterraneus*) and Golden grey mullet (*Liza aurata*) caught from Sinop coast of the Black Sea during fishing season in 2012 and 2013. Metal analyses in fish were performed using ICP-MS (Inductively Coupled Plasma – Mass Spectrometer). The levels of Al, Hg, Cd, Pb and Cu in all fish species except *T. mediterraneus* for Cu were below the limit of detections. Mean Cu level in Mediterranean horse mackerel was 0.67 mg/kg wet wt. Highest value of Zn (24.7 mg/kg wet wt) was also found in the *T. mediterraneus* which were about 7-8.5 times more than the other species. *L. aurata* have the highest Fe value (3.2 mg/kg wet wt), but have the lowest As value (0.25 mg/kg wet wt.) compared with the other fish species. The levels of As for *M. barbatus* were high (1.3 mg/kg wet wt.) and followed by *M. merlangus* (1.24 mg/kg wet wt). The concentrations of the studied heavy metals were far below the established values by the Turkish Food Codex and Commission Regulation (EC). Therefore, their contribution to the total body burden of these heavy metals can be considered as negligibly small.

**Keywords:** Black Sea, heavy metal, *Mullus barbatus*, *Merlangius merlangus*, *Trachurus mediterraneus*, *Liza aurata*.

### Karadeniz'in Sinop Kıyılarında (Türkiye) Yakalanan Dört Ticari Balıkta Ağır Metal Düzeyleri

#### Özet

Bu çalışmanın amacı 2012-2013 balıkçılık sezonunda Karadeniz'in Sinop kıyılarından yakalanan barbun (*Mullus barbatus*), mezgit (*Merlangius merlangus*), istavrit (*Trachurus mediterraneus*) ve altınbaş kefal (*Liza aurata*) balıklarının kas dokularında sekiz ağır metalin (alüminyum, arsenik, bakır, çinko, cıva, demir, kadmiyum ve kurşun) düzeylerini değerlendirmektir. Balıklardaki metal analizleri ICP-MS (endüktif eşleşmiş plazma – kütle spektrometresi) kullanılarak gerçekleştirilmiştir. *T. mediterraneus* türündeki Cu miktarı hariç tüm balık türlerindeki Al, Hg, Cd, Pb ve Cu düzeyleri ölçüm düzeylerinin altında bulunmuştur. İstavritteki ortalama Cu miktarı 0.67 mg/kg yaş ağırlıktır. En yüksek Zn (24,7 mg/kg yaş ağırlık) yine *T. mediterraneus* türünde bulunmuş olup diğer türlerdeki değerlerden 7-8,5 kez daha fazladır. Diğer türler ile karşılaştırıldığında *L. aurata* türü en yüksek Fe (3,2 mg/kg yaş ağırlık) ancak en düşük As (0,25 mg/kg yaş ağırlık) değerine sahiptir. *M. barbatus* türündeki As düzeyleri yüksek (1,3 mg/kg yaş ağırlık) olup bunu *M. merlangus* türü (1,24 mg/kg yaş ağırlık) izlemiştir. Çalışılan ağır metal konsantrasyonları Türk Gıda Kodeksi ve Avrupa Birliği Tüzüğü'nce belirlenen değerlerin oldukça altında bulunmuştur. Bu nedenle, bu ağır metallerin toplam vücut yükündeki payının önemsenmeyecek kadar küçük olarak nitelendirilebilir.

**Anahtar Kelimeler:** Karadeniz, ağır metal, *Mullus barbatus*, *Merlangius merlangus*, *Trachurus mediterraneus*, *Liza aurata*.

#### Introduction

The pollution of marine environment with chemical contaminants especially heavy metals has become one of the most critical environmental problems of the century. As a result of the heavy metals transport from industrial areas into the

environment and their chemical persistence, many marine ecosystems like the Black Sea are faced with spatially or temporally serious high levels of heavy metals. The Black Sea is one of the youngest seas in the world and it is nearly-enclosed basin between southern Europe and the Anatolian peninsula and has historically been one of the most biologically

productive regions in the world (Özsoy and Ünlüata, 1997). The Black Sea ecosystem has been seriously damaged as a result of pollution (Bat, 2014). In terms of heavy metal, the Black Sea is vulnerable to pollution from untreated industrial effluents and municipal wastewater, runoff from chemical fertilizers in the coastal area and rivers. Bakan *et al.* (1996) pointed out that Ordu and Giresun city centres have separate sewerage systems where Sinop, Trabzon and Zonguldak have combined systems and Samsun city centre have both combined and separate sewerage system. The dumping of wastes via rivers in the Black Sea provides a significant source of metal input (Bakan and Büyükgüngör, 2000; Altas and Büyükgüngör, 2007; Bat *et al.*, 2009).

Some of essential heavy metals are required by humans and quickly decay into harmless or less harmful forms, while non-essentials are non-biodegradable and remain dangerous for a long time. Now, there is a growing concern worldwide over the indiscriminate use of such heavy metals, resulting in environmental pollution and toxicity risk to marine organisms. Thus, ecological damage of the environment caused by anthropogenic factors as well as the presence of heavy metals may affect people. Heavy metals tend to accumulate in advanced organisms through bio-magnification effects in the food chain. Thus they can enter into the human body and accumulate in the human tissues to pose chronic toxicity. Chronic assimilation of heavy metals is a known cause of cancer (The Earth Report 3, 1992).

The Marine Strategy Framework Directive (MSFD, 2008/56/EC) establishes a framework for the development of marine strategies designed to achieve Good Ecological Status (GES) in the marine environment, by the year 2020, using 11 qualitative descriptors (Official Journal of the European Union, 2008). The concentration of contaminants including heavy metals in the marine environment and their effects needs to be assessed taking into account the impacts and threats to the ecosystem in Directive 2008/56/EC. Contaminants in fish and other seafood for human consumption do not exceed levels established by Community legislation or other relevant standards (Descriptor 9; Directive 2000/60/EC) (Official Journal of the European Communities, 2000).

Fish is a healthy food because of its nutritional benefits related to its proteins of high biological quality, desirable lipid composition, valuable mineral compounds and vitamins. The particular composition of its lipid fraction, rich in essential  $\omega$ -3 polyunsaturated fatty acids (PUFA), especially eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), and low cholesterol levels makes it an important item in human food. Thus, it is very important to determine the heavy metals concentrations in commercial fish in order to evaluate the possible risk of fish consumption for human health.

Total fish production in Turkey was reported as 644.852 tons in 2012 and 607.515 tons in 2013, where 315.636,5 tons in 2012 and 295.167,9 tons in 2013 of this total production came from sea fish (TUIK, 2014). Catches of Red mullet, Whiting, Mediterranean horse mackerel and Golden grey mullet from Turkish waters in 2013 were 2.055,4 tons, 9.396,9 tons, 21.817,8 tons and 2.504,9 tons, respectively (TUIK, 2014). These data show that these species of fish are economically important for human consumption and there is a need for more information on contaminant levels in fish of Turkey. To protect and develop fishery resources, it is important to undertake research work on this environmental issue.

The present study provides information on eight heavy metal concentrations in dorsal muscle tissue of four commercial fish species (*M. barbatus*, *M. merlangus*, *T. mediterraneus* and *L. aurata*) from the Sinop coast of the Black Sea, Turkey during fishing season (i.e. autumn, winter) in 2012 and 2013. The heavy metals investigated include aluminium, arsenic, copper, zinc, mercury, iron, cadmium and lead because they are known to accumulate in the environment and in fish, and are known to cause adverse health effects if consumed in sufficient quantities.

The aims of this study are to determine and to compare the concentrations of aluminium, arsenic, copper, zinc, mercury, iron, cadmium and lead in edible parts of four selected commercial Sinop coast of the Black Sea fish species and to compare with the guidelines set down by the Ministry of Agriculture, Fisheries and Food (MAFF), the Turkish Food Codex, Commission Regulation (EC) for the safe consumption limits of fish and the other studies.

## Materials and Methods

### Sampling

The fish species selected were; Red mullet (*M. barbatus*), Whiting (*M. merlangus*), Mediterranean horse mackerel (*T. mediterraneus*) and Golden grey mullet (*L. aurata*). Red mullet, Whiting and Mediterranean horse mackerel were sampled during fish season in 2012; Golden grey mullet were sampled during fish season in 2012 and 2013 directly from the Turkish fishing vessels in Sinop coasts of the Black Sea. The total length was measured from snout to the tip of the caudal fin and the body weight (g) was taken on a top loading electronic balance to the nearest 0.1 g.

### Study Area

To understand the bioavailability of heavy metals and assess their potential impact on marine biota, fish samples were obtained during the fishing period of 2012 and 2103. The study area is presented

in Figure 1.

### Preparation Samples and Determination of Heavy Metals


Twenty fish individuals from each species taken randomly were rinsed in uncontaminated sea water then were washed with double distilled water. The muscle tissues of the fish were prepared for analysis according to the method described by Bernhard (1976). All samples were stored deep frozen at  $-21^{\circ}\text{C}$  until their analysis. Metal analysis in fish was performed using m-AOAC 999.10- ICP/MS (Inductively Coupled Plasma – Mass Spectrometer) method by accredited ÇEVRE Industrial Analysis Laboratory Services Trade Company (TÜRKAK Test TS EN ISO IEC 17025 AB-0364-T). EN 15763 European Standard methods was applied. The limits of detection used for analysis of aluminium, arsenic, copper, zinc, mercury, iron, cadmium and lead were 0.5, 0.05, 0.5, 0.5, 0.05, 0.5, 0.02 and 0.05, respectively.

### Statistical Analysis

A one- way analysis of variance (ANOVA) was performed, followed by Tukey post hoc comparisons for the source of statistically significant difference. The significance was set at 0.05 and P-values less of 0.05 were considered statistically significant (Zar, 1984). All values were being expressed on an mg/kg wet wt. basis.

### Results

The mean lengths (cm) and weights (g) of the fish samples are given in Table 1. The analytical data for these metals also showed that there was difference in heavy metal content in various fish. Figure 2 shows the mean concentrations of eight heavy metals in the muscle of fish species (*M. barbatus*, *M. merlangus*, *T. mediterraneus* and *L. aurata*) from the Sinop coast of the Black Sea. The levels of Al, Hg, Cd and Pb in all fish species were below the limit of detection. The levels of Cu in Red mullet, Whiting and Golden grey



**Figure 1.** Fish sampling area from Sinop coasts of the Black Sea, Turkey.

**Table 1.** The mean and standard deviations of total length (cm) and total weight (g) of fish species from Sinop coasts of the Black Sea

Fish species	Mean $\pm$ SD Lengths (cm)	Mean $\pm$ SD Weights (g)
<i>Mullus barbatus</i>	13 $\pm$ 3	25 $\pm$ 9
<i>Merlangius merlangus</i>	12 $\pm$ 2	21 $\pm$ 5
<i>Trachurus mediterraneus</i>	11 $\pm$ 1	17 $\pm$ 3
<i>Liza aurata</i>	25 $\pm$ 4	185 $\pm$ 14

**Table 2.** Metal concentrations (mean mg/kg wet wt.) in the muscle tissues of the fish species from the Black Sea. Vertically, letters a, b and c show statistically significant differences ( $P < 0.05$ )

Species	Heavy metals							
	Cd	Hg	Pb	Al	Cu	As	Fe	Zn
<i>Mullus barbatus</i>	<0.02 <sup>a</sup>	<0.05 <sup>a</sup>	<0.05 <sup>a</sup>	<0.5 <sup>a</sup>	<0.5 <sup>a</sup>	1.30 <sup>a</sup>	2.3 <sup>a</sup>	3.2 <sup>a</sup>
<i>Merlangius merlangus</i>	<0.02 <sup>a</sup>	<0.05 <sup>a</sup>	<0.05 <sup>a</sup>	<0.5 <sup>a</sup>	<0.5 <sup>a</sup>	1.24 <sup>a</sup>	0.87 <sup>b</sup>	3.4 <sup>a</sup>
<i>Liza aurata</i>	<0.02 <sup>a</sup>	<0.05 <sup>a</sup>	<0.05 <sup>a</sup>	<0.5 <sup>a</sup>	<0.5 <sup>a</sup>	0.25 <sup>b</sup>	3.2 <sup>c</sup>	2.9 <sup>b</sup>
<i>Trachurus mediterraneus</i>	<0.02 <sup>a</sup>	<0.05 <sup>a</sup>	<0.05 <sup>a</sup>	<0.5 <sup>a</sup>	0.67 <sup>b</sup>	0.39 <sup>c</sup>	2.2 <sup>a</sup>	24.7 <sup>c</sup>

mullet were also below the limit of detection. However, Cu level in Mediterranean horse mackerel was 0.67 mg/kg. In general all the species Zn is the most abundant of the metals examined. Highest values were found in the *T. mediterraneus* which were about 7-8.5 times more than the other species. *L. aurata* have the highest Fe values (3.2 mg/kg), but have the lowest As values (0.25 mg/kg) compared with the other fish species. The levels of As for *M. barbatus* were high (1.3 mg/kg) and followed by *M. merlangus* (1.24 mg/kg). Metal concentrations in the different fish species by ANOVA and Tukey test are shown in Table 2. The results indicated that there were significant differences within fish species ( $P < 0.05$ ).

## Discussion

Recently, Bat (2014) reviewed the heavy metals in biota from the Black Sea coast of Turkey and suggested that fish can be considered good bioindicators for heavy metal contamination in marine ecosystems because they occupy different trophic levels and are widely consumed in many parts of the world by humans. Heavy metals whether essential or not may be toxic to living organisms including fish and this can lead to deleterious effects on fisheries and, potentially, public health. In this study, the metal concentrations decreased in the order  $Zn > Fe > As > Cu > Al > Pb > Hg > Cd$ . Zn and Cu accumulated in higher concentrations in *T. mediterraneus* than another species studied. The highest Fe level was in *L. aurata*. Similar results were found by other studies (Tepe et al., 2007; Türkmen et al., 2008, 2009). Differences in metal concentrations were related to diet and feeding habits of benthic and pelagic fish species (Bustamente et al., 2003). They show that benthic fish generally accumulate higher concentrations of heavy metals than pelagic fish. Topping (1973) suggested that mainly plankton feeding fish contain much higher concentrations of some heavy metals than bottom feeding fish. Metals, such as Fe, Cu and Zn, are essential metals since they play important roles in biological systems (Türkmen et al., 2009) and the contaminant uptake rate is positively linked to the metabolic rate in marine organisms, it can be supposed that metal accumulation would be high (Kojadinovic et al., 2007). Among the four species, the highest As was found in *M. barbatus* followed by *M. merlangus*. The red mullet is a carnivorous species, found mostly on muddy bottoms and feed on Polychaeta, Lamellibranchiata and Crustacea. Whiting is benthopelagic and is also a carnivorous species, found mainly on mud and gravel bottoms, but also on sand and rock. They feed on shrimps, crabs, molluscs, small fish and polychaetes. Kojadinovic et al. (2007) suggested that that fish muscle impregnation is the result of the accumulation of heavy metals from preys living in different environments than where the fish

were caught and which contained high metal levels. The remaining metals including Al, Pb, Hg and Cd were below the limit of detection in all fish samples. Hg, Pb and Cd belong to the group of non-essential and toxic metals and they have no known function in biochemical processes and are accumulated in multiple organs (Phillips, 1977).

Studies on the distribution of heavy metals show that there exists a certain variation in accumulation patterns among fish species. The metal concentrations in these fish species were compared among the Turkish Black Sea and other coasts and presented in Table 3. The maximum concentrations of Pb ( $6.86 \pm 0.26$  mg/kg dry wt.) and Cu ( $9.10 \pm 5.9$  mg/kg dry wt.) were found to be highest in *M. barbatus* of Trabzon coasts (Topçuoğlu et al., 1990). Fe ( $163 \pm 12$  mg/kg dry wt.) and Zn ( $106 \pm 9.1$  mg/kg dry wt.) concentrations were found to be the highest in *M. barbatus* (Uluozlu et al., 2007). The highest As level ( $2.375$  mg/kg dry wt.) was also found in *M. barbatus* of Sinop coast (Das et al., 2009). The maximum concentrations of Al and Hg were  $86.3 \pm 11$  mg/kg dry wt. (Turan et al., 2009) and  $0.084 \pm 0.005$  mg/kg wet wt. (Tüzen, 2009) in *M. merlangus*, respectively. Cd concentrations were found to be highest ( $0.81$ - $1.61$  mg/kg dry wt.) in *L. aurata* from Egypt and it was followed by *M. merlangus* with  $0.55 \pm 0.04$  mg/kg dry wt. (Türkmen et al., 2008). Boran and Altinok (2010) concluded that heavy metal pollution in living organisms of the Black Sea has attracted considerable research attention since last 20 years.

In this study the levels of Al, Hg, Cd and Pb in all fish species were below the limit of detection. However, the average concentrations of other metals were slight lower than the maximum tolerance levels for human consumption established by the Turkish Food Codex, Commission Regulation (EC) and MAFF (TGK, 2002; Council of Europe, 2001; MAFF, 1995). Furthermore, the tolerable weekly intakes were estimated by means of references for muscles tissues of fishes consumed by people. The average yearly fish consumption in Turkey is 7.1 kg per person (TUIK, 2014), which are equivalent to 19.45 g/day for Turkey.

The highest Cu level in Mediterranean horse mackerel was 0.67 mg/kg. The Joint FAO/WHO Expert Committee on Food Additives established a PTWI for Cu of 3.5 mg/kg body weight/week which was equivalent to 245 mg/week for a 70 kg adult (WHO, 1996; Council of Europe, 2001). By using the means of weekly fish consumption in Turkey of 136.15 g per person and intake of Cu in fish was calculated as 0.09 mg ( $136.15 \text{ g} \times 0.67 \text{ mg}/1000 \text{ g}$ ) per person. *L. aurata* have the highest Fe values (3.2 mg/kg), the Joint FAO/WHO Expert Committee on Food Additives established a PTWI for Fe of 5.6 mg/kg body weight/week which was equivalent to 392 mg/week for a 70 kg adult (FAO/WHO, 2010). Intake of Fe in fish was calculated as 0.44 mg ( $136.15 \text{ g} \times 3.2 \text{ mg}/1000 \text{ g}$ ) per person. Highest values of Zn

**Table 3.** Heavy metal levels in fish from the Black Sea coasts and other country expressed in mg/kg wet wt. (\*= dry wt., BS= Black Sea)

Species	Region	Heavy metals								References
		Cd	Hg	Pb	Al	Cu	As	Fe	Zn	
<i>M. barbatus</i>	Sinop	<0.02	<0.05	<0.05	<0.5	<0.5	1.30	2.3	3.2	Present study
<i>M. barbatus</i> *	Trabzon	<0.1	-	6.86±0.26	-	9.10±5.9	-	39±1	11.5±3.5	Topçuoğlu et al., 1990
<i>M. barbatus</i>	Sinop	0.023±0.002	-	0.28±0.06	-	0.76±0.07	-	4.18±0.81	2.42±0.27	Bat et al., 1996
<i>M. barbatus</i>	Sinop	0.076	-	0.424	-	8.968	-	21.33	9.90	Bat et al., 2006
<i>M. barbatus</i> *	BS	0.017	-	0.077	-	0.01	-	4.5	4.3	Dalman et al., 2006
<i>M. barbatus</i> *	Sinop	-	-	-	-	0.38-2.714	-	-	1.424-63.29	Turk Culha et al., 2007
<i>M. barbatus</i> *	BS	0.45±0.04	-	0.84±0.07	-	0.98±0.07	-	163±12	106±9.1	Uluozlu et al., 2007
<i>M. barbatus</i> *	BS	0.208±0.017	-	0.727±0.141	9.85±14	-	-	21.2 ±1.47	7.573±0.389	Turan et al., 2009
<i>M. barbatus</i>	BS	0.17 ± 0.02	0.036±0.002	0.36 ± 0.03	-	0.96±0.08	0.11±0.01	53.2 ± 2.8	75.5 ± 5.3	Tuzen, 2009
<i>M. barbatus</i> *	Samsun	<0.02	<0.050	0.0815±0.003	-	-	1.33±0.02	-	-	Das et al., 2009
<i>M. barbatus</i> *	Sinop	<0.02	<0.050	0.051±0.0005	-	-	2.375	-	-	Das et al., 2009
<i>M. barbatus</i> *	BS	0.020±0.002	low	0.92±0.12	-	3.14±0.31	-	29.17±2.18	23.71±0.71	Nisbet et al., 2010
<i>M. barbatus</i> *	Sinop	0.08-0.18	-	0.02	-	2.85-5.26	-	-	22.82-34.33	Bat et al., 2013
<i>M. merlangus</i>	Sinop	<0.02	<0.05	<0.05	<0.5	<0.5	1.24	0.87	3.4	Present study
<i>M. merlangus</i>	Kastamonu	-	0.01-0.22	0.02-0.11	-	0.62-3.25	-	-	-	Ünsal et al., 1993
<i>M. merlangus</i>	Zonguldak	-	0.006-0.04	0.05-2.26	-	0.37-7.72	-	-	-	Ünsal et al., 1993
<i>M. merlangus</i> *	Perşembe	<0.02	-	<0.05	-	1.86±0.04	-	57±1	43.1±0.1	Topçuoğlu et al., 2002
<i>M. merlangus</i> *	Rize	<0.02	-	<0.05	-	4.54±0.11	-	46±1	30.2±0.1	Topçuoğlu et al., 2002
<i>M. merlangus</i> *	BS	0.0131	-	0.088	-	1.3	-	2.5	3.3	Dalman et al., 2006
<i>M. merlangus</i> *	BS	0.55 ± 0.04	-	0.93 ± 0.07	-	1.25±0.10	-	104±9.8	48.6 ± 3.9	Türkmen et al., 2008
<i>M. merlangus</i> *	BS	0.192 ±0.02	-	0.502 ± 0.104	86.3±11	-	-	4.48 ±0.441	6.03 ± 0.54	Turan et al., 2009
<i>M. merlangus</i>	BS	0.21 ± 0.02	0.084±0.005	0.53 ± 0.04	-	1.32±0.11	0.17±0.01	98.1±7.7	65.4 ± 4.2	Tuzen, 2009
<i>M. merlangus</i> *	BS	0.18 ± 0.02	-	0.46 ± 0.05	-	1.8 ± 0.2	-	27.7±2.8	20.6 ± 2.1	Mendil et al., 2010
<i>L. aurata</i>	Sinop	<0.02	<0.05	<0.05	<0.5	<0.5	0.25	3.2	2.9	Present study
<i>L. aurata</i> *	Egypt	0.81-1.61	-	1.41-2.66	-	3.43-5.49	-	-	12.98-35.45	Bahnasawy et al.,
<i>L. aurata</i> *	Caspian Sea	0.35±0.23	-	1.50±0.53	-	4.54±1.07	-	67.52±33.53	13.69±7.23	Taghavi Jelodar et al., 2011
<i>T. mediterraneus</i>	Sinop	0.03±0.00	-	0.63±0.25	-	16.7±2.08	-	57.2±17.9	42.6±9.14	Türkmen et al., 2008
<i>T. mediterraneus</i>	Bartın	0.30±0.04	-	1.31±0.34	-	1.68±0.11	-	57.6±3.63	8.15±1.81	Türkmen et al., 2008
<i>T. mediterraneus</i>	Sinop	0.043-0.048	-	0.17-0.23	-	2.22-6.21	-	-	17.89-32.38	Bat et al., 2012
<i>T. mediterraneus</i>	Sinop	<0.02	<0.05	<0.05	<0.5	0.67	0.39	2.2	24.7	Present study

were found in the *T. mediterraneus* (27.4 mg/kg wet wt.) which were about 7-8.5 times more than the other species. The Joint FAO/WHO Expert Committee on Food Additives established a PTWI for Zn of 7 mg/kg body weight/week which was equivalent to 490 mg/week for a 70 kg adult (WHO, 1996; Council of Europe, 2001). Intake of Zn in fish was calculated as 3.36 mg (136.15 g x 24.7 mg/1000 g) per person. The highest levels of As were found in *M. barbatus* (1.3 mg/kg), but there is no available data on the As levels in fish to compare. However, the EU Scientific Committee for Food reports indicated that large consumption of fish had as intakes of about 0.050 mg/kg body weight, which corresponds to a daily intake of 3.5 mg for a 70 kg adult. Moreover, it is allowed that fish and seafood consumers may reach an intake of 1 mg/day from these foods alone (Directorate-General Health and Consumer Protection, 2004). By using the means of weekly intake calculated as 0.177 mg (136.15g x 1.3 mg/1000 g) per person for As in muscles of the fishes. As it can be seen that, the concentration of metals in the muscle tissues were generally low and within the ranges expected for the metals in muscle tissues of fish from Sinop coasts of the Black Sea. The concentrations of these metals found in the examined fish species were well below the acceptable levels for human consumption recommended by the Turkish Food Codex and Commission Regulation (EC) which means that they are not toxic to the health of consumers.

## Conclusion

Based on the results, heavy metal concentrations found in the edible tissues of the fish samples were considerably lower than the maximum levels set by the Ministry of Agriculture, Fisheries and Food (MAFF), the Turkish Food Codex, Commission Regulation (EC). Even though, there is no risk from toxic heavy metals in fish, a possible hazard may develop in the future depending on the industrial, agricultural, urban and fishing development in the Black Sea coasts. These data provide a useful baseline to measure any future changes in metal pollution of the Black Sea coasts. Furthermore, constant monitoring of the marine ecosystems in the Black Sea is required by the concerned agency due to increase in anthropogenic activities for safe supply of fish to the consumers.

## Acknowledgments

This work was supported in the Sinop University for Scientific Research Project; project number SÜF-1901-12-02.

## References

- Altas, L. and Büyükgüngör, H. 2007. Heavy metal pollution in the Black Sea shore and offshore of Turkey. Environmental Geology, 52: 469-476. doi: 10.1007/s00254-006-0480-1.

- Bahnasawy, M., Khidr, A.Z. and Dheena, N. 2011. Assessment of heavy metal concentrations in water, plankton and fish of Lake Manzala, Egypt. Turk. J. Zool., 35(2): 271-280. doi:10.3906/zoo-0810-6.
- Bakan, G., Özkoç, H.B., Büyükgüngör, H., Ergun, O.N. and Onar, N. 1996. Evaluation of the Black Sea and-based sources inventory results of the coastal region of Turkey. Proc. of the International Workshop on MED and Black Sea ICZM; 1996 November 2-5; pp: 39-52.
- Bakan, G. and Büyükgüngör, H. 2000. The Black Sea. Marine Pollution Bulletin, 41: 24-43. doi: 10.1016/S0025-326X(00)00100-4.
- Bat, L., Öztürk, M. and Öztürk, M. 1996. Heavy metal amounts in some commercial teleost fish from the Black Sea. O.M.Ü. Fen Dergisi, 7: 117-135.
- Bat, L., Gündoğdu, A., Yardım, Ö., Zoral, T. and Çulha, S. 2006. Heavy metal amounts in zooplankton and some commercial teleost fish from inner harbour of Sinop, Black Sea. SÜMDER (Su Ürünleri Müh. Dergisi), 25: 22-27.
- Bat, L., Gökkurt, O., Sezgin, M., Üstün, F. and Şahin, F. 2009. Evaluation of the Black Sea land based sources of pollution the coastal region of Turkey. The Open Marine Biology Journal, 3: 112-124.
- Bat, L., Sezgin, M., Üstün, F. and Şahin, F. 2012. Heavy metal concentrations in ten species of fishes caught in Sinop coastal waters of the Black Sea, Turkey. Turkish Journal of Fisheries and Aquatic Sciences, 12: 371-376. doi: 10.4194/1303-2712-v12\_2\_24
- Bat, L., Sezgin, M., Gökkurt Baki, O., Üstün, F. and Şahin, F. 2013. Determination of heavy metals in some commercial fish from the Black Sea coast of Turkey. Walailak J. Sci. and Tech., 10: 581-589.
- Bat, L. 2014. Heavy metal pollution in the Black Sea. In: E. Düzgüneş, B. Öztürk, and M. Zengin (Eds.), Turkish Fisheries in the Black Sea. Published by Turkish Marine Research Foundation (TUDAV), Publication number: 40, Istanbul: 71-107.
- Boran, M. and Altinok, I. 2010. A review of heavy metals in water, sediment and living organisms in the Black Sea. Turkish Journal of Fisheries and Aquatic Sciences, 10(4): 565-572. doi: 10.4194/trjfas.2010.0418
- Bryan, G.W. 1976. Heavy Metal Contamination in the Sea. In: R. Johnston (Ed.), Marine Pollution, Academic Press Inc., London: 185-302.
- Bustamante, P., Bocher, P., Chérel, Y., Miramand, P. and Caurant, A. 2003. Distribution of trace elements in the tissues of benthic and pelagic fish from the Kerguelen Islands. The Science of the Total Environment 313: 25-39. doi:10.1016/S0048-9697(03)00265-1
- Council of Europe 2001. Council of Europe's policy statements concerning materials and articles intended to come into contact with foodstuffs. Policy Statement concerning materials and alloys. Technical Document. Guidelines on metals and alloys used as food contact materials (09.03.2001), Strasbourg, 67 pp.
- Dalman, Ö., Demirak, A. and Balcı, A. 2006. Determination of heavy metals (Cd, Pb) and trace element (Cu, Zn) in sediment and fish of the Southeastern Aegean Sea (Turkey) by Atomic Absorption Spectrometry. Food Chemistry, 95: 157-62. doi: 10.1016/j.foodchem.2005.02.009
- Das, Y.K., Aksoy, A., Baskaya, R., Duyar, H.A., Güvenc, D. and Boz, V. 2009. Heavy metal levels of some

- marine organisms collected in Samsun and Sinop Coasts of Black Sea, in Turkey. *Journal of Animal and Veterinary Advances*, 8(3): 496-499.
- Directorate-General Health and Consumer Protection 2004. Reports on tasks for scientific cooperation. Assessment of the dietary exposure to arsenic, cadmium, lead and mercury of the population of the EU Member States. Report of experts participating in Task 3.2.11. 125 pp. [http://ec.europa.eu/food/food/chemicalsafety/contaminants/scoop\\_3-2-11\\_heavy\\_metals\\_report\\_en.pdf](http://ec.europa.eu/food/food/chemicalsafety/contaminants/scoop_3-2-11_heavy_metals_report_en.pdf). Accessed on 06.06.2014.
- FAO/WHO 2010. Summary report of the seventy-third meeting of JECFA. Joint FAO/WHO Expert Committee on Food Additives. Geneva.
- Kojadinovic, J., Potier, M., Le Corre, M., Cosson, R.P. and Bustamante, P. 2007. Bioaccumulation of trace elements in pelagic fish from the Western Indian Ocean *Environmental Pollution*, 146(2): 548-566. doi: 10.1016/j.envpol.2006.07.015.
- MAFF (The Ministry of Agriculture, Forestry and Fisheries) 1995. Monitoring and surveillance of non-radioactive contaminants in the aquatic environment and activities regulating the disposal wastes at sea. Directorate of Fisheries research, Lowestoft, Aquatic Environment Monitoring Report, No 44.
- Mendil, D., Demirci, Z., Tüzen, M. and Soylak, M. 2010. Seasonal investigation of trace element contents in commercially valuable fish species From the Black Sea, Turkey. *Food and Chemical Toxicology*, 48: 865-870. doi:10.1016/j.fct.2009.12.023
- Nisbet, C., Terzi, G., Pilger, O. and Sarac, N. 2010. Determination of heavy metal levels in fish sample collected from the Middle Black Sea. *Kafkas Üniv.Vet. Fak. Der.*, 16: 119-125.
- Official Journal of the European Communities 2000. Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy, 327: 1-72.
- Official Journal of the European Union 2008. Directives Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008 establishing a framework for community action in the field of marine environmental policy (Marine Strategy Framework Directive). L 164: 19-40.
- Özsoy, E. and Ünlüata, Ü. 1997. Oceanography of the Black Sea: A review of some recent results. *Earth Sci. Rev.*, 42(4): 231-272.
- Phillips, D.J.H. 1977. The use of biological indicator organisms to monitor trace metal pollution in marine and estuarine environments. A review. *Environ. Pollut.*, 13: 281-317.
- Taghavi Jelodar, H., Sharifzadeh Baei, M., Najafpour, Sh. and Fazli, H. 2011. The comparison of heavy metals concentrations in different organs of *Liza aurata* inhabiting in Southern Part of Caspian Sea. *World Applied Sciences Journal (Special Issue of Food and Environment)*, 14: 96-100.
- Tepe, Y., Türkmen, M. and Türkmen, A. 2007. Assessment of heavy metals in two commercial fish species of four Turkish seas. *Environmental Monitoring and Assessment*.doi: 10.1007/s10661-007-0079-3.
- TGK 2002. Official Gazette of Republic of Turkey. Notifications about determination of the maximum levels for certain contaminants in foodstuffs of Turkish Food Codex (in Turkish). (Notification No: 2002/63), Issue: 24885.
- The Earth Report 3. 1992. An A-Z Guide to environmental issues. (General Eds.) Goldsmith, E. and Hildyard, N.) London, 175 pp.
- Topçuoğlu, S., Erentürk, N., Saygı, N., Kut, D., Esen, N., Başsarı, A. and Seddigh, E. 1990. Trace metal levels of fish from the Marmara and Black Sea. *Toxicological and Environ. Chemistry*, 29: 95-99.
- Topçuoğlu, S., Kirbasoglu, Ç. and Güngör, N. 2002. Heavy metals in organisms and sediments from Turkish coast of the Black Sea, 1997-1998. *Environment International*, 27: 521-526. PII: S0160-4120(01)00099-X.
- Topping, G. 1973. Heavy metals in fish from Scottish waters. *Aquaculture*, 11: 373-377.
- TUIK, Turkish Fishery Statistics. 2014. <http://www.tuik.gov.tr/>. Accessed 9.12.2014.
- Turan, C., Dural, M., Oksuz, A. and Öztürk, B. 2009. Levels of heavy metals in some commercial fish species captured from the Black Sea and Mediterranean coast of Turkey. *Bull Environ. Contam. Toxicol.*, 82: 601-604. doi:10.1007/s00128-008-9624-1
- Türk Çulha, S., Bat, L., Çulha, M., Efendioglu, A. andaç, M. and Bati, B. 2007. Heavy metals levels in some fishes and molluscs from Sinop Peninsula of the Southern Black Sea, Turkey. *Rapp. Comm. Int. Mer. Medit.*, 38: 323.
- Türkmen, M., Türkmen, A. and Tepe, Y. 2008. Metal contaminations in five fish species from Black, Marmara, Aegean and Mediterranean Seas, Turkey. *J. Chil. Chem. Soc.*, 53(1): 1435-1439. Doi:10.4067/S0717-97072008000100021.
- Türkmen, M., Türkmen, A., Tepe, Y., Töre, Y. and Ateş, A. 2009. Determination of metals in fish species from Aegean and Mediterranean seas. *Food Chemistry*, 113: 233-237. doi:10.1016/j.foodchem.2008.06.071.
- Tüzen, M. 2009. Toxic and essential trace elemental contents in fish species from the Black Sea, Turkey. *Food and Chemical Toxicology*, 47:1785-1790. doi:10.1016/j.fct.2009.04.029
- Uluozlu, O.D., Tüzen, M., Mendil, D. and Soylak, M. 2007. Trace metal content in nine species of fish from the Black and Aegean Seas, Turkey. *Food Chemistry*, 104(2): 835-840. doi:10.1016/j.foodchem.2007.01.03
- Ünsal, M., Bekiroğlu, Y., Akdoğan, Ş, Ataç, Ü., Kayıkçı, Y., Alemdağ, N., Aktaş, M. and Yıldırım C. 1993. Determination of heavy metals in some economically important marine organisms in southwestern Black Sea. TUBITAK Project No: DEBAG-80/G, Ankara, 78 pp. (in Turkish).
- WHO 1996. Trace elements in human nutrition and health. ISBN 92 4 156173 4 (NLM Classification: QU 130). Geneva.
- Zar, J.H. 1984. *Biostatistical analysis*. Prentice Hall, Int., New Jersey, 718 pp.